

The Newsletter of the International Fission-Track Community June 2006, Volume 14, Number 1, Issue 27

# CONTENTS

Editor's Notes	2
SHORT TRACKS	2
FORTHCOMING MEETINGS	3
PAST MEETINGS	10
CURRENT PHD STUDIES	13
Perturbation of isotherms below topography: constraints from tunnel transects through the Alps .	13
Low Temperature Thermochronology of the Krishna-Godavari Basin, India and its Hinterland	13
Neogene Extensional Deformation in Northern Baja California, Mexico: Implications from Combir Low-Temperature Thermochronology and Structural Studies	ned 13
Modelling the architecture of coupled denudation-sedimentation systems in Australian sedimenta basin evolution	ary 14
New Approaches to Low-Temperature Thermochronology	14
The post break-up margin evolution of northern Namibia (Kaokoveld)	15
Low-temperature thermochronology of the northern Canadian shield, and new insights into fissio track behaviour in solids	on 15
RECENT PHD THESES	16
Calibration of the fission-track method and application in central China	16
The low-temperature thermochronology of cratonic terranes	17
Cooling history and relief evolution of Corsica (France) as constrained by fission track and (U- Th)/He thermochronology	18
ARTICLES	20
Measuring wt% CI to assess and allow for variation in fission track annealing rates between different apatite species	20
Call for interlaboratory comparison of U & Th measurements in sub mg apatite samples	30
Do I need to average all my zetas? — Outlier test by Out?Lier	31
New Features. Tricks and Tips for the use of Trackkey	
Irradiation Eacilities at the Forschungsneutronenguelle Heinz Maier Leibnitz (FRM II)	41
INTERNATIONAL FISSION-TRACK DIRECTORY	
CALL FOR CONTRIBUTIONS	72

On Track is a newsletter of the international fission-track community published electronically in the months of July and February. The views expressed in On Track articles are those of the authors and do not reflect those of the fission-track community or the editor of On Track. The copyright of articles belongs to the authors. Articles may not be reproduced without written permission. Trademarks and patents are the property of the corporations and/or individuals indicated in the article.

#### PREVIOUS EDITORS:

Dave Coyle (1990-91), La Trobe University Trevor Dumitru (1991-92), Stanford University Rasoul Sorkhabi (1992-93), Arizona State University Dennis Arne (1993-94), Dalhousie University Richard J. Weiland (1994-95), University of Texas Ruth Siddall (1995-96), University College London Danny Stockli (1996-97), Stanford University Paul O'Sullivan (1997-98), La Trobe University Alexander Grist (1998-99), Dalhousie University Richard Spikings (1999-2000), ETH Zürich Raymond Jonckheere (2000-2001) Max-Plank Heidelberg Andy Carter (2001-2002) Birkbeck College, London Ed Sobel (2002-2003) Inst. f. Geow., Potsdam Roderick Brown (2003-2006) Glasgow University

CURRENT EDITOR: Matthias Raab The University of Melbourne School of Earth Sciences Melbourne, 3010, Australia Phone: + 61 3 8344 9868 Fax: + 61 3 8344 7761 E-mail: mraab@unimelb.edu.au

# **Editor's Notes**

I am delighted to announce that OnTrack is back on track again.

Shortly after the 10th International Conference on Fission Track Dating and Thermochronology in Amsterdam in 2004, four manuscripts were submitted to OnTrack. Big thanks to the authors for their contributions and for their patience in particular.

Because of the forthcoming European Conference on Thermochronology (ECTC) in Bremen, Germany (30/7 - 4/8), I was trying to get this issue out as soon as the program had been finalised by the ECTC-committee. Despite the very short notice, my call for contributions in early May received a magnificent response from many people in the fission track community. Comprising 74 pages, this seems to be the largest OnTrack ever.

In this issue you will find details for the forthcoming ECTC and the Goldschmidt conferences, directly followed by nostalgic images from our past meeting in Amsterdam. OnTrack

always encourages PhD students to tell us about their research. Commencing on page 13 seven students give us insight into *their* exciting projects, and most of them are in the first half of their candidature. I know that there are many more students in the community, and hopefully you will see the seven abstracts as an encouragement to write your own two paragraph project description for the next issue. From my personal point of view PhD students should start very early in their career to become visible in the scientific community. OnTrack is a great platform to start with, followed by conferences, workshops, etc. The better you are known for your research the better are your chances for a smooth transition into a post-doctoral position in a different research group. I encourage every student to use this opportunity and publish early versions of your work, and hope that supervisors encourage their students too. You will see it is a very beneficial exercise.

Three recent PhD theses give an overview of their base and applied research, demonstrating that basic assumptions should always be questioned.

In the article section Paul Green responded immediately to a heated (and ongoing) discussion in Amsterdam about the variability of track annealing behaviour in apatite, and outlines recommendations to integrate chlorine measurements. István Dunkl sent three articles, where he: a) invites researchers to participate in an interlaboratory comparison of U and Th measurements on gem-quality apatites; b) introduces his new program to test for outliers in datasets, and c) provides new features, tricks and tips for the use of Trackkey, a program to calculate fission track ages. A new research reactor in Germany commenced operations in 2004. Heiko Gerstenberg gives us great insight into the irradiation facility in Garching.

An amazing number of at least 324 fission track and U-Th/He papers has been published over the last three years.

Also, many people have moved in the last three years and it was quite a challenge to keep track of everybody. Therefore the fission track directory is almost up to date, and I believe that with combined forces and lot of chatting in Bremen we may have it as close as it gets in the next issue.

[Matthias Raab]

# Short Tracks

**Joachim Jacobs** was appointed in late 2005 as a Professor of Structural Geology/Tectonics at the Department of Earth Science at the University of Bergen, Norway. He has already established a new fission track laboratory. **David Belton**, formerly a member of Professor Andrew Gleadow's Fission Track Research Group at the University of Melbourne, has recently taken up a position with the Australia's national research organisation CSIRO. Within the Division of Exploration and Mining, David is now responsible for the operation of the nuclear microprobe. This role is based in the School of Physics at the University of Melbourne – so the move hasn't dramatically changed his travel arrangements! David continues aspects of his research in thermochronology as well as collaborative efforts with Professor Gleadow's group. Delegates to the upcoming ECTC Conference in Bremen might be interested to know that David will also be representing Alphachron – through which CSIRO and Patterson Instruments have commercialised the provision of (U-Th)/Helium analytical equipment.

**Michael Krochmal** from Autoscan Systems and **Matthias Raab** are building the first complete electronic archive of OnTrack. Many old versions only exist as hardcopies, and with the help of our intern Ute Eckardt, who scanned the early versions of OnTrack, we are now having a complete set of issues available. The archive is available on: http://www.autoscan.com.au/download.html. Whoever wants to mirror this archive is invited to download all issues from Mike's webpage or contact either of us for a CD.

# Forthcoming Meetings

# ECTC 2006 Bremen

The European Conference on Thermochronology 2006 will take place at Bremen (Germany). The meeting begins on Sunday afternoon July 30 and lasts till Friday afternoon August 04. The Conference is hosted in the lecture theatre of the GEO building, domicile of the Earth Science Department of the Universität Bremen. Registration for conference participants and accompanying persons is open on Sunday afternoon from 15:00 to 20:00 at the GEO building, where the Icebreaker Party will be organized to welcome the delegates.

The scientific program is planned as a series of single sessions and includes oral presentations and posters. Methodological aspects of low-temperature thermochronology will be discussed on Monday July geological interpretation of while 31. the within thermochronological data long-term landscape evolution studies will be the focus during the following days. Posters will be presented on Tuesday, August 01 afternoon. A Round Table discussion, a field trip and the conference dinner

are also part of the program (see below). The abstracts will be published as a volume of the "Schriftenreihe der Deutschen Gesellschaft für Geowissenschaften". Moreover, it is considered to publish a selection of conference papers in a special issue of a leading international journal.

European Conference on Thermochronology

July 30 – August 04, 2006 Bremen, Germany

Conference language is English. Both Information to the Scientific Program (changes to the Program, details for oral and poster presenters, awards) and General Information (registration, internet access, social program, accompanying persons, eating, travel/accommodation) can be obtained from the 3rd Circular. To download the 3rd Circular and the latest version of the Scientific Program, and for further information and update please check: http://www.geopol.unibremen.de/ectc.htm or contact the ECTC2006 Organising Committee (ectc2006@uni-bremen.de).

The scientific program is published below:

Welcome Address			
Time: 8:30–9:00			
Session 1: Methods I			
	Chair: G.	A. Wagner	
Time	Authors	Title	
9:00–9:30	Gleadow, A., Gleadow, S., Belton, D., Kohn, B., Krochmal, M. & Brown, R.	Coincidence Mapping <sup>™</sup> - a breakthrough strategy for the automatic counting of fission tracks in natural minerals	
9:30–9:50	Marpu, P.R., Gloaguen, R. & Jonckheere, R.	Automatic identification of fission tracks using object-based image analysis	
9:50–10:10	Sobel, E.R. & Seward, D.	Influence of etching conditions on apatite fission track etch pit diameter	
10:10–10:30	Ketcham, R.A., Carter, A., Donelick, R.A., Barbarand, J. & Hurford A.J.	An apatite fission-track calibration for all etching protocols: can we all get along now?	
	Coffe	e Break	
Time	Authors	Title	
11:00–11:20	Gerstenberg, H. & Li, X.	Irradiation facilities at the <i>Forschungsneutronenquelle</i> Heinz Maier Leibnitz (FRM II)	
11:20–11:40	Hasebe, N., Ito, K., Carter, A., Hurford, A. & Arai, S.	LA-ICP-MS FT dating: zircon and volcanic glass	
11:40–12:00	Weber, U.D., Prowatke, S., Glasmacher, U.A., Ludwig, T. & Wagner, G.A.	Apatite fission track dating: determination of the <sup>238</sup> U-content using Secondary Ion Mass Spectrometry (SIMS)	
12:00-12:20	20 Suzuki, T. An absolute approach to fission track dating		
12:20–12:40	Foeken, J.P.T., Stuart, F.M., Dobson, K.J., Persano, C. & Vilbert, D.	A diode laser system for heating minerals for (U-Th)/He chronometry	
	Lunch		
Session 2: Methods II			
Chair: G. Bigazzi			
Time	Authors	Title	
14:00–14:30	Garver, J.I.	The significance of radiation damage in zircon for fission track dating	
14:30–14:50	<b>Dobson, K.J.</b> , Stuart, F.M., Dempster, T.J. & Hinton, R.	U and Th zonation in Fish Canyon Tuff zircon: implications for use as a zircon (U-Th)/He mineral standard	
14:50–15:10	Weise, C., Jonckheere, R. & Ratschbacher, L.	Annealing of heavy-ion tracks in monazite	
15:10–15:30	Stübner, K., Jonckheere, R. & Ratschbacher, L.	Helium diffusion in apatite assessed by Elastic Recoil Detection Analysis (ERDA) and implications for (U-Th)/He dating	
15:30–15:50	Spiegel, C., Kohn, B., Donelick, R., Belton, D., Raza, A. & Gleadow, A.	The effect of long-term low-temperature exposure on fission track stability and helium diffusion: constraints from deep sea drillings	
	Coffe	e Break	
Time	Authors	Title	
16:20–16:40	Belton, D.X., Gleadow, A.J.W. & Kohn, B.P.	A universal annealing model: the chlorine-compesated solution	
16:40–17:00	Zattin, M., Bersani, D. & Carter, A.	Raman microspectroscopy: a non-destructive tool for routine calibration of apatite composition for fission-track analyses	
	Roun	d Table	
Chair: A. Hurford			
Time	Speakers	Topics	
	Andriessen, P.A.M.	Initiative for European thermochronology network	
17.00–18.30	Glasmacher, U.A.	TOPO-EUROPE	
	Jonckheere, R.	Etching protocol	
	Raab, M.J.	OnTrack	

Session 3: Asia			
Chair: L. Ratschbacher			
Time	Authors	Title	
9:00–9:30	Carter, A. & Foster, G.	Improving source characterisation through combined fission track and Sm-Nd-Rb-Sr and Lu-Hf-U-Pb analysis on single detrital apatite and zircon grains	
9:30–9:50	Khattak, N.U., Akram, M. & Khan, M.A.	Recognition of emplacement time of the Loe-Shilman Carbonatite Deposit from NW Pakistan: constraints from apatite fission-track dating	
9:50–10:10	Thiede, R.C., Arrowsmith, J.R., Bookhagen, B., McWilliams, M., Sobel, E.R. & Strecker, M.R.	Timing of dome formation in the Tethyan Himalaya, Leo Pargil (NW India)	
10:10–10:30	Campanile, D., Brown, R., Stuart, F., Widdowson, M. & Bishop, P.	The post break-up evolution of the Western Indian high elevation passive margin	
	Coffe	e Break	
Time	Authors	Title	
11:00–11:20	De Grave, J., <b>Van den haute, P.</b> , Buslov, M.M. & Dehandschutter, B.	Thermotectonic evolution of the Teletskoye graben, northern Siberian Altai-Sayan Mountains, by means of multi-method chronometry	
11:20–11:40	<b>Zhou, Z.</b> , Ding, R., Xu, C., Liu, Y., Cheng, H. & Xu, F.	Modelling of low-temperature exhumation rate in Dabie Shan based on (U-Th)/He and fission-track thermochronological data	
11:40–12:00	Liu, S., Weber, U., Xu, ZQ., Glasmacher, U.A. & Wagner, G.A.	The Dabie Shan-Sulu Connection, China: Cause of strong East-West topography change as revealed by low temperature thermochronological data	
12:00–12:20	Tagami, T., Murakami, M. & Nagahara, K.	Probing fault zone heterogeneity on the Nojima fault: constraints from zircon fission-track analysis of borehole and trench samples	
	Lu	unch	
	Session 4	: Gondwana	
	Chair:	J. Jacobs	
Time	Authors	Title	
13:30–14:00	Lisker, F., Läufer, A.L.	The home of the <i>break in slope</i> buried: the Cretaceous Victoria Basin between Antarctica and Australia	
14:00–14:20	14:00–14:20 Emmel, B., Jacobs, J., Dasszinies, M.C. & Combined apatite fission-track and single grain a possibility to identify thermal overprinted cruin Maud Land (East Antarctica)		
14:20–14:40	20–14:40 Raab, M.J., Brown, R.W. & New sub-surface insights into the structural evolution of the Escarpment, South Africa		
14:40–15:00	Persano, C., van der Beek, P., Braun, J. &   The Eritrean high elevation passive margin: is the escarpment retror     Balestrieri, M.L.   or is the plateau degrading?		
15:00–15:20	<b>Daszinnies, M.C.</b> , Emmel, B., Jacobs, J., Grantham, G. & Wartho, JA.	Karoo age rift flank uplift and its implication for Gondwana rifting: constraints from <sup>40</sup> Ar/ <sup>39</sup> Ar hornblende, biotite and titanite fission track analyses in northern Mozambique	
	Coffe	e Break	

#### **Session 5: Posters** Time: 16:00-18:00 Title Poster # Authors Alvarez-Marron, J., Glasmacher, U.A., Thermochronology of selected Variscan basement rocks in the Western 1. Menéndez, R., Marquínez, J. & Fernández, S. Cantabrian Mountain Range Low-temperature thermochronology as a tool for reconstructing scarp Balestrieri, M.L., Abbate, E., Bigazzi, G. & Omer evolution along Red Sea margins: preliminary apatite fission-track data 2. El Bedri Ali from the Sudan coastal strip and continental interior and comparison with data from the contiguous Eritrea margin Low-temperature thermochronology, uplift and denudation history of the Bauer, F., Glasmacher, U.A., Reiners, P., 3. East African Rift system with special emphasis to the Ruwenzori Bechstädt, T., Förster, A. Mountains (Uganda) Bauer, F., Glasmacher, U.A., Reiners, P., The Sabiñánigo delta complex: coupled basin and delta evolution of the 4. Zühlke, R. Jaca Basin, Northern Spain – a thermochronological approach P-partition and Methodology Alternating Proposal (MAP): new statistic-5. Bermúdez, M. & Alson, P. mathematical methodology that involves uranium densities to estimate ages of rocks by means of fission track dating Detrital apatite fission track thermochronology of the Argentine Puna: 6. Carrapa, B. & DeCelles, P.G. implications for early plateau development Carrière, K.L., Glasmacher, U.A., Wagner, G.A. Neoproterozoic to Holocene tectonothermal evolution of the southern 7. & Bechstädt, T. Cantabrian Mountains, NW Spain as revealed by apatite fission tracks Construction of the southern Eastern Cordillera, NW-Argentina: from Deeken, A., Sobel, E.R., Coutand, I., Haschke, 8. early Cretaceous extension to middle Miocene shortening, constrained M., Riller, U. & Strecker, M.R. by apatite fission track thermochronometry Timing of tectonic uplift in the Sierra de Cameros (Iberian Range, Spain): 9. Del Rio, P., Barbero, L., Stuart, F. & Casas, A. constraints from fission track analysis and (U-Th)/He geochronology in apatites Emmerich, A., Bechstädt, T., Glasmacher, U.A. Modelling the basin history of the SW Dolomites with apatite fission 10. & Zühlke, R. tracks (Upper Permian to recent, Northern Italy) Post-Jurassic tectonic evolution of the western part of the Antarctic 11. Faundez, V., Hervé, F., Stöckhert, B., Brix, M.R. Peninsula: thermochronologic constraints from apatite and zircon fission track analyses - a progress report Apatite fission track thermochronology in southern Brazil: Cretaceous-Franco, A.O.B., Hackspacher, P.C., Hadler 12. Tertiary evolution in Ponta Grossa Arch and its relationship with the Neto, J.C., Saad, A.R., Guedes, S. & Iunes, P.J. South Atlantic rift Glasmacher, U.A., Krochmal, M., Bauer, F., Reliability of confined fission-track length measurement with the new 13. Dreyer, C., Dowsey, G., Repenning, C., Warter, Autoscan automated 3-axis high precision stage V., Hackspacher, P. Reaction of an upper plate to varying subduction regimes - concept and 14. Heberer, B., Behrmann, J.H. & Rahn, M. initial results from the Southern Chile Trench The break-up of the Pangea megacontinent: erosional, geothermal and 15. Juez-Larré, J. & Andriessen, P.A.M. climatic implications in Western Europe Krugh, W.C., Densmore, A.L. & Displacement and denudation patterns from a linked normal fault array, 16. Wassuk Range (Nevada, USA) Seward, D. Segmentation and spatial variation of rifting of the Lambert Graben, 17. Lisker, F. & Brown, R. Antarctica Lisker, F., Läufer, A.L., Olesch, M., Rossetti, F. The denudation history of the Outback Shoulder of northern Victoria 18. & Schäfer, T. Land (Antarctica): a key to the evolution of the Transantarctic Basin Can we define the end of Molasse sedimentation? - A concept and first 19. Rahn, M.K. results from dating Hegau tuffs and volcanogenic layers in the uppermost Swiss Molasse sediments Reinecker, J., Spiegel, C., Glotzbach, C., Rahn, Investigating isotherm perturbation: 1. The Lötschberg transect 20. M. & Frisch, W. (Switzerland) Poster # Authors Title

21.	Spiegel, C., Kohn, B.P., Belton, D.X. & Gleadow, A.J.W.	Cooling history of the Kenya rift valley flanks: implications for Late Cenozoic climate change in East Africa	
22.	Spiegel,C., Sachsenhofer, R., Danišík, M. & Privalov, V.	Thermal evolution of an intracratonic rift basin: the Donbas foldbelt, Ukraine	
23.	Swift, D.A., Persano, C., Nienow, P.W., Dowdeswell, J.A. & Evans, J.	Investigating rates and patterns of late Neogene denudation using detrital apatite fission track thermochronometry: Kangerlussuaq, East Greenland	
24.	Timar-Geng, Z., Henk, A. & Wetzel, A.	The influence of convective heat transport on the interpretation of fission- track data – numerical models and case studies	
25.	Ullrich, A., <b>Glasmacher, U.A.</b> , Miletich, R. & Kohn, B.P.	High pressure effects on fission-track formation in apatite	
26.	Ventura, B., Lisker, F., Ehling, BC., Kopp, J. & Zeh, A.	From the NW Saxo-Bohemian Massif to the southern margin of the Northeastern German Basin: apatite fission track thermochronological constraints on Late Cretaceous tectonic inversion	
27.	Wölfler, A., Danišík, M., Dunkl, I., Dekant, C. & Frisch, W.	Exhumation and thermal history of the Kreuzeck basement complex (Eastern Alps) as inferred from apatite fission track and apatite (U-Th)/He thermochronometry	
28.	Wüthrich, E., Seward, D. & Dimov, D.	Preliminary thermochronological data on the tectonic evolution of the Bulgarian Rhodope	
29.	Xu, C.H., Mansy, J.L., van den haute, P., Guillot, F., Zhou, Z.Y., De Grave, J. & Selen, N.	Apatite fission-track thermochronology of Paleozoic rocks in the Ardenne	
30.	Zentilli, M., Grist, A.M. & Williamson, M-C.	Paleothermal effects of salt diapirs detected by apatite fission track thermochronology and (U-Th)/He dating	

Session 6: Europe I			
Chair: L. Barbero			
Time	Authors	Title	
9:00–9:30	Gibson, M., Sinclair, H.D., Lynn, G.J. & <b>Stuart, F.M.</b>	Last to post-orogenic exhumation of the Pyrenees, Western Europe	
9:30–9:50	Fellin, M.G., Vance, J.A., Garver, J.I. & Zattin, M.	Long low-temperature thermal histories and zircon fission-track annealing: a case study from Corsica	
9:50–10:10	Kuhlemann, J., van der Borg, K., Bons, P.D., Danišík, M. & Frisch, W.	Erosion rates on subalpine paleosurfaces in the western Mediterranean by in-situ <sup>10</sup> Be concentrations in granites: implications for surface processes and long-term landscape evolution in Corsica (France)	
10:10–10:30	Persano, C., Stuart, F.M., Barfod, D.N., Bishop, P. & Dempster, T.J.	Tertiary denudation of Scotland during impact of the proto-Iceland plume from low temperature thermochronometry	
	Coffe	e Break	
Time	Authors	Title	
11:00–11:20	<b>Dobson, K.J.</b> , Stuart, F.M., Dempster, T.J., Persano, C. & Bell, B.R.	The timing and mechanisms of post rift denudation within the Palaeogene Hebridean Igneous Province: a multiple thermochronometer study	
11:20–11:40	Dreyer, C., Glasmacher, U.A., Bauer, F., Stockli, D., Weber, U. & Wagner, G.A.	Cretaceous to Paleogene temperature and exhumation history of the Odenwald revealed by apatite fission-track and (U-Th)/He data	
11:40–12:00	Ventura, B., Lisker, F. & Kopp, J.	Varying denudation and uplift pattern across the Elbe Fault System, East Germany	
12:00–12:20	Wauschkuhn, B., Jonckheere, R., Renno, A. & Ratschbacher, L	Methodical limits of the apatite fission-track-temperature-time-path- modelling: borehole data from the <i>Kontinentale Tiefbohrung (KTB)</i>	
	Lu	inch	
	Session	7: Europe II	
	Chair: I	D. Seward	
Time	Authors	Title	
13:30–14:00	Barbarand, J., Cathelineau, M., Carter, A., Fourcade, S., Pagel, M. & Zeyen, H.	Uplift and erosion of western Europe: what might be the role of hot fluid circulation?	
14:00–14:20	Cederbom, C., Schlunegger, F., van der Beek, P. & Sinclair, H.	Climate versus tectonically driven Pliocene uplift of the European Alps: new fission track data from the North Alpine Foreland Basin	
14:20–14:40	Glotzbach, C., Spiegel, C., Rahn, M. K., Reinecker, J. & Frisch, W.	Investigating isotherm perturbations: 2. The Gotthard transect (Switzerland)	
14:40–15:00	Malusà, M.G., Zattin, M., Andò, S., Garzanti, E. & Vezzoli, G.	Erosional patterns constrained by fission-track grain-age distributions of detrital apatites: preliminary results from the Po River modern sands	
15:00–15:20	<b>Boztuğ, D.</b> , Tichomirowa, M., Jonckheere, R., Harlavan, Y. & Bombach, K.	<sup>207</sup> Pb/ <sup>206</sup> Pb, K/Ar and fission-track geothermochronology of the S-I-A- type granitoids revealing the continent-oceanic island arc and continent- continent collision stages of the Neo-Tethyan convergence system in central Anatolia, Turkey	
	Coffe	e Break	
Session 8: From fission track dating to landscape evolution			
Chair: R.W. Brown			
Time	Speakers	Торіс	
	Garver, J.I.,		
15:50–16:50	Van den haute, P.,	A short history of fission track research	
	Kohn, B.P.		
Laudations			

Session 9: America			
Chair: M.R. Brix			
Time	Authors	Title	
9:30–10:00	Kohn, B., Gleadow, A., Kohlmann, F., Belton, D., Osadetz, K. & Brown, R.	Low temperature thermochronology on cratons: rechecking the rules of the game	
10:00–10:20	Thomson, S.N., Tomkin, J.H., Brandon, M.T. & Reiners, P.W.	Does glacial erosion control the height of mountains? Using thermochronology to find an answer in the Patagonian Andes	
10:20–10:40	Juez-Larré, J., Dunai, T.J. & González, G.	Subduction dynamics and climate change along the Andean margin of Chile, assessed by low-temperature thermochronology and cosmogenic nuclides	
10:40–11:00	Montario, M.J., Garver, J.I. & Reiners, P.W.	(U-Th)/He dating of fission-track-dated zircon with an example from Peru	
Coffee Break			
Time	Authors	Title	
11:30–11:50	Hackspacher, P.C., Saad, A.R., Saenz, C.T. & Hadler Neto, J.C.	Constraints on the evolution of the Cenozoic continental rift system of SE-Brazil: fission track on apatite and zircon	
11:50–12:10	Riley, B.C.D. & Garver, J.I.	Controls on low-temperature resetting of natural damaged detrital zircons: case study from Arizona	
12:10–12:30	Min, M., Ratschbacher, L., Jonckheere, R., Enkelmann, E., Tichomirowa, M., Bachmann, R., Nelson, B., Martens, U., McWilliams, M. & Weber, B.	Long-term evolution of continental transform fault zones: thermochronology along the northern Caribbean plate boundary (Guatemala-Honduras)	
Closing Ceremony			



# 16th Annual V.M. Goldschmidt Conference 2006

### 27 August - 1 September 2006

Melbourne Exhibition and Convention Centre, Melbourne , Australia

The Goldschmidt Conference has a number of sessions related to thermochronology under the Themes 7 and 10. The most relevant sessions amongst them are listed below. The full scientific program is available under www.goldschmidt2006.org.

# Theme 7: Geochemical constraints on timescales and mechanisms of tectonic processes

S7-02: Up and down: Geochemical constraints on paleotopography and tectonic geomorphology *Convenors:* Matt Kohn Barry Kohn

Topography reflects the complex interplay between tectonic and climatic processes, so topographic changes have fundamental influences on regional climate, flora and fauna, erosion, and structural and sedimentation styles. In addition, topography is a direct reflection of the internal force balance of an orogenic system, so paleoelevation histories can discriminate between competing orogenic models. In this symposium, we invite contributions that use geochemical proxies of paleoelevation to investigate a large range of geologic, climatic, and tectonic

processes, e.g. thermochronology and exhumation histories, stable isotopes and plateau/range heights, atmospheric thermodynamics and paleoaltimetry, cosmogenic radionuclide exposure dating of slowly eroded surfaces, and the interplay among climate, erosion and topography.

S7-03: Fast and furious versus slow and steady: rates of geological processes *Convenors:* Gordon Lister Simon Turner

Speakers in this session will be able to address the myriad questions that surround the issues of gradual versus episodic evolution in Earth process. Presentations will range from discussions as to the rate of evolution of landforms, the rate of magma formation, differentiation, degassing and emplacement, how fast can deformation and metamorphism take place, and how fast can rocks cool. Can vastly different processes be synchronized on a planetary scale? What does this imply in terms of System Earth? Many different isotopic systems can be utilized to address the questions that are of interest.

#### S7-06: Fault systems: their geochronology and geochemistry Convenors:

Mike Cosca (mcosca@unil.ch) Institute of Mineralogy and Geochemistry, University Lausanne Simon Kelley (s.p.kelley@open.ac.uk) Department of Earth Sciences, The Open University

Fault systems are ubiquitous features of the Earth's lithosphere and play critical roles in the exhumation of deeply buried crust, fluid transport, and the development of local and regional topography. This symposium is intended to highlight recent geochemical and geochronological advances in our understanding of the varied geological processes associated with deep and shallow fault systems. We invite contributions from a broad subject range that includes, for example, radiogenic and cosmogenic dating of mylonites, pseudotachylytes, extensional detachments, fault scarps and gouges, and the geochemistry and isotopic signatures of fault systems associated with oil and mineral deposits.

# Theme 10: Surface processes, low temperature systems and landscape evolution weathering

# S10-01: Geochemistry, chronology and global consequences of terrestrial weathering

*Convenors:* Paulo Vasconcelos Jerome Gaillardet

Distinct experimental approaches – the mineralogical and isotopic composition of marine sediments, and the age and rate of formation of weathering profiles on continental landscapes – reveal variations in the intensity of weathering processes through time. This session will examine compatibilities and discrepancies between the two approaches; it will explore new approaches suitable for investigating changes in weathering rates; and it will scrutinize the causes and consequences of varying weathering rates through time.

S10-02: Low temperature thermochronometry: models, methods and applications

Convenors: Roderick Brown Andrew Gleadow Cornelia Spiegel

Rapid progress is being made in extending the reach of low temperature thermochronometry through the development of new analytical protocols for new minerals (Fe-Mn oxides, zircon etc) and in more sophisticated models, both for calculating cooling ages and for converting the ages (including track lengths and diffusion data) into thermal history information. We invite papers on any aspect of lowtemperature

thermochronometry especially those reporting new analytical procedures. Papers on novel applications and/or which use an intergrated approach to inverse modelling of diverse thermochronometry data are also particularly encouraged.

S10-03: Terrestrial cosmogenic nuclides: surface process rates and/or dates? Convenors:

Tibor Dunai Fin Stuart Derek Fabel

In-situ produced cosmogenic isotope abundance measurements have enabled acompletely new approach to studying the Earth's surface environment. Advances insample preparation, measurement protocols and analyses of new isotopes in a widerrange of materials are producing larger data sets with more precision and more rapidly. These data offer opportunities for more sophisticated and ambitious applications which are driving development of more complex interpretive strategies and models. The resultant range of quantitative constraints on near surface earth processes now extends well beyond the exposure age, versus steady state erosion rate, options. The aim of this session is to discuss these recent developments, encompassing all aspects of measuring and applying cosmogenic isotope abundances to studying the Earth's surface evolution. We particularly encourage presentations on theoretical and practical issues related to studying catchment scale processes, and sampling and analytical strategies involving stable and/or radionuclides enabling extension to longer time scales.

# **Past Meetings**

Every good meeting needs a good finale. The pictures below are a courtesy of Roderick Brown - who is also responsible for the captions - and Michael Krochmal. Enjoy reminiscent memories of the 10<sup>th</sup> International Conference of Fission Track Dating and Thermochronology in Amsterdam in summer 2004:



Iconic Amsterdam



Stewart Clark, Simon Holdsworth, Justine Tinker, Kate Dobson, Daniel Campanile, Darrel Swift



Three 'old hands'; Andrew Gleadow, Ed Sobel and Trevor Dumitru



Three more 'old hands'; Paul Green, Paul Andriessen and Mike Krochmal



Günther Wagner and Gulio Bigazzi



Margaret and Ray from A to Z



Canal boat trip



Paul Green surprised that Paul Andriessen has turned down a second glass of red!



Glasgow stalwarts; Simon Holdsworth (who's actually from Birmingham), Cristina Persano, Darrel Swift and Stewart Clark



Jurgen Foeken and Cristina Persano



Paul Green supervising Rod Brown attempting to count to ten.



Kate and Callum



Günther Wagner and Andy Gleadow



Günther Wagner's assessment of the success of the Amsterdam meeting!

# Current PhD Studies

#### Perturbation of isotherms below topography: constraints from tunnel transects through the Alps

**Christoph Glotzbach**, supervised by John Reinecker & Cornelia Spiegel, Universität Tübingen, Germany

The effect of perturbation of near surface isotherms by topography has been described early in the 20th century. The magnitude of perturbation depends on several parameters: exhumation rate, geothermal gradient, wavelength and amplitude of topography, and finally on the age of surface relief change. Recently, a number of studies were pursued to quantify the effect of topography on low temperature isotherms using analytical calculations of a simplified topography. Modelling results suggests that in rapidly exhuming areas lowtemperature thermochronologic data are affected by the perturbation of near surface isotherms. Thus, resulting exhumation rates deduced from age elevation profiles may be overestimated.

Our study aims to directly measure the perturbation of isotherms under a given framework of denudation rate, topographic wavelength, relief amplitude, and geothermal gradient by applying low-temperature thermochronology (zircon fission track, apatite fission track and (U-Th)/He analysis). We sampled three tunnel transects through the Alps (Gotthard and Mont Blanc road tunnels and Lötschberg railway tunnel), as well as their corresponding surface lines. All transects are situated in areas with high local relief and presentday uplift rates between 0.7 and 1.5 mm/y. Thermochronological analysis will he complemented by steady state and time variable 3-D thermal modelling. This will allow comparing different model-scenarios with the measured age patterns. and testing paleotopographic reconstructions of the investigated areas. Additionally, our study aims to quantify the perturbation-induced overestimation of exhumation rates inherent to different dating techniques in orogenic settings, and thus to provide correction factors.

# Low Temperature Thermochronology of the Krishna-Godavari Basin, India and its Hinterland

**Himansu S. Sahu**, supervised by Matthias Raab and Andrew Gleadow, The University of Melbourne, Australia In an attempt to bring additional and robust tools towards hydrocarbon exploration, the current study focuses on the application of low temperature thermochronology (AFTT and U-Th/He dating technique) to understand the hydrocarbon habitat in the Krishna-Godavari basin, India and, to generate a picture of the landscape evolution in its hinterland through geological time. We are collaborating with ONGC (Oil and Natural Gas Corporation of India), which has provided 153 borehole samples (both onshore and offshore) from 10 different wells. In addition to that, 49 surface samples have been collected from the different areas of the hinterland giving special attention on the past tectonic activities and present drainage pattern in the area.

A combination of organic maturity data (Vitrinite Reflectance) and the analysis of low temperature thermochronology data will allow the identification of significant heating episodes in the past in which rocks reached palaeotemperatures which might have been higher than their present temperatures in the basin. It will quantify the effects of these palaeothermal episodes and the timing of cooling from maximum palaeotemperatures. Thus a thorough understanding of the thermal history of the basin as well as hydrocarbon habitat will be achieved.

The results of the low temperature thermochronology analysis on hinterland rock types will provide information on the amount of denudation through geological time as well as the sediment types being discharged into the basin. Models on mass balancing will incorporate the removal of crustal material from the continent and its redistribution onto the continental shelf, taking the volume of sediment deposited offshore into account.

#### Neogene Extensional Deformation in Northern Baja California, Mexico: Implications from Combined Low-Temperature Thermochronology and Structural Studies

**Christian Seiler**, supervised by Andrew Gleadow and Chris Wilson, The University of Melbourne, Australia

The Gulf of California provides the unique opportunity to investigate the transition from continental break-up to the development of passive continental margins. Despite its exemplary characteristics for many of the more evolved passive margins around the world, the complex evolution of this highly oblique rift-drift system is still poorly understood. The interaction between time, temperature and strain localization plays a central role in the evolution of the incipient margin adjacent to the Gulf of California and needs to be resolved in further detail if we are to understand when, where and how extensional deformation occurred.

Using the powerful combination of both the fission track and (U-Th)/He thermochronometers in conjunction with structural geology, the project focuses on mapping and structural work in a highly extended portion of the rift system as well as the acquisition and interpretation of a regional low-temperature thermochronological dataset in an area where other timing constraints on extensional deformation are scarce.

#### Modelling the architecture of coupled denudation-sedimentation systems in Australian sedimentary basin evolution

**Stefan Frei**, supervised by Andrew Gleadow, Barry Kohn and Matthias Raab, The University of Melbourne, Australia

Most of the coastal regions of the Australian continental landmass are passive margins and as they developed during and after Gondwana breakup, accumulated an enormous amount of onshore and offshore sediments. Some of these areas now host prolific hydrocarbon deposits and are therefore of economic importance. As the burial history of sedimentary basins is the controlling factor of the thermal maturation process, an understanding of the sources of sediment supply through time and their interactions with basin architecture and stratigraphy potentially provide important insights into basin evolution and the hydrocarbon systems they contain.

The aim of this PhD project is to provide a capability to map out the source regions for large sedimentarv accumulations and to better understand the basin architecture and evolution of important hydrocarbon-bearing environments. Using low-temperature thermochronology, particularly the fission-track method, this technique presents a new perspective on sedimentary basin evolution by investigating the timing, magnitude and patterns of erosion in the basinal hinterland regions and revealing the course of broad-scale drainage patterns in the evolving landscape. The study will focus mainly on the southeastern and northwestern regions of Australia.

#### New Approaches to Low-Temperature Thermochronology

**Steven Spencer**, supervised by Andrew Gleadow, Barry Kohn, The University of Melbourne, Australia, and Shengbiao Hu, Chinese Academy of Sciences, Beijing.

This project aims to identify and characterise potential causes of excessively old apatite He ages, which are substantially older than their coexisting FT ages. Such ages have been attributed to different causes and some of these have been further tested, but this study in particular, evaluates the potential for alpha particles to be implanted into apatite grains 'in situ' by decay in high U- and Thbearing neighbour minerals.

In addition, a combined apatite fission track and (U-Th)/He thermochronometry study has been undertaken on the eastern North China Craton (NCC). Basement samples were collected from the Shandong Province, south of the Bohai Basin, as a transect across the >2000km long, NE-SW trending Tan-Lu Fault and from a vertical profile at Mount Taishan. The eastern NCC margin has experienced predominantly compressive tectonic activity since the Palaeozoic with rifting dominating in the Cenozoic. This study will elucidate the timing of uplift and denudation associated with the interplay of regional compressional and rifting episodes, which have impacted the later stage tectonic evolution of the continental margin.

# The post break-up margin evolution of northern Namibia (Kaokoveld)

**Fernanda F. Luft**, supervised by Matthias Raab and Barry Kohn, The University of Melbourne, Australia,

The Kaoko Belt in northern Namibia is a typical transpressional orogen characterised by km-scale NNW structures generated during the Late Neoproterozoic collision between Congo and Rio de La Plata cratons. These old regional structures (i.e. Purros Mylonite Zone, Khudobib and Sesfontein Thrust, and Ambrosiusberg Fault) have been reactivated during the Phanerozoic and played an important role in the tectonic evolution of the Kaoko Belt and influenced the sedimentary history of the Namibe and Walvis basins.

This project aims to investigate the denudation and structural history for this belt since continental break-up (ca 130 Ma). It will provide crucial information on plate readjustments and the effects on passive margin evolution. 75 basement samples, collected in transects orthogonal to the structural trend of the Kaoko Belt, will be analysed by apatite fission track and (U-Th)/He dating. This aims to enhance our understanding of the low temperature denudation history for the study area with a particlular emphasis on: 1. Significant structural offsets related to extensional and post-rifting tectonics, 2. potential application in petroleum exploration in the adjacent offshore basins

#### Low-temperature thermochronology of the northern Canadian shield, and new insights into fission track behaviour in solids

**Fabian Kohlmann**, supervised by Barry Kohn, Andrew Gleadow and Matthias Raab, The University of Melbourne, Australia

The project comprises two parts. One part, carried out in collaboration with the Geological Survey of Canada, aims to reconstruct the low temperature thermal history of the Slave-Rae craton in the Northwest Territories, Canada, principally using a combination of (U-Th)/He and fission track thermochronology. Particular attention will be given to kimberlite fields and different dyke swarms intruding the shield. The study will test quantitatively the paradigm of long-term craton stability and the recently proposed mechanism of radiation-enhanced annealing.

Another part of the project will investigate the formation of fission tracks in solids, using an Atomic Force Microscope (AFM), an instrument which is able to image the surface of minerals down to the atomic scale. The AFM approach aims to obtain new insights into fission track development in different minerals and their etching behaviour close to the nanoscale. In work so far, fission tracks were implanted in mica and fluorapatite using a 50µCi 252Cf source and imaged with an AFM. Preliminary data in mica suggests that unetched fission track openings form crater-like structures on the mineral surface with diameters between 16-28 nm. Some openings have a pronounced raised rim that probably represents a region of strained lattice around the track core.



3D-AFM image of an unetched fission track in mica

### **Recent PhD Theses**

# CALIBRATION OF THE FISSION-TRACK METHOD AND APPLICATION IN CENTRAL CHINA

#### EVA ENKELMANN

Technische Universität Bergakademie Freiberg Institut für Geologie, Bernhard-von-Cotta Strasse 2, 09599 Freiberg, Germany

This thesis deals with the determination of independent fission-track ages, i.e. ages that are not calibrated against the reference ages of age standards. Recent methodical advances imply that problems related to the value of the fission-decay constant and neutron fluence measurements need not remain obstacles to establish an independent fission-track method, provided that sources of systematic error related to the track counting can be accounted for. The proposed z<sub>0</sub>-factor (GQR; Jonckheere, 2003) was determined for apatite using three different approaches; calculation, experiment and based on age standards. The results are consistent and consequently endorse the validity of the individual applied calibration approaches. The independent fission-track ages (f-ages) of the apatite age standards show that the ages of the accepted standards (DUR; FCT) are consistent with their reference ages but that those of the proposed age standards (MTD; LIM) are not. The latter are therefore not recommended for calibration. The f-, Z-, and z- ages of samples dated in the course of geological investigations are consistent and agree within their 1s errors. The fact that the f-ages agree with their Z- and z-ages is, however, inconsistent with the ~10 % length reduction of confined spontaneous tracks in the accepted standards; the same applies for the agreement between the independent ages obtained on accepted age standards. It is concluded that these results indicate that ~10 % reduction of the spontaneous tracklengths in FCT and DUR has no measurable effect on their fission-track ages. The calibration approach is extended to titanite, which presents the specific problem that the track etching and counting efficiencies depend on the accumulated radiation damages. The experimental calibration approach leads to a significant overestimation of GQR, because it requires prior annealing of the titanite. The GQR-values determined on the basis of age standards (FCT; MTD) are, however, consistent with the calculated value. The GQR-value determined on the proposed Limberg t3 standard is different, probably not because of an invalid

reference age but due to numerous defects and dislocations that obstruct track counting. The fact that identical GQR-factors were obtained on standards of different age and uranium content suggests that a single Z-, z- or GQR-value is appropriate for dating titanites within a broad range of densities. An improved equation for calculating the uranium concentration in thick samples using external fission-track detectors is proposed and tested on standard uranium glasses (CN-5; IRMM-540R) and apatite samples (Durango; horse tooth) in which the uranium content was also determined with independent methods (INAA; ENAA; TIMS). The results show that accurate measurements with the fission-track method are possible within a broad range of uranium concentrations and that uranium determinations based on standards are only accurate if the standard and sample are made of the same material.

The Tibet Plateau is the result of the Cenozoic India-Asia collision that doubled the crust underneath Tibet and elevated the topography to 4-5 km. The area north of the Indian indenter can be mainly attributed to north-south convergence; however, east of the longitude of the eastern Himalayan syntaxis, high topography extends far beyond the eastern margin of India. This thesis also deals with the application of the fission-track method to study the exhumation processes in time and space in the eastern Tibet Plateau and the Qinling and to test the existing models of Plateau formation. Late Cretaceous to Miocene ages indicate a phase of very slow exhumation (<0.02 mm/a) and reflect tectonic guiescence and peneplanation, followed by more rapid exhumation (>0.2 mm/a) since the middle Miocene. Westerly younging ages across west-northwest dipping faults indicate out-of-sequence thrusting within the Longmen Shan belt and the Min Shan. Rapid exhumation is estimated to start at 18-11 Ma in the Min Shan area; the other areas of eastern Tibet give lower (7, 10, and 11 Ma in central eastern Tibet) or upper limits (13 and 30 Ma in southeastern Tibet) for the onset. Denudation was significant ≤10 km along the Longmen Shan, Min Shan and in the central Plateau, but the areas north and south of the Longmen Shan were denuded little, likely due to younger uplift. Cretaceous to Eocene ages indicate continuous cooling and slow denudation in the northern and eastern Qinling, which contrasts with rapid cooling at the end of the thermal history in the southwestern Qinling. There, the rapid cooling started at ~9–4 Ma, a few million years later than in the eastern Tibetan Plateau. A compilation of major Cenozoic faults in the eastern Tibetan Plateau and the Qinling, and their kinematic and dynamic characterization show that deformation in the Qinling is dominant strike-slip. Sinistral and dextral strike-slip boundaries, active at the end of the deformation history, bounding the area of rapid Late Cenozoic cooling identified by apatite fission-track thermochronology, outline an area of eastward rock flow.

### THE LOW-TEMPERATURE THERMOCHRONOLOGY OF CRATONIC TERRANES

#### DAVID X BELTON

The University of Melbourne School of Earth Sciences, Melbourne 3010, Australia

Cratonic terranes present many problems for geologists attempting to define those regions of the continental crust that are the core of today's continents. Inherent in the term is the great passage of time, and typically, the term defines consolidated Archaean or Proterozoic crust (Park and Jaroszewski, 1994). Cratons are further distinguished on the basis of tectonic activity. Marshak and his colleagues (1999) suggest that the lack of penetrative deformation or metamorphism is a useful definition but they further narrow this with the restriction of a Precambrian timeframe. Central to either view is the assumption of stability and perhaps senescence. And, since many aspects of geological research involve the detection of stratigraphic, structural or mineralogical change, stability implies a lack, or at least a minimum, of change. Thus the absence of these traditional markers of geological evolution or change, related to these processes, presents significant challenges in the study of cratons. This is particularly so in shield areas - those cratons with exposed basement rocks.

The purpose of this thesis is to investigate key aspects of landscape evolution in two regions, central Australia and southern Africa, where models have been espoused, arguing for extraordinary surface stability or alternatively a simple erosional history or pediplanation. Contemporary thermochronological techniques now permit us to investigate these regions in previously unavailable detail.

The primary technique used in this work was apatite fission track analysis and an introduction to the fundamentals of the method are given in Chapter 1. The theoretical and practical aspects of the fission track method provided the basis for an innovative approach presented in Chapter 2. TASC is a scheme for analysing the raw fission track data so as to extract additional information about the rock's thermal history prior to undertaking traditional inverse modelling techniques. This method (recently described by the author in Ehlers et al., 2005) proved to be a powerful complement to the routine fission track analysis undertaken as part of the Australian and African case studies.

Although first proposed for geological use in the 1960's, the fission track technique really only gained serious application following technical and theoretical breakthroughs in the 1980's. Since then, growing understanding of the processes of annealing and how they might be modelled has allowed the technique continue developing. Chapter 3 is a discussion of this topic that expands on material previously published by the author and colleagues (Gleadow et al., 2002) and presents additional new work. Nevertheless, despite its wide application in tectonic and basin studies amongst others, there remain many improvements to be made and problems to be solved. As part of this project, research into several areas presented opportunities to contribute toward improvement in the apatite fission track technique, with the potential to aid the study of cratonic terranes.

The chlorine content of apatite has a profound influence on the sensitivity of the mineral for recording thermal events. Few current annealing models are capable of comprehensively addressing the variation of chlorine and other trace elements that appear to play a role in the annealing process. This issue is addressed in Chapter 4 where a universal annealing model is proposed to deal with the wide chemical variability observed in real apatites. For this theme, a fresh consideration of the empirical mathematical models was undertaken and all the published annealing data was considered.

Modern inverse modelling is based on a series of robust, but nonetheless empirical, equations that have withstood the test of time. However, with the aim of developing a more realistic and thus predictive model, Chapter 5 introduces an alternative, physico-chemical to modelling the thermal annealing of fission tracks. This work attempts to draw firmer links between the processes of fission track formation, the mechanics of diffusion and the predicted response to variable temperature regimes.

The first of the case studies is presented in Chapter 6 and is a comprehensive investigation of the long-term landscape evolution of the Davenport Ranges in the central Australian Craton. The study employs traditional petrographic methods as well as thermochronology and combines cosmogenic isotope analysis in an assessment of early landscape models. This chapter expands on work previously published by the author and co-workers (Belton et al., 2004) and has implications for our understanding of landscape evolution in the broader context of the Australian Craton.

In order to maximise temperature sensitivity in slow cooled terranes, the relatively new thermochronological technique of (U-Th)/Helium analysis of apatite was trialed on a suite of central Australian samples. The inconclusive results of this experiment prompted an investigation into the possible causes, and an important baseline study was conducted (Chapter 7). The study has implications for routine application of this new thermochronometer in cratonic and other terranes. More importantly the research identified a potential new thermochronometer with an even greater temperature sensitivity and near surface application for use in future landscape studies.

Chapter 8 documents a larger, craton-wide study of the Mesozoic to recent landscape evolution of the Zimbabwe Craton. This work builds on material presented in earlier chapters and provides a broader view of the nature of crustal cooling, structural reactivation and landform development in the cratonic setting of southern Africa

### COOLING HISTORY AND RELIEF EVOLUTION OF CORSICA (FRANCE) AS CONSTRAINED BY FISSION TRACK AND (U-TH)/HE THERMOCHRONOLOGY

#### MARTIN DANIŠÍK

Eberhard-Karls-Universität Tübingen Institute of Geosciences, Sigwartstrasse 10, D-72076 Tübingen, Germany

The island of Corsica in the Western Mediterranean represents an excellent study area for studying the processes of cooling, exhumation, and relief formation. In a relatively small area, two distinct principal domains are distinguished: (i) Alpine Corsica, a complex nappe stack dominated by metamorphic rocks of oceanic origin; and (ii) Variscan Corsica, a well-exposed Variscan crystalline basement made up mainly of unmetamorphosed granitoid rocks. Geodynamic evolution is characterized by Mesozoic extension, followed by middle Late Cretaceous to Eocene shortening, resulting in Alpine collision when the Alpine nappes were thrust onto the Variscan basement. In the Oligocene, the tectonic regime changed from a compressional to extensional, which led to collapse of the Alpine wedge, rifting leading to ocean spreading, and drifting of the Corsica-Sardinia block away from the European mainland. From the geomorphological point of view. Variscan Corsica represents an outstanding ridgeand-valley structure with peaks exceeding 2000 m

in elevation, and with paleosurface remnants of unknown age preserved at elevations between 200 and 2300 m.

The aim of this study is to investigate the thermal and exhumation history and relief evolution of Corsica, and to constrain the age of paleosurface remnants through the use of zircon fission track (ZFT), apatite fission track (AFT) and (U-Th)/He dating techniques. Previous thermochronological studies focused on investigating the inversion of the Alpine wedge in NE Corsica, whereas the major part of Variscan Corsica is virtually unexplored. This study concentrates specifically on these regions with the highest relief and paleosurface remnants.

Altogether 3 ZFT, 67 AFT and 40 (U-Th)/He analyses were performed on samples from the Variscan basement and Eocene flysch sediments. The data published in other studies were carefully revised according to strictly defined criteria and fully integrated into our dataset. ZFT analyses from the central part of the Variscan basement yield ages between 159.2±9.8 and 144.6±10 Ma. These ages record a thermal event related to the Jurassic opening of the Ligurian-Piedmont Ocean. Data from other studies suggest that the ZFT system in the eastern margin of Variscan Corsica was partially reset during Eocene metamorphism, and that during the Cretaceous there was no thermal event affecting Variscan Corsica. In Alpine Corsica, ZFT data together with 40Ar/39Ar data bracket the time of tectonic denudation of Alpine units between ~33 and 19 Ma.

AFT data range from 105.3±7.2 to 16.4±1.4 Ma. AFT ages from Eocene flysch are all completely reset. Based on the AFT data and thermal modeling results, the cooling history of Corsica can be reconstructed as follows: after the Jurassic thermal event related to opening of the Ligurian-Piedmont Ocean, the basement was cooling to near-surface temperatures where it remained until the Early Paleocene. Since the Late Paleocene, the basement was progressively buried below sediments of the foreland basin, and later during Eocene collision it was partly covered by Alpine nappes. At the end of the Eocene, all of Variscan Corsica had been covered by a thick pile of rocks, leading to a total reset of the AFT system in the major part of the basement. In the Oligocene, after tectonic reorganization, the basement started to exhume, which led to the removal of the thick cover from the top of the basement. The removal of the cover occurred by erosional denudation in most of Variscan Corsica, as well as by tectonic denudation in the NE part of the island. AFT ages from the NW part of the basement record an Early Miocene

thermal event associated with the opening of the Ligurian-Provençal Ocean.

(U-Th)/He analyses of 12 samples from nine different paleosurface remnants vielded ages between 22.1±0.9 and 16.0±2.3 Ma. There are no obvious differences in He age between different paleosurface remnants, although the vertical distance in some cases is more than 1600 m. This indicates that all paleosurface remnants in Corsica have their origin in one single paleosurface. Segmentation of the paleosurface and differential uplift of individual blocks happened after the time of cooling through the temperature zone sensitive to the (U-Th)/He system. The age of the paleosurface formation is bracketed between ~120 and 60 Ma, based on FT data, thermal modeling results and considering stratigraphic and structural data. During Eocene collision, the paleosurface was buried by a thick flysch pile and was thus protected from destruction. In the Oligocene, the paleosurface started to exhume and the cover was removed. During the Miocene rotation, the paleosurface was cut by faults, and at ~17 Ma the region was uplifted by differential block movements, creating relief and inducing valley incision. This event can be viewed as the onset of peneplain destruction that occurred mainly by fault-induced valley incision and widening.

The data presented in this study bring new understanding of the thermal, tectonic and morphological evolution of Corsica and provide new insight into the geodynamic evolution of the Western Mediterranean realm. Moreover, it is shown that after proper revision, it is possible to integrate several, otherwise contradicting, FT datasets.

# Articles

### MEASURING WT% CL TO ASSESS AND ALLOW FOR VARIATION IN FISSION TRACK ANNEALING RATES BETWEEN DIFFERENT APATITE SPECIES

#### PAUL F. GREEN

Geotrack International, 37 Melville Road, Brunswick West, Victoria 3055, Australia (submitted 10/04)

#### Introduction

discussion at the 10th International During Conference on Fission Track Dating and Thermochronology, held in Amsterdam, August 8<sup>th</sup>-14<sup>th</sup> 2004, a growing acceptance of the variability in annealing behaviour of different apatite species, often within a single sample, led to contemplation of the way that such variation can be incorporated into routine application of apatite fission track techniques. Two basic approaches were espoused, one in which the differential annealing kinetics are explained in terms of chlorine content (an approach "championed", if you will, by yours truly), and an alternative approach in which etch pit sizes are used to monitor these effects (favoured by Ray Donelick and co-workers). As a result of an (almost) impromptu discussion session chaired in highly capable fashion by Barry Kohn, the suggestion arose that the rival camps might set out protocols by which their favoured approaches could be implemented in routine application. With this in mind, the aim of this contribution is to outline some recommendations for routine integration of wt% CI measurements with the more usual measurements of fission track age and track length data.

#### **Background / Historical review**

Studies of apatite fission track parameters in subsurface samples of volcanogenic sandstone from the Otway Basin in Southeastern Australia (Gleadow and Duddy, 1981) provided the first systematic evidence that apatite grains within a single rock sample could show a significant variation in the degree of annealing (i.e. age and/or length reduction). Since all the apatite grains within a single sample shared a common volcanic origin, and had undergone the same thermal history, an increasing dispersion in fission track ages downhole must clearly reflect differences in annealing rates which must be due to variation in composition and/or structure between the different apatite grains.

Green et al. (1985, 1986) subsequently reported that within one particular sample, from a presentday temperature of ~95°C, the fission track age measured in individual apatite grains correlated closely with the chlorine content of that grain, such that grains containing 0 wt% Cl were totally annealed (i.e. gave a zero age) while grains containing ~2 wt% Cl or above still retained fission



**Figure 1:** Patterns of fission track age and length expected in detrital apatites within sedimentary rocks currently at their maximum post-depositional temperature.



**Figure 2:** Patterns of fission track age and length expected in detrital apatites within sedimentary rocks which have been hotter in the past.

track ages close to the depositional age of the host sediment (i.e. these grains had undergone very little annealing). Peter Tingate made the original measurements at the University of Melbourne as part of his PhD study. Peter's role in this is often overlooked, so we should acknowledge it here.

More recent laboratory studies have confirmed the existence of differences in annealing sensitivity linked to apatite composition. While such studies have consistently shown that wt% Cl exerts the dominant control (e.g. Barbarand et al. 2003; Ravenhurst et al., 2003), elements such as Sr, Mn and Fe have also been suggested as causing differences in annealing rates (Burtner et al., 1994; Carlson et al., 1999), although Crowley et al. (1991) reported that a Sr-rich apatite and a fluorapatite showed similar sensitivities. Other workers have suggested a possible role of rare earth elements (e.g. Carpena, 1998; Barbarand et al. 2003).

But no evidence has yet been reported for the systematic influence of any element other than Cl in natural data, and the influence of these other elements seems to be sub-ordinate to that of Cl. In contrast, a number of recent sedimentary basin studies have consistently illustrated the systematic effect of Cl on annealing sensitivity (e.g. Argent et al., 2002; Crowhurst et al., 2002; Green et al., 2002; Green, 2004).

# Routine implementation of wt% CI measurements

Geotrack's in-house approach involves measurement of wt% CI on every apatite grain in which either age or track length data are measured. Data are then allocated into compositional groups, and apatite grains in each group are treated as separate (but not independent) systems characterised by their own specific annealing



**Figure 3:** Fission track age vs wt% CI in detrital apatites from a Triassic sandstone outcrop in NE England. Each square represents the fission track age measured in an individual apatite grain, with colours dividing the data into groups for analysis. The horizontal black lines show the predictions of the best-fit thermal history.

#### Barbarand et al. (2002): Analyst-1



**Figure 4:** Mean track length in various annealed low-Cl apatites (from Barbarand et al. 2003) plotted against the mean length in Durango apatite from the same annealing run. The diagonal black line defines the 1:1 relationship. Mean track lengths an apatite containing 0.41 wt% Cl are close to the value measured in Durango apatite, whereas mean lengths in low-Cl apatites are always less than in Durango, by a substantial amount which increases as the degree of annealing (length reduction) increases).

constants. Usina our proprietary "multicompositional" annealing kinetic model which incorporates the differential effect of CI on kinetics. model expected parameters in each we compositional group and compare these with the observed parameters in each group. The modelling approach is essentially identical to that described by Green et al. (1989), but using a number of parallel systems rather than a single mono-compositional system. Using maximum likelihood theory similar to that employed by Gallagher (1995) in MonteTrax, we can define the range of thermal histories which give predictions that are consistent with the measured parameters and their variation with wt% CI within 95% confidence limits (and of course the best-fit history). There are a variety of ways this problem can be approached, but the important thing for this discussion is that the variation in annealing kinetics with wt% CI should be taken into account.

Someone pointed out during the discussion in Amsterdam that this is all very well but there is no annealing model currently available that embodies the differential effect of wt% Cl. But in fact, the laboratory annealing data reported by Carlson et al. (1999) and Barbarand et al. (2003) for a variety of apatites provide a useful basis for erecting such a model. The key to this is the approach described by Ketcham et al. (1999) (which is similar to our own approach), where the degree of annealing in different apatites can be inter-calibrated such that predictions for one composition can be immediately converted to equivalent values for other compositions. In this way, for example, the predictions of the Laslett et al. (1987) model could be easily converted to apatites of other CI contents using those recently published experimental data.

Even without using a multi-compositional model for fully quantitative interpretation, the variation of annealing rates with wt% CI could still be utilised in qualitative fashion to extract useful information. Patterns of single grain age vs wt% CI within individual samples are very much like Ar release spectra, and contain information on the degree of annealing. For example, in a sandstone that is currently at its maximum post-depositional paleotemperature, we would expect to see patterns like those shown in Figure 1, whereas in a sandstone that has been hotter in the past, expected patterns are rather different (Figure 2). So this type of data can be very useful in gualitative terms in understanding the thermal history of a sample.

Measuring wt% Cl on any individual grains that stand out as obviously inconsistent with other grains from the same sample provides a basis for



**Figure 5:** Repeated measurements of various elements at a single site within a slab of Durango apatite, using conditions decribed in the text. These results show no systematic shift with time, emphasising that judicious choice of analytical conditions can eliminate spurious effects that can be caused by intense beam currents focussed in a small area.

identifying and excluding possible contaminants (and/or grains dominated by second order controls

on annealing). Figure 3 shows an example of data in apatites from a Triassic sandstone from NE England. The two grains containing between 0.0 and 0.1 wt% CI with older ages are clearly aberrant, and can be excluded from further analysis.

A number of people have commented to us that they don't see any relationship between fission track age and wt% CI in their own data. It's important to remember that such a relationship is only to be expected in samples which have been heated to the temperature range in which the differential effect of the range of CI contents within the sample is maximised (as in the above sample). This is typically ~90 to 120°C, depending on the specific mix of chlorine contents. Even then, a clear relationship will only emerge if the time of cooling is sufficiently different to the time at which track retention began.

But even at lower temperatures the differential effect in the degree of track length reduction is significant, as shown in Figure 4. For annealing runs where the mean length measured in Durango apatite is 14 microns, mean lengths in the more Frich (i.e. low CI) apatites are around 13 microns, while where the mean length in Durango is 13 microns the values in the low-Cl apatites are between 11 and 12 microns. In contrast, an apatite containing 0.41 wt% CI (similar to the value in Durango apatite) gives very similar mean lengths to the values in Durango. This emphasises the greater sensitivity of the low-CI apatites, compared to Durango (so next time you hear someone refer to Durango as a fluor-apatite, be sure to correct them!).

The obvious effect of this is that applying models based on Durango apatite to data dominated by more sensitive (i.e. low CI) apatites will require erroneously high paleotemperatures to produce a given degree of annealing, in turn implying unreasonably high amounts of erosion etc. As I suggested in Amsterdam, I believe this is a major contributor to the "anomalous Late Miocene cooling" effect often seen in MonteTrax solutions derived using the Laslett et al. (1987) annealing algorithm. Use of a modified model appropriate to the F-rich apatites, based on a conversion using the data in Figure 4, should reduce much of this "anomalous" cooling", and should provide more realistic results using data from basement samples which are commonly (but not always!) dominated by CI contents close to zero.

#### **Practical issues**

<u>Measurement of CI</u>: We routinely measure only wt% CI in every grain that is analysed for either age or



**Figure 6:** To test the uniformity of CI in typical detrital apatites, we measured three (more, occasionally) 20  $\mu$ m spots within each grain in two samples chosen at random from samples currently being processed. Results are grouped by colour and symbol for each grain, and emphasise the lack of significant CI heterogeneity in the great majority of grains, with results in most cases falling within a single division of 0.1 wt% width, as employed in routine interpretation.

length. Chlorine contents are measured using a fully automated Jeol JXA-5A electron microprobe equipped with a computer controlled X-Y-Z stage and three computer controlled wavelength dispersive crystal spectrometers, with an accelerating voltage of 15kV and beam current of 25nA. The beam is defocussed to 20  $\mu$ m diameter to avoid problems associated with apatite decomposition (see below), which occur under a fully focussed 1 - 2  $\mu$ m beam. The X-Y co-ordinates of analysed apatite grains within each grain mount are transferred directly from the Autoscan Fission

Track Stage System to a file suitable for input into the electron microprobe (we can do that for "Trevor stages" too). The system is semi-automated, so that after the location of each grain is verified optically the stage shifts to each grain in turn and analyses for CI. As we have three detector channels, we also monitor Ca and P count rates, as an additional check. CI count rates from the analysed grains are converted to wt% CI by reference to those from a Durango apatite standard (Melbourne University Standard APT151), analysed at regular intervals. This approach implicitly takes into account atomic number absorption and fluorescence matrix effects, which are normally calculated explicitly when analysing for all elements. A value of 0.43 wt% Cl is used for the Durango standard, based on repeated measurements on the same single fragment using pure rock salt (NaCl) as a standard for chlorine.

This approach gives essentially identical results to CI contents determined from full compositional measurements, but has the advantage of reducing analytical time by a factor of ten or more. Measuring full compositions would be time prohibitive. We routinely make just a single 30 second measurement in each grain, with every measurement broken up into three sequential 10 second intervals . Monitoring the consistency of counts in each of the three intervals provides an additional check on beam stability etc.

<u>Mobilisation of F:</u> Stormer et al. (1993) showed that electron microprobe measurements in apatite could be severely affected by beam effects producing diffusion of fluorine within the apatite, leading to erroneous results. To illustrate that these effects can be eliminated by judicious choice of analytical conditions, Figure 5 shows the results of repeated determinations of various elements in a slab of Durango apatite, using our standard measuring conditions (15kv accelerating voltage, 25nA beam current, 20µm beam spot) and a longer than usual counting time of 6 minutes, in order to obtain good statistics for fluorine.

These measurements were all made on exactly the same spot, without moving the beam between measurements. Clearly there is no change in the resulting compositions over the total run time of one hour, showing that the effects reported by Stormer et al. (1993) can be eliminated by suitable choice of beam spot size, current etc. Indeed, such effects have been well known for over 30 years, and measuring conditions adopted for routine use since the late 1960's in the University of Melbourne, and more recently in our own electron microprobe facility, have been designed specifically to avoid these effects.

Non-uniformity in CI: At the Amsterdam Conference, Andy Carter showed some results suggesting this could be a major effect in common apatites. And in conversation, many people have expressed the opinion that apatites are likely to be non-uniform. In contrast, we have never seen any evidence of major differences in wt% CI within individual apatite grains. But to test this possibility more rigorously, we recently measured three spots within each grain in two samples selected at random. The results are shown in Figure 6. In my view, this is pretty convincing evidence that there isn't much variation in wt% CI within any of these grains. The gridlines in these plots divide the data into 0.1 wt% CI divisions, in exactly the way that we routinely split our data, and in most cases, all three measurements fall well within a single group.

<u>Choice of group size:</u> As above, we routinely divide the data into 0.1 wt% divisions. This is partly for simplicity, and partly because this is somewhat larger than the typical resolution of probe measurements. A third reason is that this seems to provide a convenient level of discrimination between differential annealing effects, as illustrated for example by Figure 3.

As before, a range of alternative approaches to this problem would be possible, e.g. a logarithmic division, involving break down 0.0-0.1 wt% CI into finer divisions. This could be considered particularly beneficial since Barbarand et al. (2003) suggested that significant differences exist in annealing properties between apatites within the 0.0 to 0.1 wt% CI range. But experience suggests that the temperature differential corresponding to variation within this interval is only a couple of degrees over geological timescales, so we see no practical reason to pursue any finer division of the data. The case study of data from Taranaki Basin Fresne-1, reported by Crowhurst et al. (2002), and further examples on our website (www.geotrack.com.au) further illustrate this point.

<u>Second order effects:</u> There is no doubt that other elements (Sr, Mn, CO<sub>2</sub>, REE, OH?) also influence fission track annealing rates in apatite, as demonstrated by the laboratory annealing results reported by Carlson et al. (1999) and Barbarand et al. (2003), but no evidence of systematic effects similar to Cl have as yet been shown. In routine implementation, analysis of data as a function of wt% Cl allows those few grains which don't follow the trend with Cl to be identified and eliminated (see Crowhurst et al. 2002, and Figure 3, above).



**Figure 7:** Track length distributions in different compositional groups within the Triassic sandstone from which the fission track age vs wt% Cl data were shown in Figure 3. Red histograms show measured data. Blue distributions show the distributions predicted by the best-fit thermal history.

#### Why should we worry about wt% CI?

One obvious reason for worrying about it is accuracy – using a model based on a single composition to extract thermal history information will only provide accurate answers if applied to data in grains of the same composition.

But another important but more subtle reason is that compositional effects can introduce severe biases into data, which might not be obvious at first sight. One example might be of a sediment which has been heated to say 110°C and then cooled to lower temperatures fairly recently. Most of the F-rich grains will have been totally annealed and contain low spontaneous track densities (young ages), while the more CI-rich grains will have undergone only partial age reduction. Since the F-rich grains tend to be the most common (in most cases), the age data would most likely be dominated by these arains, while the track lengths would come dominantly from the more CI-rich grains which will have much older ages and therefore much higher spontaneous track densities. Treating all grains as

equivalent in this case will inevitably distort the data to such a degree that any interpretations are meaningless.

If any further illustration is required of the need to pay explicit attention to compositional effects, Figure 7 shows the track length distributions in different compositional groups from the same Triassic sandstone sample for which the age vs wt% CI data are shown in Figure 3.

Figure 7 shows how the data in various compositional groups undergo a transition from limited annealing of tracks formed up until the paleo-thermal maximum in grains containing high Cl contents to almost total annealing of the corresponding tracks in grains containing 0.0 wt% Cl. The track length distribution passes from a unimodal distribution in high Cl apatites through bimodal forms in intermediate compositions with shorter and longer tracks easily resolved, to a unimodal distribution with a small tail to shorter lengths in the most sensitive apatites. This is very reminiscent of the patterns of variation identified in



**Figure 8:** Relationships between mean track length and fission track age in apatites from a suite of outcropping Paleozoic basement samples from the UK Lake District (from Green, 1986). The track length distributions show the same systematic change as the fission track age is reduced that is seen in compositional groups within the single sample illustrated in Figure 7. For these outcrop samples, the differences largely represent differences in maximum paleotemperature, but in Figure 7 the differences are induced purely by differential annealing, with all grains having reached the same maximum paleotemperature.

a batch of outcrop samples from the English Lake District (Green, 1986), as illustrated in Figure 8.

The point here is that everyone would surely agree that there is no way that all the outcrop samples in Figure 8 could be treated as identical and given a common interpretation. But in the same way, it is equally inadmissible to treat all of the compositional groups within the Triassic sandstone sample in Figures 3 and 7 as identical (even though they do share an identical history).

#### The future

Quantitative definition of the influence of wt% CI on annealing kinetics has to be a major direction of future research in the fission track community, if accurate results are to be obtained. Almost 20 years after it was first reported (Green et al., 1985), the influence of CI on annealing kinetics is still largely ignored in the majority of studies, and it is time that the "head in the sand" attitude adopted in many quarters since that time was replaced by a more rigorous approach. Another major subject for future research should be identification of second order controls. These undoubtedly exist, as discussed earlier, but a systematic influence of elements other than chlorine in "real data" has yet to be demonstrated.

#### **Closing remarks**

I've made the case here for using wt% CI from electron microprobe analyses as the best indicator of differential annealing, and hopefully provided some practical suggestions regarding how it could be routinely implemented. At Geotrack we prefer this approach over the use of etch pit size, for a variety of reasons (but that's a separate issue for another day). But whatever your preference, one thing is clear - any route you take to dealing with differential annealing is better than ignoring it! And our commercial colleagues in Idaho are to be congratulated on their efforts in this direction. It's time for everyone else to join in!

#### References

- Argent, J.D., Stewart, S.A., Green, P.F. and 2002. Underhill, J.R. Heterogeneous exhumation in the Inner Moray Firth, UK North Sea: constraints from new  $\mathsf{AFTA}^{\texttt{B}}$  and seismic data. Journal of the Geological Society of London. 159, 715-729.
- Barbarand, J., Carter, A., Wood, I. and Hurford, A.J. 2003. Compositional and structural control of fission track annealing in apatite. Chemical Geology, 198, 107-137.
- Burtner, R.L., Nigrini, A. and Donelick, R.A. 1994. Thermochronology of lower Cretaceous source rocks in the Idaho-Wyoming thrust belt. AAPG Bulletin, 78, 1613-1636.
- Carlson, W.D., Donelick, R.A. & Ketcham, R.A. 1999. Variability of apatite fission-track annealing kinetics: I. Experimental results. American Mineralogist, 84, 1213 - 1223.
- Carpena, J. 1998. Uranium-235 fission track annealing in minerals of the apatite group: an experimental study. In: Van den haute, P. and De Corte, F. (eds), Advances in Fission-Track Geochronology. Kluwer Academic Publishers, Dordrecht, pp81-92.
- Crowhurst. P.V., Green, P.F. & Kamp, P.J.J. 2002. (U-Th)/He Appraisal of apatite thermochronology as a thermal history tool for hydrocarbon exploration: an example from

the Taranaki Basin, New Zealand. AAPG Bulletin, 86, 1801-1819.

- Crowley, K.D., Cameron, M. & Schaeffer, R.L. 1991. Experimental studies of annealing of etched fission tracks in fluorapatite. Geochimica et Cosmochimica Acta, 55, 1449 - 1465.
- Gallagher, K. 1995. Evolving temperature histories from apatite fission-track data. Earth and Planetary Science Letters, 136, 421-435.
- Gleadow, A.J.W. and Duddy, I.R. 1981. Fission track analysis: a new tool for the evaluation of thermal histories and hydrocarbon potential. APEA Journal, 23, 93-102.
- Green P.F. 1986. On the thermo-tectonic evolution of Northern England: evidence from fission track analysis. Geological Magazine, 123, 493 - 506.
- Green, P.F. 2004. Burial and exhumation histories of Carboniferous rocks of the Southern North Sea and Onshore UK, with particular emphasis on post-Carboniferous events. Proceedings of the Yorkshire Geological Society, (in press).
- Green, P.F., Duddy, I.R., Gleadow, A.J.W., Tingate, P.R. & Laslett, G.M. 1985. Fission-track annealing in apatite: track lenath measurements and the form of the Arrhenius plot. Nuclear Tracks, 10, 323-328.
- Green, P.F., Duddy, I.R., Gleadow, A.J.W., Tingate, P.R. & Laslett, G.M. 1986, Thermal annealing of fission tracks in apatite 1. A qualitative description. Chemical Geology (Isotope Geoscience Section), 59, 237-253.
- Green, P.F., Duddy, I.R., Laslett, G.M., Hegarty, K.A., Gleadow, A.J.W. & Lovering, J.F. 1989b. Thermal annealing of fission tracks in apatite 4. Quantitative modelling techniques and extension to geological timescales. Chemical Geology (Isotope Geoscience Section), 79, 155-182.
- Green, P.F., I.R. Duddy, & K.A. Hegarty, 2002. Quantifying exhumation from apatite fissiontrack analysis and vitrinite reflectance data: precision, accuracy and latest results from the Atlantic margin of NW Europe. In: Doré, A.G., Cartwright, J., Stoker, M.S., Turner, J.P. and White, N. Exhumation of the North Atlantic Margin: Timing, mechanisms and

*Implications for Petroleum Exploration.* Geological Society Special Publication, **196**, 331-354.

- Japsen, P., Green, P.F. and Chalmers, J. in press. Separation of Palaeogene and Neogene uplift on Nuussuaq, West Greenland. *Journal of the Geological Society, London*.
- Ketcham, R.A., Donelick, R.A. & Carlson, W.D. 1999. Variability of apatite fission-track annealing kinetics: III. Extrapolation to geological timescales. *American Mineralogist*, 84, 1235 – 1255.
- Laslett, G.M., Green, P.F., Duddy, I.R. & Gleadow, A.J.W. 1987. Thermal annealing of fission tracks in apatite 2. A quantitative analysis.

Chemical Geology (Isotope Geoscience Section), **65**, 1-13.

- Ravenhurst C.E.; Roden-Tice M.K.; Miller, D.S. 2003. Thermal annealing of fission tracks in fluorapatite, chlorapatite, manganoanapatite, and Durango apatite: experimental results. <u>Canadian Journal of Earth Sciences</u>, **40**, 995-1007.
- Stormer, J.C., Pierson, L.M. and Tacker, R.C. 1993. Variation of F and Cl X-ray intensity due to anisotropic diffusion in apatite during electron microprobe analysis. American Mineralogist, 78, 641-648.

# CALL FOR INTERLABORATORY COMPARISON OF U & TH MEASUREMENTS IN SUB MG APATITE SAMPLES

#### ISTVÁN DUNKL, VOLKER KARIUS, ANDREAS KRONZ, KLAUS SIMON, ALFONS M. VAN DEN KERKHOF AND HILMAR VON EYNATTEN

Geoscience Center, University of Göttingen, Goldschmidtstrasse 3, Göttingen, D-37077 Germany, istvan.dunkl@geo.uni-goettingen.de (submitted 8/2/05)

The (U-Th)/He thermochronology needs precise determination of uranium and thorium in single grain accessory apatite crystals. Their weight is usually between 10 and 50 µg and the amount of actinide elements range from 0.01 to 10 ng in a single crystal. The low weigth, the low actinide concentrations and the chemical behaviour of thorium (tendency to adsorb) cause difficulties in these measurements. The Durango apatite is the most widely used age standard, however, only few actinide concentrations have been published so far. The reported Th/U ratios vary between 7.7 and 25. This broad range of the Th/U ratios is mainly due to the different origins of the aliquots (different mines, crystals and perhaps also zoning within the crystals). Hence, we can hardly estimate the consistency of the trace element analytical techniques used in different laboratories. Thus Tibor Dunai recommended already at the East Kilbride He-meeting in May 2004 to distribute an apatite powder for an inter-laboratory crosscheck.

We have performed homogeneity tests on several gem-like apatite crystals using CL imaging, (sensitive for manganese, and the main lanthanides), ICP-MS (28 elements in 52 isotopes), electron microprobe (quantitative traverses, 18 elements) and fission track mapping of uranium (laser ICP-MS mapping, IR spectroscopy and halogen determinations are also planned for the near future). The most homogeneous material contains low amounts of fluide inclusions, relatively few cracks and it is available in the form of several cm sized euhedral crystals (AS-16). This apatite was chosen as test material No.1 and we got a proper mass (over 4.5 kg) of more-or-less gem quality crystals from one locality. We perform a careful selection and pulverisation of ca. 100 g and we will start to distribute the 0.5 g aliquots of AS-16 in January or in February 2005. In order to detect a possible systematic deviation of methods/laboratories we plan to distribute a second material. This showed less perfect results in the homogeneity tests. However the significance of the inter-laboratory comparison will be increased by using two materials. None of these materials come from Durango.

The suggested amount taken for an analysis is 10 mg because (i) this relatively high amount will minimize the weighting error and (ii) this amount will assure the representativeness of these rather homogeneous solids. After the dissolution it is requested to take for analysis an aliquot of the solution corresponding to 15  $\mu$ g apatite what is the average weight of an accessory apatite crystal.

We plan to present the results in the following, splitted form:

Table 1: Person or group & affiliation

Table 2: Method (icp-ms or other); with/without isotope dilution; acid composition and concentration; ml used for analysis; with/without ion-exchange; U, Th and Ca in ppm [solid]; blanks; special comments.

There will be no link between the tables and only one of us (ID) will handle the original raw data of the participants. The relation between the results and laboratories will be kept absolutely confidential. Further on the brand of instruments used for analysis will not be presented in order to keep the anonymity of the laboratories.

The results of the homogeneity tests of the gemlike apatite species and the compilation of their geochemical (and partly crystallographical) survey will be published only after the end of this proposed inter-laboratory comparison. The homogeneity of some of the tested crystals is rather promising; thus we hope that at the end of this survey we will be able to issue a new standard for electron microprobe as well as SIMS and Laser-ICP-MS.

Contact us in case of interest and we will mail an aliquot soon.

# DO I NEED TO AVERAGE ALL MY ZETAS? — OUTLIER TEST BY OUT?LIER

#### ISTVÁN DUNKL

Geoscience Center, University of Göttingen, Goldschmidtstrasse 3, Göttingen, D-37077 Germany, http://www.sediment.uni-goettingen.de/staff/dunkl/ (submitted 9/12/2004)

#### Windows program for testing outliers in data sets having normal or "close-to-normal" distribution, and low number of observations

It is always a dilemma: can we reject the "not nice" results or not? Which criterion can be used? For example in case of zetas if one value is far from all the other: is it necessary to include it in the averaging always, or not, or can we reject a measured values at all? I have compiled in this version of Out?Lier four widely used tests which may give some more exact approaches in these questions. These tests are probably not useful to judge the rejection of a single grain FT age within an EDM sample (grain age population). In the majority of the samples the single grain ages follow some oblic distribution (lognormal, gamma or beta). Thus the usage of these tests - which are based on normal distributions - would give false results.

The main window:



💭 Outlier Test		
File Edit Help		
	Tests of out	liers
(These tes	ts serve only as a very coarse estimati	ion; study the Help for careful usage.)
File: C:\Program	Files\OutLier\O_L-19-test.csv	
example for	general app	lication Mean: 20.09
No. of data: 19		Std. Dev.: 13.22
		Std. Error: 3.03
	Minimum (#3):	Maximum (# 18):
	Value: 10.50	Value: 66.17
<b>Grubbs test</b> (for3-100 data)	<i>T</i> : 0.725 <i>Prob.</i> : > 10 %	7: 3.486 Prob.: <1 %
<b>Dixon test</b> (for 3-25 data)	R: 0.069 Prob.: > 10 %	R: 0.566 Prob.: 0.5-1 %
IQR test (for >5 data) Q1	Q1: 13.875 - 1.5 x IQR: 5.0 within inner fences	Q3: 19.79 mild outlier Q3 + 1.5 x IQR: 28.66
Gauss g-test (for 3-50 data)	g: 0.76 at 5%: OK	g: 6.67 at 5%: outlier
	Reject this grain, copy and recalculate	Reject this grain, copy and recalculate

#### 1. Specifications

1.1. Hardware: Min. 200 MHz .

1.2. Operating system: Tested on WinMe, Win2000 and XP.

1.3. Decimal setting in operating system: DOT (Control Panel/Regional Options: "UK" or "US"; Decimal symbol: "."). Unfortunately the comma as decimal symbol makes troubles (sorry for .fr, .it, .de, .sk and .hu users)

1.4. Maximum of data: 300.

1.5. Input

1.5.1. From comma delimited text files

Order the data in maximum three columns (see an example in the table below; suggested software: any spreadsheet calculator, e.g. Excel, Grapher or Origin). First line: three cells for the description of sample locality, measuring conditions, etc. The first record must contain something.

All other lines:

1st cell: data (obligatory, must be a number)

2nd cell: standard deviation of the data (can be empty, but it is practical to save this value, if the file will be opened by other statistical software).

#### 1.5.2. Paste from the clipboard

Copy an individual, continuous column or a complete block with numbers only, having no empty cells, and paste it into the running Out?Lier.

ОК	ОК	FALSE
13.76	13.76 14.55	13.76 14.55
17.2	17.2 15.29	17.2 15.29
10.5	10.5 13.15	10.5 13.15
14.55		35.65
15.29		

#### 1.6. Output

The results of the tests are presented on screen and can be copied to the clipboard in table form. Paste these data into a spreadsheet calculator program.

#### 2. Usage

#### 2.1. Purpose

These four tests are used to detect outliers in univariate data sets. Tested samples are usually repeated measurements performed in similar conditions. The supposed reason of the aberrant behavior of an outlier is related to some nonsystematic, unknown, technical factor that added the unexpected deviation to the scatter of the data

	sample ID	text	text	
	13.76	3.28	data ID	
	17.2	2.35	data ID	
	10.5	2.69	data ID	
	14.55	4.84	data ID	
	-15.29	2.73	data ID	
sample ID	<b>Iple ID</b> Some identification of the sample; obligatory, text or number Data: obligatory, continuous, only numbers			e; obligatory, text or number y numbers
	Error of the	data; recom	mended, typ	be when available
	Comments	and data ide	entification; c	an be empty, but do not use comma

3rd cell: ID of data point (can be empty)

Export the table as 'Comma delimited' or as 'Comma Separated Variables' text file (extension: '.csv').

which is typical for the given measurement.

2.2. What are the results?

The program shows the results of four most commonly used outlier tests developed for low number of observations. The T, R, Qx and g values are calculated by testing the minimum and maximum values of the sample. These values are compared with the critical values of the tables (or test algorithms). The tables of critical values are listed below. The results of the tests are in framed text fields in the main window of the program.

In the Grubbs and Dixon tests the 'Prob.' values indicate the probability that the tested data belongs to the supposed normal distribution determined by the whole population. Below 5% the pink coloring warns to the low probability. The Grubbs and Dixon tests have complex tables with critical values, thus the results can be expressed by probability ranges (e.g. P=5-7.5%). The IQR test has three categories at P=5%: 'within inner fences' = no outlier, 'mild outlier' and 'extreme outlier'. The Gauss g-test has only two categories: 'OK' and 'outlier' at P=5%.

#### 2.3. What is important to consider?

#### Normality

These tests are developed to detect outliers in samples having normal distributions. If the investigated sample has some other, especially asymmetric distribution (e.g. lognormal), then these tests give false results!

#### Very low number of observations

I have serious doubts on the usage of these tests when the number of observations is below 10. Each test somehow requires the empirical distribution of the data and relates the most extreme value(s) to some parameter of the distribution. If the number of observations is very low (e.g. below 5) then the model is poorly constrained and the estimation is rather approximate.

#### Data with different precision

If the elements of the data set are determined by very different precision, then the application of the tests need special care. End-member of such a population can be just a consequence of lower precision resulting from the given measurement.

#### When real outlier --> then few

If more than 20% of elements of the sample is identified as 'outlier' then (1) assumption on the normality of the distribution was not appropriate, and/or (2) quality of data is not proper, and/or (3) tested sample is composite in character, the elements were derived from two or more independent populations.

#### 2.4. Summary

- All statistical tests can be just a very coarse approximation when the number of observations is low.

- Handle the results of the tests only as suggestions; consider rejection only when the population fulfils the requirements of 2.3.

- A really good reason to remove data from the data set exists only when some independent sign or evidence (e.g. technical problem) is known on the given data having 'aberrant behavior'.

- If the four tests do not give consistent results then it is worth to study the equations of the tests and trace the reason of the inconsistency. In these cases the rejection of the tested data can be problematic.

#### 3. Equations of the tests

#### 3.1. Grubbs test

The test is based on the difference of the mean of the sample and the most extreme data considering the standard deviation (Grubbs, 1950, 1969; DIN 32645; DIN 38402).

The test can detect one outlier at a time with different probabilities (see table below) from a data set with assumed normal distribution. If n>25 then the result is just a coarse approximation.

$$T_{\max} = \frac{x_n - X_{mean}}{s}$$
  $T_{\min} = \frac{X_{mean} - X_1}{s}$ 

where

 $x_i$  or  $x_n$  = the suspected single outlier (max or min) s = standard deviation of the whole data set  $X_{mean}$  = mean

#### 3.2. Dixon test

The test is based on the differences between the data members at the tails of the sample (Dixon, 1953). The test detects one outliers at a time (see the table of critical values below). Normal or close-to-normal distribution of the sample is required.

$$r_{\min} = \frac{x_2 - x_1}{x_n - x_1}$$
  $r_{\max} = \frac{x_n - x_{n-1}}{x_n - x_1}$  when n = 3 to 7

$$r_{\min} = \frac{x_2 - x_1}{x_{n-1} - x_1}$$
  $r_{\max} = \frac{x_n - x_{n-1}}{x_n - x_2}$  when n = 8 to 10

$$r_{\min} = \frac{x_3 - x_1}{x_{n-1} - x_1}$$
  $r_{\max} = \frac{x_n - x_{n-2}}{x_n - x_2}$  when n = 11 to 13

$$r_{\min} = \frac{x_3 - x_1}{x_{n-2} - x_1}$$
  $r_{\max} = \frac{x_n - x_{n-2}}{x_n - x_3}$  when n = 14 to 25

where

 $x_i$  or  $x_n$  = minimum and maximum of the sample

$$g = \frac{x_{extr} - \overline{x}}{s}$$

#### where

 $X_{extr}$  = extreme value of the sample

X = the mean of all other data

s = standard deviation of all other data

#### 3.3. IQR (interquartile range) test

The test is based on the difference between the first and third quartile of the sample (Q1 and Q3). If the most extreme data has 1.5 times IQR distance from the next quartile then it is considered as 'mild outlier'; if the distance is over 3 times IQR then it is considered as 'extreme outlier'.

#### 3.4. Gauss g-test

The test is based on the difference between the suspicious extreme value and the mean and standard deviation calculated from the rest of the sample (Szalma, 1984).

#### 4. The tables of critical values

Are in the Help of the program.

#### References

- DIN 32645. (1994): Nachweis-, Erfassungs- und Bestimmungsgrenze. Beuth Verlag, Berlin.
- DIN 38402, Teil 51. (1986): Kalibrierung von Analysenverfahren. Beuth Verlag, Berlin.
- Dixon, W. J. (1953): Processing data for outliers. Biometrica, 9/1.
- Grubbs, F. E. (1950): Sample criteria for testing outlying observations. Annals of Mathematical Statistics, 21.
- Grubbs, F. (1969): Procedures for Detecting Outlying Observations in Samples, Technometrics, 11/1, 1-21.
- Szalma, J. (1984): Introduction to evaluation of measured data. Tankönyvkiadó, Budapest.
- Weber, E. (1980): Grundriss der biologischen Statistik, 8th Edition, Fischer, Jena.

#### Acknowledgements

The program was tested and the description was improved by Martin Danišik and Balázs Székely.



# Versions of Trakscan packages supplied by Autoscan Systems Pty. Ltd.

### Autoscan Systems now supplies 4 versions of the Trakscan package :

#### A. Trakscan DeLuxe (based on Zeiss microscope)

This new version of Trakscan contains a number of significant improvements over its predecessor. It is based on the formidable Zeiss Axio Imager microscope:

1. The use of a very high-resolution colour digital camera and two monitor screens of matching resolution exploits the human brain's superior object detection and discrimination mechanism.

2. The incorporation of highly accurate external linear sensors allows our stages to achieve excellent positioning accuracy.

3. The new features of on-screen track counting and track length measurement have been made possible through the use of highly accurate imaging components : the microscope, camera and monitor screens.

4. The timing of our new version has been fortuitous : at this time Carl Zeiss (the foremost manufacturer of optical research microscopes) has decided to no longer manufacture or supply the drawing tubes (which were previously used in conjunction with our Trakscan programs and digitiser tablets to measure track lengths).

5. The use of two monitor screens in the latest version of our Trakscan software allows simultaneous viewing of both the grain and mica mounts (with a corresponding increase in usable image size), or simultaneous viewing of the camera image and the program control screen (thus enhancing program usability).

6. Our latest version of Trakscan is supplied with EasyLength, a new facility that allows the user to quickly assess the usability of specific mounts without going through the usual file housekeeping procedures. EasyLength also incorporates a useful graphing facility that provides instant feedback about the quality of the results obtained.

7. The Trakscan DeLuxe package incorporates a Zeiss Axio Imager microscope which has been configured to optimise the utility of this package for FTD work, but also incorporates optical components which facilitate a broad range of microscopic applications. It represents the absolute state-of-the-art in Zeiss microscopes at this time.

#### B. Trakscan Professional (based on Zeiss microscope)

This version contains most of the useful features of the DeLuxe version, but with some modifications to bring it into the range of a lower budget. It is based on a modified version of the Zeiss Axio Imager:

1. A specially configured Zeiss Axio Imager microscope, with most of the features of the DeLuxe microscope, but some economies in objective technology. This version of the Zeiss Axio Imager has been specially configured to maximise its usefulness for FTD work.

2. A monochrome high-resolution digital camera, which retains the crispness and detail of the DeLuxe version, but without the addition of colour information.

3. Standard motor shaft encoders on the stage. While these are not as accurate as the external linear position sensors, they still provide positioning repeatability of the order of 2 microns in X and Y, with positioning repeatability of the order of 0.2 microns in Z.

4. On-screen track counting and length measurement are still incorporated, thus providing a convenient and accurate method of carrying out these functions.

5. The EasyLength package is included.

AUTOSCAN SYSTEMS PTY. LTD. ABN 73 006 275 136

P.O. Box 112, Ormond, Victoria, Australia, 3204 \* Suite 4, 293 Bay Street, Brighton 3186 Australia Telephone: + 61 3 9596 8065 Facsimile: + 61 3 9596 8369 \* Home Page: www.autoscan.com.au Email: autoscan@autoscan.com.au



#### C. Trakscan DeLuxe (based on Olympus microscope)

This version is based on the popular Olympus BX51 microscope:

 The use of a very high-resolution colour digital camera and two monitor screens of matching resolution exploits the human brain's superior object detection and discrimination mechanism, when aided by colour.
The incorporation of highly accurate external linear sensors allows our stages to achieve unprecedented positioning accuracy.

3. The new features of on-screen track counting and track length measurement have been made possible through the use of the highly accurate imaging components : the camera and monitor screens. This was not a practical method with previously available imaging technology.

4. The use of two monitor screens in the latest version of our Trakscan software allows simultaneous viewing of both the grain and mica mounts (with a corresponding increase in usable image size), or simultaneous viewing of the camera image and the program control screen (thus enhancing program usability).

5. Our latest version of Trakscan is supplied with EasyLength, a new facility that allows the user to quickly assess the usability of specific mounts without going through the usual file housekeeping procedures. EasyLength also incorporates a useful graphing facility that provides instant feedback about the quality of the results obtained.

6. This Trakscan DeLuxe package incorporates an Olympus BX51 microscope which has been specially configured to optimise the utility of this package for FTD. It represents the state-of-the-art in Olympus microscopes at this time.

#### D. Trakscan Professional (based on Olympus microscope)

This version contains most of the useful features of the DeLuxe version which is based on the Olympus BX51 microscope, but with further modifications to bring it into the range of a lower budget. It was specially configured to cater for the market segment which prefers to use a digitiser and drawing tube for counting and measuring tracks:

1. A specially configured BX51 microscope as above, but with a drawing tube to allow the use of a digitiser tablet for counting and length measurement of fission tracks.

2. A monochrome high-resolution digital camera, which retains the crispness and detail of the DeLuxe version, but without the addition of colour information.

3. A single high-resolution monitor screen.

4. Standard motor shaft encoders on the stage. While these are not as accurate as the external linear position sensors, they still provide positioning repeatability of the order of 2 microns in X and Y, with positioning repeatability of the order of 0.2 microns in Z.

5. A digitiser tablet for counting and measuring of fission track lengths. This method is preferred by some operators, who believe that the live view through the microscope afford a better means of visual discrimination between fission tracks and other mineral features.

6. The EasyLength package is included.

All of the above versions are supplied with our popular high-performance 3-axis AS3000i/f stage (supplied complete with EL300 electronic controller, JS300 joystick, universal power supply and all necessary connecting cables). The systems are also supplied with a suitable state-of-the-art computer package, which runs the Trakscan program under a current version of Windows.

We look forward to once again meeting our FTD friends at the European Conference on Thermochronology in Bremen, Germany (30 July-04 August 2006).

We would be delighted to assist you in deciding on the best package for your particular needs. Please contact us at Autoscan Systems Pty. Ltd.

Note : The product names above are trade names of their respective owners.

AUTOSCAN SYSTEMS PTY. LTD. ABN 73 006 275 136

P.O. Box 112, Ormond, Victoria, Australia, 3204 \* Suite 4, 293 Bay Street, Brighton 3186 Australia Telephone: + 61 3 9596 8065 Facsimile: + 61 3 9596 8369 \* Home Page: www.autoscan.com.au Email: autoscan@autoscan.com.au
# NEW FEATURES. TRICKS AND TIPS FOR THE USE OF TRACKKEY (TK-NEWS NO. 3. — FOR VERSIONS 4.2.F AND 4.3)

## ISTVÁN DUNKL

Geoscience Center, University of Göttingen, Goldschmidtstrasse 3, Göttingen, D-37077 Germany, http://www.sediment.uni-goettingen.de/staff/dunkl/ (submitted 24/10/2004)

The purpose of Trackkey is to calculate and visualize the single grain ages measured by the external detector method. In the last time I have made some improvements and here I would like to complete the former descriptions of this software (Dunkl, 1991; 1992). At the daily work by data made in different systems the file conversion is a really nerve-racking and unnecessary amusement. That is why the new TK versions contain more import and export filters to help the users and achieve more flexibility. Another issue is the type of the distribution of the single grain ages. It is especially important at the identification of age components at the age-provenance studies of siliciclastic rocks. A new plot of TK (Pearson type distribution) may help us for the better understanding.

Running on MAC: Frank Lisker runs without hindrance Trackkey on MAC with a PC emulation. from his letter of 31/8/2004: Citation Α comprehensive overview about Windows emulation programs is given at http://emulation.net/windoze/. The easiest, smoothest running, most distributed (and most expensive) one is "Virtual PC" (current version: 6.1). Trackkey works excellently on this platform, and the respective files can be transferred straightly from and to the Mac disc. The only alternative I tested is "MacBochs", a rather intricate freeware which can be downloaded from http://bochs.sourceforge.net/.

Import: Trackkey uses a double file system. The data files (with extension: .APA and .ZIR) contain the counting data, irradiation code, chemical, Dpar & morphological information of the crystals and some optional data on the locality, petrography etc. The Nd of the irradiation, the name and U content of the monitoring glass and the zetas (+errors) are stored in one or more "user defined" setting file(s). Thus, at the opening of fission track data files of other systems, some data of the setting file must be added for the proper calculations.

BinomFit-import: At the import of a data file with .FTZ or .FTA extension (Brandon, 1992) Trackkey can calculate the age, but can not extract the (i) type of U glass, (ii) the number of tracks on detector (Nd), (iii) the code of irradiation and (iv) the standard deviation of zeta. A warning message appears, and for the fast usage these values are temporarily supposed (for example glass name: "unknown", irradiation code: "# phantom" and standard deviation of zeta: 10).

MacTrack-import: The first problem is that the text exported MacTrack files have .TXT extension and that should make confusion: not all .TXT files can be imported in this way. At the import Tracckey can not extract the type of the U glass and the code of irradiation; thus the running of the program is stopped and the user is asked to specify these data.

Paste from Clipboard: If the user stores the counting data in a spreadsheet calculator (e.g. in an Excel table) then the new "Paste" function can be useful for a rapid conversion of the data to the Trackkey system (menu: File/New and press the Paste button for more guidance). By this function the user can paste from the clipboard the Ns. Ni and Area columns in the table of the Edit Data window.

Export: A part of the export files help at the presentation of results in tables and drawings and an other part of the export files help at the communication with other systems (e.g. AFTSolve and Binomfit).

New extension of "Grain-age files": The grainage files contain the calculated ages of the grains (Trackkey data files contain only the rough counting data). Formerly both AFTSolve and Trackkey used the ".AGE" extension, but the two file types had basically different formats. In the latest TK versions the extension of the grain-age files has changed to ".GRA" to avoid confusions with Richard Ketcham's file system.

DXF (AutoCad) export: These export files contain the most important plots in vector graphic form. These files of .DXF extension can be imported and further edited by many graphical software.

**AFTINV export:** These files are used for modelling program of Dale Issler and they contain both track counts and lengths. Thus before the export the user must load a length file.

*Copy results to clipboard:* The calculated results can be placed on the clipboard in two formats.

"Menu: Edit / Copy All Results to Clipboard" will result in a table with 34 columns. This complex data record can be useful for an update of the user's internal data base, but too complex to insert in a report or publication as it is. The "Copy Selected Results" option will place on the clipboard the data, which are recommended by the IUGS (Hurford, 1990). The order of columns (see Table 1) is a little bit different from Tony's sequence, but contains all necessary data used in the tables of publications. For the N values square bracket is used because the normal bracket () has an assigned meaning in Excel. 4. By selection of the grain population having chisquare value more than 1 % TK can calculate the so-called chi-square age and draws this population on the age spectrum plot.

**Bar plot:** The latest version can draw bar plot with ages below 1 million year; the only restriction is the maximum number of bins. It is limited to 250, because when the bins are too narrow then they can be simply invisible on the screen.

**Type of distribution:** There is rather robust method for the graphical estimation of the type of the distribution just using its kurtosis and skewness (beta1 and beta2). Trackkey contains a new module to plot the projection point of the data on a diagram showing major distribution types (Menu / Plots / Pearson Distribution). The disadvantage of this method is that it requires relative high number of data. There is a very simple empirical test to check how reliable are the beta values: the user can reject the latest 1.2...5 grains and observe the drift of the recalculated projection point on the plot. The

File	Locality	Code	Strat.	Petrogr.	Cryst.	RhoS	[Ns]	Rhol	[Ni]	RhoD	[Nd]	Chi-sq. P (%)	Disp.	Central Age	±	1s
Pop- 293.zir	Kallis	Gö-11	Mioc 2	sandst	50	66.53	[2531]	55.33	[2105]	4.54	[4798]	65	0.09	32.4	±	3.4

**Table 1**: Summary table placed on the clipboard(track densities are 105 tracks per cm2)

*Length files:* The latest TK versions can open both AFTSolve (Ketcham et al., 2000) and Dumitru's stage files (Dumitru, 1993) containing track length data.

**Chi-square age:** There was an interesting discussion at the Amsterdam meeting on the possible treatment methods of partly reset detrital apatite ages in sandstones. One possibility would be to apply the chi-square age (Brandon, 1992) that aims to identify a population at the youngest part of the single grain age distribution, where the chi-square value guarantees the homogeneity of this population. Using TK the isolation of this population is rather simply:

- 1. Menu: Plots / Spectrum Plot Options / Fix Actual Spectrum.
- 2. Menu: Calculations / Chi-square Age.
- The appearing table shows the single grain ages in increasing order. The eighth column indicates the chi-square value computed from the first n single grain ages.

distribution types of Pearson family are plotted on Fig. 1. A detailed analysis on the distribution of single grain ages is in preparation (comes maybe in the next OnTrack).



Figure 1: Distribution types of Pearson family

Goodness measure of the grains (e.g. PMA): The individual counts in the grains contain always some extra-Poissonal error. This comes mainly form the heterogeneous uranium distribution of the dated phases (see details in Green, 1981). We suggested a method and apply the "Proportion of Measurable Area" to characterize somehow the goodness of the dated grains (PMA; Dunkl and Székely, 2004). This is a number between 10 and 100. The latest TK version presents on the main window the average of the "goodness" estimations if they are performed on all grains. If the user applies some other numeric parameter and type these numbers into the "Goodness" column (the last but one) of the data table in the "Edit Data" window then their average will appear on the main window.

**Calculation of uranium content:** Former TK versions did not consider the different ranges of the fission track particles in glass and in the dated minerals. The new TK-BASE.SET setting file contains computed and the recommended values thanks to the letter communications of Eva Enkelmann and Raymond Jonckheere of 25/10/2004 and according to Jonckheere (2003) further considering the densities of the different materials (see Help on the window Settings / Uranium calculation).

**Online Help:** In the Help menu the user can develop an on-line connection to the Trackkey homepage and it hopefully help to check the latest version and visit the online help pages.

**Example files:** By upgrade delete the old example files from your machine, download the new set of files, and place them in the root of drive c (into the C:\TK-DATA directory). The new example files have speaking names and they will help to test the new functions (e.g.: binomfit-example, composite-file, distinction, L-Dumitru-s-stage, L-in-AFTSolve-format, MACTracks-example, test-activation, test-euhedral-population, with-PMA).

Suggestions, problems and points to be discussed:

- Independently of Trackkey I recommend the usage of <u>Total Commander</u> (<u>http://www.ghisler.</u> <u>com</u>). It is similar to the former Norton Commander. File operations are much more transparent and very easy with such a system having always two lists by side than using "My computer" which shows only one list. Very useful function of it is the F3 function key that shows the content of the files promptly. Of course the unzipping and installation of TK is also very fast and simple with it. - It is recommended to show the extensions of the files in your system. It can make serious troubles and wasting of time if you store the results of a sample in more files having the same file name (which is logic) and different extension/format. Thus it is worth to show the extension and for this to disable the following option: My computer / Tools / Folder Options / View / Hide file extensions for known file types.

- Unfortunately TK runs precisely only when the decimal setting of your Windows is dot (Control Panel / Regional Options: UK/US; Decimal Symbol: . dot!). This is an obscure history, sorry.

There are two points where we need some discussion and agreement for the proper und uniform presentation of FT data (and also for further upgrade of Trackkey):

- How to present the zero-track grains on the age spectrum? If a grain have zero spontaneous track then the zero FT age can be plotted by the bar diagram, but the zero age also involves zero s.d. and this makes impossible to draw a Gauss curve from this grain data. Thus, if a sample contains such grains the bar plot and the age spectrum show different characters. Ann Blythe suggested to apply one spontaneous track for such grains just to calculate some s.d. and draw the age spectrum. I was thinking also in this way and speculating even to apply 1/2 track. If someone has a concrete idea and can argue exactly how many virtual tracks can we apply for this computing then I would be happy to receive suggestions. If I will not get any constraints then the 4.3 version (see below) will perform the computing the age spectra with 1/2track.

- If several samples can be added (as the sample localities are very close, or the ages are identical, etc.) and these mounts were irradiated with identical fluent (same irradiation package, close position) then the situation is easy: Menu / File / Combine Files / Load Another Data File. However, if the mounts were irradiated in different conditions (having different RhoD) we have to export them as "grain-age file" and later amalgamate those files. In this case only the mean age is calculated from the grain-age files. The Ns and Ni numbers can not be used for the computing of pooled and central ages as the grains experienced different irradiations. What to do in these cases? Can we normalize the Ni counts at all and when yes then how? The question is especially sharp when the individual samples pass the chisquare test, but for the amalgamated sample we still do not have the procedure and convention how to calculate the pooled or central age. I would be grateful if we achieve some wise convention for these cases.

**Upgrade:** The actually downloadable 4.2.f version contains all the above listed new functions. However, if you have no serious trouble with your running TK version I suggest to wait a few weeks with the download and upgrade because the coming 4.3 version will contain some more import/export filters (e.g. for the new stage software of Trevor Dumitru, for AFTINV of Dale Issler and for the new AFTSolve of Rich Ketcham). I will launch version 4.3 as soon as I get the example files with the new formats.

## Steps of upgrade

(i) It is recommended to perform the installation of the latest version of TK into a new directory (e.g. Program Files\Trackkey.43).

(ii) Copy the existing personal setting files (having .SET extension e.g. myzircon.set) from the old program directory into the new TK directory. In this way the new version will use all your pre-set values (zetas, Nd numbers, graphical settings, etc.).

(iii) For the correct calculation of the uranium content modify some constants in all your personal setting files: Menu / Settings / Uranium calculation / Zeta method / Apatite = 0.945; Zircon = 0.797; Sphene = 0.935. If you install TK first time you can skip steps (ii) and (iii); in this case just create the personal setting files by modification of TK-BASE.SET.

(iv) Pull an icon from the new TK directory to the Desktop and delete the old icon to avoid confusion.

I am open for suggestions and ready for further upgrades.

## References

- Brandon, M.T. (1992): Decomposition of fissiontrack grain-age distributions. Amer. J. of Science, 292, 535-564.
- Dumitru, T.A. (1993): A new computer-automated microscope stage system for fission-track analysis. Nucl. Tracks Radiat. Meas. 21. 575-580.

- Dunkl, I. (2001): The additional parameters a short introduction to the data handling with TRACKKEY 4.1. OnTrack, 11, 19-22.
- Dunkl, I. (2002): TRACKKEY: a Windows program for calculation and graphical presentation of fission track data. Computers and Geosciences. 28(1), 3-12.
- Dunkl, I. and Székely, B. (2004): Proportion of Measurable Area (PMA) as weighting factor for fission track age calculation compensation of the bias made by the heterogeneity of U distribution in dated crystals. Abstracts of the 10th International Conference on Fission Track Dating and Thermochronology, Amsterdam, p. 10.
- Green, P.F. (1981): A new look at statistics in fission track dating. Nuclear Tracks, 5, 77-86.
- Hurford, A.J. (1990): Standardization of fission track dating calibration: Recommendations by the Fission Track Working Group of the I.U.G.S. Subcommission on Geochronology. Chemical Geology, 80, 171-178.
- Jonckheere, R. (2003): On the densities of etchable fission tracks in a mineral and co-irradiated detector with reference to fission-track dating of minerals. Chemical Geology, 200, 41-58.
- Ketcham, R.A., Donelick, R.A., and Donelick, M.B. (2000): AFTSolve: A program for multi-kinetic modeling of apatite fission-track data. Geological Material Research. 2(1), 1-32.

# IRRADIATION FACILITIES AT THE FORSCHUNGSNEUTRONENQUELLE HEINZ MAIER LEIBNITZ (FRM II)

## H. GERSTENBERG, X. LI

Forschungsneutronenquelle Heinz Maier-Leibnitz (FRM II), TU München, Lichtenbergstrasse 1, D-85748 Garching, Germany

In Garching near Munich, Germany, the Technische Universität München operates the Forschungsneutronen-quelle Heinz Maier-Leibnitz (FRM II), a 20 MW research reactor which had been taken in operation in 2004 and which is open for scientific research since April 2005.

The central component of the reactor is the single cylindrical fuel element - the so called compact core - containing approximately 8 kg of highly enriched uranium. The reactor core is embedded in a large (≈ 13 m³) cylindrical tank containing heavy water, which serves for the reflection and the moderation of the fission neutrons. The most important advantage of this design is that the maximum of the thermal neutron flux density is located within the moderator tank but outside the fuel element in an area which is accessible for scientific installations. Although the reactor has clearly been optimized for neutron beam tube experiments it offers several irradiation devices. Among those the pneumatic rabbit device, the hydraulic rabbit device and in particular the socalled fishing line setup, i.e. a very simple purely mechanical irradiation device, are regarded to be suitable for the irradiation of fission track specimens.

The pneumatic rabbit device is made up of 6 independent irradiation channels, using carbon dioxide instead of air as process gas in order to avoid the disturbing production of Ar-41. Due to its individual vertical position within the moderator tank each channel offers a specific neutron spectrum. The available thermal neutron flux densities in the pneumatic rabbit device range between 4.8E12 cm-2s-1 and 7.3E13 cm-2s-1, the corresponding ratio of thermal/fast neutron flux density covers values between 1300 and 67000. The neutron flux parameters for all of the channels are shown in detail in Table 1. They have been evaluated by means of the irradiation of Al:Au (0.1%), Al:Co (1%), Zr and Ni flux neutron monitors and subsequent gammaspectrometric analysis.\*

The main field of operation of the pneumatic rabbit device is the low dose irradiation of small

Irradiation Position	$\Phi_{\text{th}} (\text{cm}^{-2}\text{s}^{-1})$	$\Phi_{epi}$ (cm <sup>-2</sup> s <sup>-1</sup> )	$\Phi_{\rm f} ({\rm cm}^{-2}{\rm s}^{-1})$
Channel 1	3.5×10 <sup>13</sup>	9.6×10 <sup>09</sup>	2.0×10 <sup>09</sup>
Channel 2	1.5×10 <sup>13</sup>	4.8×10 <sup>09</sup>	4.3×10 <sup>08</sup>
Channel 3	5.0×10 <sup>12</sup>	1.4×10 <sup>09</sup>	7.1×10 <sup>07</sup>
Channel 4	6.8×10 <sup>13</sup>	3.3×10 <sup>11</sup>	4.3×10 <sup>10</sup>
Channel 5	3.9×10 <sup>13</sup>	1.4×10 <sup>10</sup>	5.1×10 <sup>09</sup>
Channel 6	7.0×10 <sup>12</sup>	1.8×10 <sup>09</sup>	1.5×10 <sup>08</sup>

Table 1: Neutron flux densities in the pneumatic rabbit device

Irradiation Position	$\Phi_{\rm th}  ({\rm cm}^{-2}{\rm s}^{-1})$	$\Phi_{epi}$ (cm <sup>-2</sup> s <sup>-1</sup> )	$\Phi_{\rm f} ({\rm cm}^{-2}{\rm s}^{-1})$
Channel 1 - Pos. 1	1.3×10 <sup>14</sup>	2.6×10 <sup>11</sup>	3.9×10 <sup>11</sup>
Channel 1 - Pos. 2	9.3×10 <sup>13</sup>	9.9×10 <sup>10</sup>	2.0×10 <sup>11</sup>
Channel 2 - Pos. 1	1.1×10 <sup>14</sup>	7.5×10 <sup>10</sup>	2.1×10 <sup>11</sup>
Channel 2 - Pos. 2	7.7×10 <sup>13</sup>	3.9×10 <sup>10</sup>	1.0×10 <sup>11</sup>

Table 2: Neutron flux densities in the hydraulic rabbit device (4 most intense channels)



Figure 1: Loading and unloading device of the hydraulic rabbit device

samples typically for activation analysis. In principle, nonetheless, it is also suitable for fission track applications. The polyethylene (PE) rabbits used for the transport of the specimens into and out of the irradiation positions, however, restrict the maximum sample size to a cylinder of 14 mm in diameter and 60 mm in length and the mass of the specimens to 10 g. It is again due the use of PE as rabbit material that the integrated thermal neutron fluence is limited to approximately 2E17 cm-2. On the other hand the rather short transportation time of only about 1 s within the moderator tank allows meeting the required neutron fluence very precisely.

In contrast the hydraulic rabbit device - being based on the pool water driven transport of specimens between the loading/unloading position in the reactor building (see Fig 1) and the irradiation position of the facility - is primarily suited for long term irradiations. It exhibits 2 independent channels both of which can be loaded by up to 5 irradiation capsules at the same time. The irradiation capsules are made from high purity AIMg3 which allows them to be irradiated for several days in high thermal neutron flux densities of up to 1.3E14 cm-2s-1. The neutron flux parameters for the 4 most intense irradiation positions are given in Table 2. In order to avoid their direct contact with the reactor pool water the specimens in the hydraulic rabbit device are additionally contained in a water tight inner Al capsule which allow maximum specimen dimensions of 24 mm in diameter and 55 mm in height. The homogeneity of the neutron flux density along the axis of the irradiation capsule is well below 10%.

For the irradiation of fission track specimens, unfortunately, the hydraulic rabbit device, suffers from the problem of a too high thermal neutron flux density leading to requested irradiation times of only few seconds which due to the low transport velocity of the water driven system can not be achieved reliably. In the past this problem has been overcome by carrying out the irradiations at low reactor power of only typically 250 kW thus prolonging the irradiation times by a factor of 80. The results gained from this irradiation technique apparently met the needs of the involved geologists almost perfectly. The situation, however, is unsatisfying in the respect that an irradiation at the required low rector power is only possible during the start of the reactor after an exchange of the fuel



Figure 2: Fishing line irradiation device at the Forschungsneutronenquelle Heinz Maier-Leibnitz (FRM II)

element or a maintenance period which under the assumption of a regular reactor operation means only 5 times per year.

As a consequence from the above situation an irradiation device was needed which allows the irradiation of medium sized specimens in a moderate, to the highest extent possible pure thermal neutron field and provides at the same time short transportation times into and out of the irradiation position. For this purpose the so called fishing line irradiation setup was designed, built and licensed by the regulator. The irradiation position of this device is located in a light water filled vertical thimble in the moderator tank which can be top loaded from the reactor handling bridge. Its distance to the fuel element (center to center) is exactly 1 m. Vertically the standard irradiation position is the mid plane of the reactor fuel element. The corresponding thermal neutron flux density is approximately 1.1E13 cm-2s-1, the ratio of thermal/fast neutron flux densities is far above 1000. Again the homogeneity of the thermal neutron

field in the irradiation position is in the range of a few percent with respect to the volume of the specimen.

The irradiation technique is very simple. The specimen is contained in an AIMg3 container which is connected by means of a Nylon line to a coil being mounted on the handling bridge of the reactor. From this position the irradiation technician lowers the specimen into the irradiation position or pulls it out in the same manner after the completion of the irradiation by mechanically winding or unwinding the Nylon line from the coil (see Fig. 2). Using this device irradiations requiring neutron fluences in the range of 1E15 - 1E17 cm-2 can be done within minutes at full rector power. On the other hand the transfer time of the specimens into and out of the irradiation position is only few seconds; thus the target neutron fluence can be met reliably. Finally in case of an even considerably lower required neutron fluence the irradiation position can easily be shifted out of the mid plane of the fuel element and after a neutron flux density measurement a less intensive irradiation position may be defined keeping the irradiation times in the range of minutes at the same time.

In summary the Forschungsneutronenquelle Heinz Maier-Leibnitz (FRM II) is able to provide a considerable range of irradiation tools also for geological applications. In the collaboration with external researchers the irradiation service of the FRM II does not only perform the irradiation itself but provides an activation and dose rate calculation for each specimen and offers assistance in the transportation of activated specimens back to the customer.

\* The authors are very grateful to X. Lin, TU München, Institut für Radiochemie, for the performance of the neutron flux density measurements.



Automated microscope stage systems greatly increase operator productivity by automating tedious aspects of microscope work. Since their introduction, our systems have been adopted by far more fission track laboratories than any other system.

## **Outstanding Hardware:**

Our stage systems are based on a highly-reliable, high-precision Kinetek<sup>™</sup> computer-automated microscope scanning stage. Several hundred Kinetek stages are currently in operation, mainly in the demanding microelectronics and biotechnology industries. Compatible with almost any brand of microscope. Use of this popular, general purpose stage significantly reduces the system cost. System also includes a high-quality Calcomp<sup>™</sup> 12x12" digitizing tablet. Assembly to full operational status generally requires only a few days.

## **Outstanding Software:**

Software is a complete, highly sophisticated Apple Macintosh program developed with careful attention to all aspects of microscope work. Fully integrates track counting, track length measurement, slide scanning, and file management functions. Moves precisely from grain to mica print in 3 seconds. Very user friendly.

## Innovative, Highly-Intuitive Stage Control System:

Stage is driven primarily with the digitizing tablet cursor rather than a joystick. For example, to center a grain or track, just superimpose the cursor on it, push a button, and the stage automatically centers it. This avoids tedious manual centering via the joystick. Most software commands are driven from the cursor buttons, which are easily distinguished by feel, so there is no need to look away from the eyepieces to the computer screen or keyboard.

## Fission Track Laboratories Using the System

(year installed; \*adapted to a non-Kinetek stage)

- •Stanford University, Stanford, California (1991)
- •University of California, Santa Barbara, California (1992)
- •ARCO Inc., Plano, Texas (1992). Moved to University of
- Minnesota, Minneapolis, Minnesota, in 1999. •Universität Bremen, Bremen, Germany (1993)
- •E.T.H., Zürich, Switzerland (1993\*)
- •Kent State University, Kent, Ohio (1993)
- •University of Wyoming, Laramie, Wyoming (1993)
- •University of Arizona, Tucson, Arizona (1993). Moved to
- Syracuse University, Syracuse, New York, in 2000.
- •Max-Planck-Institut, Heidelberg, Germany (1993\*)
- •Union College, Schenectady, New York (1994)
- •Monash University, Melbourne, Australia (1994\*). Moved to University of Melbourne in 1999.
- •La Trobe University, Melbourne, Australia (two systems, 1994\*). Moved to University of Melbourne in 1999.
- •University of Pennsylvania, Philadelphia, Pennsylvania (1995)
- •Universität Tübingen, Tübingen, Germany (1995)
- •Universidad Central de Venezuela, Caracas, Venezuela (1995)
- •Brigham Young University, Provo, Utah (1995)
- •Central Research Institute of the Electric Power Industry, Chiba, Japan (1995)
- •Universität Salzburg, Salzburg, Austria (1996)
- •University of Southern California, Los Angeles, California (1996)
- •E.T.H., Zürich, Switzerland (second system, 1996\*)
- •Geologisk Centralinstitut, Copenhagen, Denmark (1996\*)
- •University of Waikato, Hamilton, New Zealand (1996\*)

- •Università di Bologna, Bologna, Italy (1997)
- •Centro di Studio di Geologia dell'Appenno e delle Catene Perimediterranee, Florence, Italy (1997)
- •University of Wyoming, Laramie, Wyoming (second system, 1997)
- •Universität Potsdam, Potsdam, Germany (1997)
- •Seoul National University, Seoul, Korea (1998)
- •E.T.H., Zürich, Switzerland (third system, 1998)
- •Universität Basel, Basel, Switzerland (1998)
- •University of Florida, Gainesville, Florida (1998)
- •Universite Paris-XI, Paris, France (1998)
- •Universität Graz, Graz, Austria (1998)
- •Göteborgs Universitet, Göteborg, Sweden (1999)
- •Universidad de Cádiz, Cádiz, Spain (1999)
- •Universite Montpellier II, Montpellier, France (1999)
- •Kurukshetra University, Kurukshetra, India (1999)
- •Universität Tübingen, Tübingen, Germany, (second system, 1999)
- •California State University, Fullerton, California (2000)
- •Geoforschungszentrum, Potsdam, Germany (2000)
- •Polish Academy of Sciences, Krakow, Poland (2000)
- •University of Glasgow, Glasgow, Scotland (two systems, 2001)
- •Yale University, New Haven, Connecticut (2001)
- •Université Joseph Fourier, Grenoble, France (2001)
- •Universität Bremen, Bremen, Germany (second system, 2002)
- •Université des Sciences et Technologies de Lille, Villeneuve
- d'Ascq, France (2002) •University of Kansas, Lawrence, Kansas (2003)

## Further Information:

An early version of the system is described in a paper in Nuclear Tracks and Radiation Measurements, vol. 21, p. 575-580, Oct. 1993 (1992 Philadelphia Fission Track Workshop volume). For detailed information please contact: Dr. Trevor Dumitru, 4100 Campana Drive, Palo Alto, California 94306, U.S.A., telephone (auto-switching voice and fax line): 1-650-725-6155

# Fission-Track Papers 2003-2006

The 324 references listed below were published since the last issue of OnTrack in September 2003. The list is a combination of references sent to the editor and a GeoRef search for peer reviewed journals considering the keywords "fission track" and "U-Th/He".

## 2003 - 2006

- Aciego, S., DePaolo, D. J., Kennedy, B., Christensen, J., & Anonymous. (2005). U-Th/He dating of basalt. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 322.
- Aciego, S., Kennedy, B. M., DePaolo, D. J., Christensen, J. N., & Hutcheon, I. (2003). U-Th/He age of phenocrystic garnet from the 79 AD eruption of Mt. Vesuvius. Earth and Planetary Science Letters, 216(1-2), 209-219.
- Adriasola, A., Thomson, S.N., Brix, M.R. & Stöckhert, B. (2006). Postmagmatic Cooling and Late Cenozoic Denudation of the North Patagonian Batholith in the Los Lagos Region of Chile, 41°-42°15'S. International Journal of Earth Sciences, 95, p. 504-528, doi: 10.1007/s00531-005-0027-9.
- Aka, F. T., Nagao, K., Kusakabe, M., Sumino, H., Tanyileke, G., Ateba, B., et al. (2004). Symmetrical helium isotope distribution on the Cameroon volcanic line, West Africa. Chemical Geology, 203(3-4), 205-223.
- Albianelli, A., & Napoleone, G. (2004). Magnetostratigraphy of the Homo-bearing Pleistocene Dandiero Basin (Danakil Depression, Eritrea). Rivista Italiana di Paleontologia e Stratigrafia, 110, Suppl., 35-44.
- Aldega L., Corrado S., Di Leo P., Giampaolo C., Invernizzi C., Martino C., Mazzoli S., Schiattarella M. & Zattin M. (2005). The Southern Apennine case history: thermal constraints and reconstruction of tectonic and sedimentary burials. Atti Ticinensi di Scienze della Terra, 10, serie sp., 45-53.
- Alloway, B. V., Pillans, B. J., Carter, L., Naish, T. R., & Westgate, J. A. (2005). Onshore-offshore correlation of Pleistocene rhyolitic eruptions from New Zealand; implications for TVZ eruptive history and paleoenvironmental construction. Quaternary Science Reviews, 24(14-15), 1601-1622.
- Alloway, B. V., Pribadi, A., Westgate, J. A., Bird, M., Fifield, L. K., Hogg, A., et al. (2004). Correspondence between glass-FT and (super 14) C ages of silicic pyroclastic flow deposits sourced from Maninjau Caldera, west-central Sumatra. Earth and Planetary Science Letters, 227(1-2), 121-133.
- Alloway, B. V., Westgate, J., Pillans, B., Pearce, N., Newnham, R., Byrami, M., et al. (2004). Stratigraphy, age and correlation of middle Pleistocene silicic tephras in the Auckland region, New Zealand; a prolific distal record of Taupo
- Armstrong, P. A. (2005). Thermochronometers in sedimentary basins. Reviews in Mineralogy and Geochemistry, 58(1), 499-525.
- Armstrong, P. A., Taylor, A. R., & Ehlers, T. A. (2004). Is the Wasatch Fault footwall (Utah, United States) segmented over million-year time scales? Geology, 32(5), 385-388.
- Arpa, M. C. B., Bornas, M. A. V., Abigania, M. I. T., Solidum, R. U., Listanco, E. L., Ozawa, A. and Tagami, T. (2006). Petrology and chemistry of lava flow and pyroclastis from the February to March 2000 eruption of Mayon volcano, Philippines. JGSP, in press.

- Balestrieri, M. L., Stuart, F. M., Persano, C., Abbate, E., & Bigazzi, G. (2005). Geomorphic development of the escarpment of the Eritrean margin, southern Red Sea from combined apatite fission-track and (U-Th)/He thermochronometry. Earth and Planetary Science Letters, 231(1-2), 97-110.
- Balogh, K. and Dunkl, I. (2005): Argon and fission track dating of Alpine metamorphism and basement exhumation in the Sopron Mts. (Eastern Alps, Hungary): Thermochronology or mineral growth? Mineralogy and Petrology, 83, 191-218.
- Barbero, L., Glasmacher, U. A., Villaseca, C., Lopez Garcia, J. A., & Martin Romera, C. (2005). Long-term thermo-tectonic evolution of the Montes de Toledo area (central Hercynian Belt, Spain); constraints from apatite fission-track analysis. Geologische Rundschau = International Journal of Earth Sciences, 94(2), 193-203.
- Baxter, E. F. (2003). Quantification of the factors controlling the presence of excess (super 40) Ar or (super 4) He. Earth and Planetary Science Letters, 216(4), 619-634.
- Belton, D. X., Brown, R. W., Kohn, B. P., Fink, D., & Farley, K. A. (2004). Quantitative resolution of the debate over antiquity of the central Australian landscape; implications for the tectonic and geomorphic stability of cratonic interiors. Earth and Planetary Science Letters, 219(1-2), 21-34.
- Bermúdez, M., Alson, P. y Mora, J. (2004): "Equivalence between formulas for the calculation of errors in the age provided by the fission tracks dating method". Revista de la Facultad de Ingeniería, UCV. Vol. 19, N° 1, pp. 119-123.
- Bermúdez, M., Alson, P. y Mora, J. (2005): "Fundamental equation of age for the minerals dating and their adaptation to the practical equation for the fission tracks dating method". Revista de la Facultad de Ingeniería, UCV. Vol. 20, N° 2.
- Bernet, M., & Garver, J. I. (2005). Fission-track analysis of detrital zircon. Reviews in Mineralogy and Geochemistry, 58(1), 205-237.
- Bernet, M., Brandon, M. T., Garver, J. I., & Molitor, B. (2003). Downstream changes of alpine zircon fission-track ages in the Rhone and Rhine Rivers. Journal of Sedimentary Research, 74(1), 82-94.
- Bernet, M., Brandon, M. T., Garver, J. I., & Molitor, B. R. (2004). Fundamentals of detrital zircon fission-track analysis for provenance and exhumation studies with examples from the European Alps. Special Paper - Geological Society of America, 378, 25-36.
- Bernet, M., Spiegel, C. (2004). Detrital thermochronology a short introduction. In: "Detrital thermochronology Provenance Analysis, Exhumation, and Landscape Evolution in Mountain Belts, edited by Matthias Bernet and Cornelia Spiegel. GSA Special Paper 378, 1-6.
- Bhandari, N., Murty, S. V. S., Shukla, P. N., Mahajan, R. R., Shukla, A. D., Suthar, K. M., et al. (2005). Bhawad LL6 chondrite; chemistry, petrology, noble gases, nuclear tracks,

and cosmogenic radionuclides. Meteoritics & Planetary Science, 40(7), 1015-1021.

- Biermeier, C., Wiesinger, M., Stuewe, K., Foster, D. A., Gibson, H. J., & Raza, A. (2003). Aspects of the structural and late thermal evolution of the Redbank thrust system, central Australia; constraints from the Speares Metamorphics. Australian Journal of Earth Sciences, 50(6), 983-999.
- Bigazzi, G., Balestrieri, M. L., Norelli, P., Oddone, M., & Tecle, T. M. (2004). Fission-track dating of a tephra layer in the Alat Formation of the Dandiero Group (Danakil Depression, Eritrea). Rivista Italiana di Paleontologia e Stratigrafia, 110, Suppl., 45-49.
- Blisniuk, P.M., Stern, L.A., Chamberlain, C.P., Zeitler, P.Z., Ramos, V.A., Haschke, M., Sobel, E.R., Strecker, M.R. and Warkus, F., in press, Links between mountain uplift, climate, and surface processes in the southern Patagonian Andes, submitted to: Oncken et al. (eds.): The Andes from top to bottom; Frontiers in Earth Sciences, v. 1.
- Bondarenko, G. E., Soloviev, A. V., Tuchkova, M. I., Garver, J. I., & Podgornyi, I. I. (2003). Age of detrital zircons from sandstones of the Mesozoic flysch formation in the South Anyui suture zone; western Chukotka. Lithology and Mineral Resources, 38(2), 162-176.
- Boyce, J. W., & Hodges, K. V. (2005). U and Th zoning in Cerro de Mercado (Durango, Mexico) fluorapatite; insights regarding the impact of recoil redistribution of radiogenic (super 4) He on (U-Th)/He thermochronology. Chemical Geology, 219(1-4), 261-274.
- Boyce, J. W., Hodges, K. V., Crowley, J. L., Chatterjee, N., Searle, M., & Anonymous. (2005). Laser microprobe (U-Th)/He thermochronometry of monazite. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 298.
- Boztug, D., Jonckheere, R., Enkelmann, E., Ratscbacher, L., Wagner, G., & Anonymous. (2005). Geodynamic implications of rapid denudation of the granitoids at about 50 and 20 Ma in the eastern Pontides, Turkey; apatite fission-track results. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 300.
- Boztug, D., Jonckheere, R., Wagner, G. A., & Yegingil, Z. (2004). Slow Senonian and fast Palaeocene-early Eocene uplift of the granitoids in the central eastern Pontides, Turkey; apatite fission-track results. Tectonophysics, 382(3-4), 213-228.
- Brady, R. J., & Spotila, J. A. (2005). Southward-younging apatite (U-Th)/He ages in the Northern California Coast Ranges due to a northward-migrating crustal welt. Earth and Planetary Science Letters, 235(1-2), 107-122.
- Brandon, M., Zattin, M., Isaacson, P., Braun, J., Reiners, P., & Anonymous. (2005). Using thermochrometry to image topographic evolution in the Northern Apennines, Italy. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 300.
- Braun, J. (2003). Pecube; a new finite-element code to solve the 3D heat transport equation including the effects of a time-varying, finite amplitude surface topography. Computers & Geosciences, 29(6), 787-794.
- Brichau, S., Ring, U., Ketcham, R.A., Carter, A., Stockli, D., and Brunel, M. (2006). Spatial and temporal variation of the slip rate for a major extensional fault system on Naxos and Paros Islands, Aegean Sea, Greece. Earth and Planetary Science Letters v241, p293-306.
- Bruegel, A., Dunkl, I., Frisch, W., Kuhlemann, J., & Balogh, K. (2003). Geochemistry and geochronology of gneiss pebbles from foreland molasse conglomerates; geodynamic and

paleogeographic implications for the Oligo-Miocene evolution of the Eastern Alps. Journal of Geology, 111(5), 543-563.

- Bruner, J., Wood, D., Wilson, M., Ting, S., Gingerich, P., Higgins, P., et al. (2004). First reported vertebrate from the Swauk Formation (lower Eocene), near Liberty, Kittitas County, Washington, is a new species of freshwater percomorph fish. Journal of Vertebrate Paleontology, 24(3, Suppl.), 42.
- Burbank , D.W., Brewer, I.D., Sobel, E.R., and Bullen, M.E., in press, Single-crystal dating and the detrital record of orogenesis, IAS Special Publication.
- Burbank, D. W., Blythe, A. E., Putkonen, J., Pratt-Sitaula, B., Gabet, E., Oskin, M., et al. (2003). Decoupling of erosion and precipitation in the Himalayas. Nature, 426(6967), 652-655.
- Calvert, A. T., & Mortimer, N. (2003). Thermal history of Transantarctic Mountains K-feldspars, southern Victoria Land. Terra Antartica, 10(1), 3-15.
- Carrapa, B., Adelmann, D., Hilley, G. E., Mortimer, E., Sobel, E.R. and Strecker, M.R. (2005), Oligocene range uplift and development of plateau morphology in the southern Central Andes: Tectonics, v. 24, TC4011, doi:10.1029/2004TC001762.
- Carrapa, B., Strecker, M.R., and Sobel, E.R., in press, Cenozoic orogenic growth in the Central Andes: Evidence from sediment provenance and apatite fission track thermochronology along the southernmost Puna Plateau margin (NW Argentina), Earth and Planetary Science Letters.
- Carter, A and Gallagher, K. (2004). Provenance signatures and the inference of thermal history models from apatite fission track data – A synthetic data study , Geol Soc. Am. Spec. Publ. 378, 7-23.
- Carter, A., and Gallagher, K. (2004). Characterising the significance of provenance on the inference of thermal history models from apatite fission track data – A synthetic data study. In Bernet, M., and Spiegel, C., eds Detrital thermochronology – Provenance analysis, exhumation, and landscape evolution of mountain belts, GSA Special Paper, 378, p7-23.
- Carter, L., Alloway, B. V., Shane, P., & Westgate, J. (2004). Deep-ocean record of major late Cenozoic rhyolitic eruptions from New Zealand. New Zealand Journal of Geology and Geophysics, 47(3), 481-500.
- Carter, T. J., Kohn, B. P., Foster, D. A., & Gleadow, A. J. W. (2004). How the Harcuvar Mountains metamorphic core complex became cool; evidence from apatite (U-Th)/He thermochronometry. Geology, 32(11), 985-988.
- Carter, T. J., Kohn, B. P., Foster, D. A., Gleadow, A. J. W., and Woodhead, J. D., (2006). Late-stage evolution of the Chemehuevi and Sacramento detachment faults from apatite (U-Th)/He thermochronometry – Evidence for mid-Miocene accelerated slip. Geological Society of America Bulletin, 118: 689-709,
- Carter, T.J., Kohn, B.P., Foster, D.A., and Gleadow, A.J.W. (2004). How the Harcuvar Mountains metamorphic core complex became cool: evidence from apatite (U-Th)/He thermochronometry. Geology, 32: 985-988.
- Cather, S. M. (2003). Polyphase Laramide tectonism and sedimentation in the San Juan Basin, New Mexico. Guidebook New Mexico Geological Society, 54, 119-132.

- Cederbom, C. E., Sinclair, H. D., Schlunegger, F., & Rahn, M. K. (2004). Climate-induced rebound and exhumation of the European Alps. Geology, 32(8), 709-712.
- Chakurian, A. M., Arehart, G. B., Donelick, R. A., Zhang, X., Reiners, P. W., Hofstra, A. H., et al. (2003). Timing constraints of gold mineralization along the Carlin Trend utilizing apatite fission-track, (super 40) Ar/ (super 39) Ar, and apatite (U-Th)/He methods. Economic Geology and the Bulletin of the Society of Economic Geologists, 98(6), 1159-1171.
- Clendenen, W. S., Fisher, D., & Byrne, T. (2003). Cooling and exhumation history of the Kodiak accretionary prism, Southwest Alaska. Special Paper - Geological Society of America, 371, 71-88.
- Clift, P. D., Schlup, M., Carter, A., & Steck, A. (2004). Exhumation history of eastern Ladakh revealed by (super 40) Ar/ (super 39) Ar and fission track ages; the Indus River-Tso Morari transect, NW Himalaya; discussion and reply [modified]. Journal of the Geological Society of London, 161(5), 893-894.
- Clift, P.D., Campbell, I.H., Zhang, X., Carter, A., Hodges, K.V., Khan, A.A., and Allen, C.M. (2004). Thermochronology of the modern Indus River bedload: New insight into the controls on the marine stratigraphic record. Tectonics 23, TC5013
- Colgan, J. P., Dumitru, T. A., & Miller, E. L. (2004). Diachroneity of Basin and Range extension and Yellowstone Hotspot volcanism in northwestern Nevada. Geology, 32(2), 121-124.
- Colgan, J. P., Dumitru, T. A., McWilliams, M. O., & Miller, E. L. (2006). Timing of Cenozoic volcanism and Basin and Range extension in northwestern Nevada; new constraints from the northern Pine Forest Range. Geological Society of America Bulletin, 118(1-2), 126-139.
- Corrado S., Aaldega L., Di Leo P., Giampaolo C., Invernizzi C. & Zattin M. (2005) Combined use of thermal indicators to study the tectono-sedimentary evolution of Southern Apennines (Italy). Terra Nova, 17, 56-65.
- Coutand, I., Carrapa, B., Deeken, A., Schmitt, A.K., Sobel, E.R., and Strecker, M.R. (2006), Propagation of orographic barriers along an active range front: insights from sandstone petrography and detrital apatite fission-track thermochronology in the intramontaneAngastaco basin, NW Argentina: Basin Research, v. 18, p. 1-26, doi: 10.1111/j.1365-2117.2006.00283.x.
- Cunningham, C. G., Austin, G. W., Naeser, C. W., Rye, R. O., Ballantyne, G. H., Stamm, R. G., et al. (2004). Formation of a paleothermal anomaly and disseminated gold deposits associated with the Bingham Canyon porphyry Cu-Au-Mo system, Utah. Economic Geology and the Bulletin of the Society of Economic Geologists, 99(4), 789-806.
- Dadson, S. J., Hovius, N., Chen, H., Dade, W. B., Hsieh, M.-L., Willett, S. D., et al. (2003). Links between erosion, runoff variability and seismicity in the Taiwan Orogen. Nature, 426(6967), 648-651.
- Daishi, M., Inoue, M., & Furuyama, K. (2003). Radiometric ages of late Cenozoic volcanic rocks and implications for volcanic activity in western area of Oita, Kyushu, Japan. Journal of Geosciences, Osaka City University, 46, 25-45.
- Danišík, M., Dunkl, I., Puti, M., Frisch, W. and Krá, J. (2004): Tertiary burial and exhumation history of basement highs along the NW margin of the Pannonian Basin an apatite fission track study. Austrian Journal of Earth Sciences, 95/96, 60-70, Vienna.

- Danišík, M. (2005): Cooling history and relief evolution of Corsica (France) as constrained by fission track and (U-Th)/He thermochronology. Tübinger Geowissenschaftliche Arbeiten, Reihe A, 72, 1-130.
- Danišík, M., Dunkl, I., Puti M., Frisch, W. & Král, J. (2004): Tertiary burial and exhumation history of basement highs along the NW margin of the Pannonian Basin - an apatite fission track study. Austrian Journal of Earth Sciences, Vienna, 95/96, 60-70.
- Davidson, J. P., Hassanzadeh, J., Berzins, R., Stockli, D. F., Bashukooh, B., Turrin, B., et al. (2004). The geology of Damavand Volcano, Alborz Mountains, northern Iran. Geological Society of America Bulletin, 116(1-2), 16-29.
- Day, J. M. D., Hilton, D. R., Pearson, D. G., Macpherson, C. G., Kjarsgaard, B. A., & Janney, P. E. (2005). Absence of a high time-integrated (super 3) He/(U+Th) source in the mantle beneath continents. Geology, 33(9), 733-736.
- DeCelles, P.G., Gehrels, G.E., Najman, Y., Martin, A.J., Carter, A., and Garzanti, E. (2004). Detrital geochronology and geochemistry of Cretaceous—Early Miocene strata of Nepal: Implications for timing and diachroneity of initial Himalayan orogenesis. Earth and Planetary Science Letters, v227, p313-330.
- Dempster, T. J., & Persano, C. (2006). Low-temperature thermochronology; resolving geotherm shapes or denudation histories? Geology, 34(2), 73-76.
- Donelick, R. A., O'Sullivan, P. B., & Ketcham, R. A. (2005). Apatite fission-track analysis. Reviews in Mineralogy and Geochemistry, 58(1), 49-94.
- Ducea, M. N., Valencia, V. A., Shoemaker, S., Reiners, P. W., DeCelles, P. G., Campa, M. F., et al. (2004). Rates of sediment recycling beneath the Acapulco Trench; constraints from (U-Th)/He thermochronology. Journal of Geophysical Research, 109, no.B9, 11.
- Ducea, M., House, M. A., & Kidder, S. (2003). Late Cenozoic denudation and uplift rates in the Santa Lucia Mountains, California. Geology, 31(2), 139-142.
- Dunai, T. J. (2005). Forward modeling and interpretation of (U-Th)/He ages. Reviews in Mineralogy and Geochemistry, 58(1), 259-274.
- Dunkl, I., Frisch, W., & Grundmann, G. (2003). Zircon fission track thermochronology of the southeastern part of the Tauern Window and the adjacent Austroalpine margin, Eastern Alps. Eclogae Geologicae Helvetiae, 96(2), 209-217.
- Dunkl, I., Kuhlemann, J., Reinecker, J. and Frisch, W. (2005): Cenozoic relief evolution of the Eastern Alps constraints from apatite fission track age-provenance of Neogene intramontane sediments. Austrian Journal of Earth Sciences, 98, 92-105.
- Eby, G. N. (2004) Petrology, geochronology, mineralogy, and geochemistry of the Beemerville alkaline complex, northern New Jersey. In Puffer, J. H. and Volkert, R. A. (eds.) Neoproterozoic, Paleozoic, and Mesozoic Intrusive Rocks of Northern New Jersey and Southeastern New York. Twenty-First Annual Meeting Geological Association of New Jersey, Mahwah, NJ, pp. 52-68.
- Ehlers, T.A., Chaudri, T., Kumar, S., Fuller, C.W., Willett, S.D., Ketcham, R.A., Brandon, M.T., Belton, D.X., Kohn, B.P., Gleadow, A.J.W., Dunai, T.J., Fu, F.Q. (2005). Computational tools for low-temperature thermochronometer interpretation. In: Reiners, P.W. and Ehlers, T.A. (Eds.), Low Temperature

Thermochronology: Techniques, Interpretations, and Applications. Reviews in Mineralogy and Geochemistry, vol. 58, pp. 589-622.

- Emmel, B., Jacobs, J., Kastowski, M., & Graser, G. (2006). Phanerozoic upper crustal tectono-thermal development of basement rocks from central Madagascar; an integrated fission-track and structural study. Tectonophysics, 412(1-2), 61-86.
- Emmerich, A., Glasmacher, U. A., Bauer, F., Bechstadt, T., & Zuehlke, R. (2005). Meso-/Cenozoic basin and carbonate platform development in the SW-Dolomites unraveled by basin modelling and apatite FT analysis; Rosengarten and Latemar (northern Italy). Sedimentary Geology, 175(1-4), 415-438.
- Enkelmann, E., & Jonckheere, R. (2003). Correction factors for systematic errors related to the track counts in fission-track dating with the external detector method. Radiation Measurements, 36(1-6), 351-356.
- Enkelmann, E., Jonckheere, R., & Wauschkuhn, B. (2005). Independent fission-track ages (phi -ages) of proposed and accepted apatite age standards and a comparison of phi -, Z-, , zeta - and zeta (sub 0) -ages; implications for method calibration. Chemical Geology, 222(3-4), 232-248.
- Enkelmann, E., Jonckheere, R., Ratschbacher, L. (2005). Absolute measurement of uranium concentration on thick samples using fission-track detectors. Nuclear Instruments and Methods B. 229, 489-498.
- Enkelmann, E., Jonckheere, R., Ratschbacher, L. (2005). The effects of radiation damage accumulation and annealing for fission-track dating of titanite. Nuclear Instruments and Methods B. 227, 581-590.
- Enkelmann, E., Ratschbacher, L., Gloaguen, R., Nestler, R., Fleischer, M., Jonckheere, R., Hacker, B.R., Yue Qiao Zhang. (2006). Is Tibetan lower crustal flow diverging around the Sichuan basin? Denudation and deformation in the Qinling. GSA Bulletin 118, 651–671; doi: 10.1130/B25805.1
- Esteban, J. J., Sanchez-Rodriguez, L., Seward, D., Cuevas, J., & Tubia, J. M. (2004). The late thermal history of the Ronda area, southern Spain. Tectonophysics, 389(1-2), 81-92.
- Evans, N. J., Wilson, N. S. F., Cline, J. S., McInnes, B. I. A., & Byrne, J. (2005). Fluorite (U-Th)/He thermochronology; constraints on the low temperature history of Yucca Mountain, Nevada. Applied Geochemistry, 20(6), 1099-1105.
- Fellin M.G., Picotti V. & Zattin M. (2005) Neogene to Quaternary multiple rifting and inversion in north-eastern Corsica: retreat and collision in the Western Mediterranean. Tectonics, 24, TC1011.
- Fellin M.G., Vance J., Garver J. & Zattin M. The thermal evolution of Corsica as recorded by zircon fission-tracks. Tectonophysics, in press.
- Fellin, M. G., Zattin, M., Picotti, V., Reiners, P. W., & Nicolescu, S. (2005). Relief evolution in northern Corsica (western Mediterranean); constraints on uplift and erosion on longterm and short-term timescales. Journal of Geophysical Research, 110, no.F1, 16.
- Ferrari, L., Petrone, C. M., Francalanci, L., Tagami, T., Eguchi, M., Conticelli, S., Manetti, P. and Salgado, S. V. (2003). Geology of the San Pedro-Ceboruco graben, western trans-Mexican volcanic belt. Revista Mexicana de Ciencias Geologicas, 20, 165-181.

- Ferrari, L., Tagami, T., Eguchi, M., Orozco-Esquivel, M.T., Petrone, C.M., Jacobo-Albarr, J., and Lez-Martez, M. (2005). Geology, geochronology and tectonic setting of late Cenozoic volcanism along the southwestern Gulf of Mexico: the Eastern Alkaline Province revisited. J. Volcanol. Geotherm. Res., 146, 284-306.
- Filip, J., & Suchy, V. (2004). Thermal and tectonic history of the Barrandian lower Paleozoic, Czech Republic; is there a fission-track evidence for Carboniferous-Permian overburden and pre-Westphalian alpinotype thrusting? Bulletin of Geosciences (Praha), 79(2), 107-112.
- Fleischer, R. L., & Anonymous. (2003). Etching of recoil tracks in solids. Geochimica et Cosmochimica Acta, 67(24), 4769-4774.
- Fleischer, R.L. (2003). "Recoil Tracks in Solids,", Geochimica et Cosmochimica Acta 67 (#24) 4769-4774.
- Fleischer, R.L. (2004). "Does Maskelynite Imply Anything About Impact Craters?", Meteoritics and Planetary Science 39, 12, 2055
- Fleischer, R.L. (2004). "Fission Tracks in Solids Production Mechanisms and Natural Origins", J. Materials Science 39, 3901-3911.
- Fleischer, R.L. (2005) "Nuclear Tracks and Nanostructures", Chapter 10 in Engineering Thin Films and Nanostructures with Ion Beams, Emile Knystautas, ed., CRC, Marcel Dekker, New York, pp.491-522.
- Fleischer, R.L. (2005). "Are tracks caused by thermal spikes?" Int. Nuclear Track Soc. Newsletter 5 (#2), 2-3.
- Fleischer, R.L. (2005). "The Distribution of Boron in AIRu: Effect on Ductility and Toughness", Acta Materialia. 53, 2623-2627
- Fleischer, R.L., Sekyung Chang, Jeremy Farrell, Rachel C. Herrmann, Jonathan MacDonald, Marek Zalesky, and Robert H. Doremus (2006). "Etched Tracks and Serendipitous Dosimetry," Radiation Protection Dosimetry
- Fleischer, R.L., Sekyung Chang, Jeremy Farrell, Stephen A. Hadley, Rachel C. Herrmann, Jonathan MacDonald, Nicholas R. Meyer, Marek Zalesky, and Robert H. Doremus (2005).
  "Studies in Nuclear Tracks at Union College," Acta Physica et Chimica Debricina 38-39, 141-152
- Flowers, R. M., Pringle, M. S., Mahan, K. H., Bowring, S. A., Williams, M. L., Hodges, K. V., et al. (2005). Temporal constraints on the juxtaposition and exhumation of deep crustal domains, east Athabasca region, western Canadian Shield. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 291.
- Flynn, J. J., Kowallis, B. J., Nunez, C., Carranza-Castaneda, O., Miller, W. E., Swisher, C. C., III, et al. (2005). Geochronology of Hemphillian-Blancan aged strata, Guanajuato, Mexico, and implications for timing of the great American biotic interchange. Journal of Geology, 113(3), 287-307.
- Foeken, J. P. T., Dunai, T. J., Bertotti, G., & Andriessen, P. A. M. (2003). Late Miocene to present exhumation in the Ligurian Alps (Southwest Alps) with evidence for accelerated denudation during the Messinian salinity crisis. Geology, 31(9), 797-800.
- Fuegenschuh, B., & Schmid, S. M. (2003). Late stages of deformation and exhumation of an orogen constrained by fission-track data; a case study in the Western Alps. Geological Society of America Bulletin, 115(11), 1425-1440.

- Fuegenschuh, B., & Schmid, S. M. (2005). Age and significance of core complex formation in a very curved orogen; evidence from fission track studies in the South Carpathians (Romania). Tectonophysics, 404(1-2), 33-53.
- Fuegenschuh, B., Froitzheim, N., Capdevila, R., & Boillot, G. (2003). Offshore granulites from the Bay of Biscay margins; fission tracks constrain a Proterozoic to Tertiary thermal history. Terra Nova, 15(5), 337-342.
- Gallagher, K. Uplift, Denudation and their causes and constraints on geological timescales, invited book chapter for Principles of Phanerozoic Geology, Roberts, D. and Bally, A. Editors. in press.
- Gallagher, K., Stephenson, J., Brown R., Holmes, C. and Ballester, P. Exploiting 3-D spatial sampling in modeling of thermochronological data, Reviews in Mineralogy and Geochemistry, special volume on Low Temperature Thermochronology, vol 58, 375-388.
- Gallagher, K., Stephenson, J., Brown, R., Holmes, C., & Fitzgerald, P. (2005). Low temperature thermochronology and modeling strategies for multiple samples; 1, Vertical profiles. Earth and Planetary Science Letters, 237(1-2), 193-208.
- Garver, J. I., Reiners, P. W., Walker, L. J., Ramage, J. M., & Perry, S. E. (2005). Implications for timing of Andean uplift from thermal resetting of radiation-damaged zircon in the Cordillera Huayhuash, northern Peru. Journal of Geology, 113(2), 117-138.
- Glasmacher, U. A., Matenaar, I., Bauer, W., & Puchkov, V. N. (2004). Diagenesis and incipient metamorphism in the western fold-and-thrust belt, SW Urals, Russia. International Journal of Earth Sciences, 93(3), 361-383.
- Gleadow, A. J. W., Raza, A., Kohn, B. P., Spencer, S. A., & Anonymous. (2005). The potential of monazite for fissiontrack dating. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 21.
- Glodny, J., Austrheim, H., Molina, J. F., Rusin, A. I., & Seward, D. (2003). Rb/Sr record of fluid-rock interaction in eclogites; the Marun-Keu Complex, Polar Urals, Russia. Geochimica et Cosmochimica Acta, 67(22), 4353-4371.
- Glodny, J., Lohrmann, J., Echtler, H., Graefe, K., Seifert, W., Collao, S., et al. (2005). Internal dynamics of a paleoaccretionary wedge; insights from combined isotope tectonochronology and sandbox modelling of the southcentral Chilean forearc. Earth and Planetary Science Letters, 231(1-2), 23-39.
- Gomez, E., Jordan, T. E., Allmendinger, R. W., Hegarty, K., & Kelley, S. (2005). Syntectonic Cenozoic sedimentation in the northern middle Magdalena Valley basin of Colombia and implications for exhumation of the Northern Andes. Geological Society of America Bulletin, 117(5-6), 547-569.
- Green, P. F., Duddy, I. R., & Hegarty, K. A. (2005). Compositional and structural control of fission track annealing in apatite; discussion [modified]. Chemical Geology, 214(3-4), 351-358.
- Grist, A. M., & Zentilli, M. (2003). Post-Paleocene cooling in the southern Canadian Atlantic region; evidence from apatite fission track models. Canadian Journal of Earth Sciences = Revue Canadienne des Sciences de la Terre, 40(9), 1279-1297.
- Grist, A. M., & Zentilli, M. (2005). The thermal history of the Nares Strait, Kane Basin, and Smith Sound region in Canada

and Greenland; constraints from apatite fission-track and (U-Th-Sm)/He dating. Canadian Journal of Earth Sciences = Revue Canadienne des Sciences de la Terre, 42(9), 1547-1569.

- Gunnell, Y., Gallagher, K., Carter, A., Widdowson, M., & Hurford, A. J. (2003). Denudation history of the continental margin of western peninsular India since the early Mesozoic; reconciling apatite fission-track data with geomorphology. Earth and Planetary Science Letters, 215(1-2), 187-201.
- Gurov, E. P., & Koeberl, C. (2004). Shocked rocks and impact glasses from the El'gygytgyn impact structure, Russia. Meteoritics & Planetary Science, 39(9), 1459-1508.
- Harrison, T. M., & Zeitler, P. K. (2005). Fundamentals of noble gas thermochronometry. Reviews in Mineralogy and Geochemistry, 58(1), 123-149.
- Harrison, T. M., Grove, M., Lovera, O. M., & Zeitler, P. K. (2005). Continuous thermal histories from inversion of closure profiles. Reviews in Mineralogy and Geochemistry, 58(1), 389-409.
- Hasebe, N., & Hoshino, H. (2003). Igneous rocks emplacement and exhumation of sedimentary basement; fission track age determination on the Osuzuyama volcano-plutonic complex and surrounding rocks, Miyazaki Prefecture, southwest Japan. Geochemical Journal, 37(5), 537-543.
- Hasebe, N., & Watanabe, H. (2004). Heat influx and exhumation of the Shimanto accretionary complex; Miocene fission track ages from the Kii Peninsula, southwest Japan. The Island Arc, 13(4), 533-543.
- Hasebe, N., Barbarand, J., Jarvis, K., Carter, A., & Hurford, A. J. (2004). Apatite fission-track chronometry using laser ablation ICP-MS. Chemical Geology, 207(3-4), 135-145.
- Heffern, E. L., & Coates, D. A. (2004). Geologic history of natural coal-bed fires, Powder River basin, USA. International Journal of Coal Geology, 59(1-2), 25-47.
- Heim, J. A., Vasconcelos, P. M., Shuster, D. L., Farley, K. A., & Broadbent, G. (2006). Dating paleochannel iron ore by (U-Th)/He analysis of supergene goethite, Hamersley Province, Australia. Geology, 34(3), 173-176.
- Hejl, E., Sekyra, G., & Friedl, G. (2003). Fission-track dating of the south-eastern Bohemian Massif (Waldviertel, Austria); thermochronology and long-term erosion. International Journal of Earth Sciences, 92(5), 677-690.
- Hendriks, B. W. H., & Redfield, T. F. (2005). Apatite fission track and (U-Th)/He data from Fennoscandia; an example of underestimation of fission track annealing in apatite. Earth and Planetary Science Letters, 236(1-2), 443-458.
- Hendriks, B. W. H., Redfield, T. F., & Anonymous. (2005). Evidence for underestimation of long-term FT annealing in apatite from natural FT and (U-Th)/He data. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 298.
- Herwegh, M., & Pfiffner, O. A. (2005). Tectono-metamorphic evolution of a nappe stack; a case study of the Swiss Alps. Tectonophysics, 404(1-2), 55-76.
- Hodges, K. V., Ruhl, K., Wobus, C., Boyce, J. W., & Anonymous. (2005). Detrital mineral thermochronology in active fluvial systems and the evolution of modern orogenic landscapes. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 296.
- Holford, S. P., Green, P. F., & Turner, J. P. (2005). Palaeothermal and compaction studies in the Mochras

borehole (NW Wales) reveal Early Cretaceous and Neogene exhumation and argue against regional Palaeogene uplift in the southern Irish Sea. Journal of the Geological Society of London, 162(5), 829-840.

- Honda, M., Nutman, A. P., Bennett, V. C., & Yatsevich, I. (2004). Radiogenic, nucleogenic and fissiogenic noble gas compositions in early Archaean magmatic zircons from Greenland. Geochemical Journal, 38(3), 265-269.
- Hourigan, J. K., Reiners, P. W., & Brandon, M. T. (2005). U/Th zonation-dependent alpha-ejection in (U-Th)/He chronometry. Geochimica et Cosmochimica Acta, 69(13), 3349-3365.
- Hourigan, J. K., Reiners, P. W., Brandon, M. T., Plank, T., & Anonymous. (2005). Laser-ablation ICP-MS zonationdependent alpha -ejection correction of zircons in (U-Th)/He chronometry. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 302.
- House, M. A., Kelley, S. A., & Roy, M. (2003). Refining the footwall cooling history of a rift flank uplift, Rio Grande Rift, New Mexico. Tectonics, 22, no.5, 18.
- Hu, S., Kohn, B.P., Raza, A., Wang, J., and Gleadow, A.J.W. (2006). Cretaceous and Cenozoic cooling history of the ultrahigh-pressure rocks across the Tongbai-Dabie orogen, central China, from apatite fission track thermochronology. Tectonophysics, (in press).
- Huang, C.-Y., Yuan, P. B., & Tsao, S.-J. (2006). Temporal and spatial records of active arc-continent collision in Taiwan; a synthesis. Geological Society of America Bulletin, 118(3-4), 274-288.
- Hurford, A. J., Barbarand, J., & Carter, A. (2005). Compositional and structural control of fission-track annealing in apatite; reply [modified]. Chemical Geology, 214(3-4), 359-361.
- Issler, D.R. and Grist, A.M. (2005). Apatite fission track age, length and kinetic parameter (CI, Dpar) data for the Northrock et al. East MacKay I-77 well, Mackenzie Valley, Northwest Territories. Geological Survey of Canada, Open File 4889. (CD-ROM).
- Issler, D.R., Grist, A.M. and Stasiuk, L.D. (2005). Post-Early Devonian thermal constraints on hydrocarbon source rock maturation in the Keele Tectonic Zone, Tulita area, NWT, Canada, from multi-kinetic apatite fission track thermochronology, vitrinite reflectance and shale compaction. Bulletin of Canadian Petroleum Geology, v. 53, no. 4, p. 405-431.
- Ito, H. (2004). Apatite fission-track dating of fault-related rocks along the Nojima and Kusumoto Faults and its tectonic significance. Geophysical Research Letters, 31, no.21, 24.
- Japsen, P., Green, P. F., & Chalmers, J. A. (2005). Separation of Palaeogene and Neogene uplift on Nuussuaq, West Greenland. Journal of the Geological Society of London, 162(2), 299-314.
- Jasra, A., Stephens, D.A. Gallagher, K. and Holmes, C.C. (2006) Analysis of geochronological data with measurement error using Bayesian mixtures, Mathematical Geology, 38(3).
- John, B. E., Foster, D. A., Murphy, J. M., Cheadle, M. J., Baines, A. G., Fanning, C. M., et al. (2004). Determining the cooling history of in situ lower oceanic crust; Atlantis Bank, SW Indian Ridge. Earth and Planetary Science Letters, 222(1), 145-160.
- Jonckheere, R., Enkelmann, E., Stübner, K. (2005). On the geometries of etched fission and alpha-recoil tracks with

reference to models of track revelation in minerals. Radiation Measurements 39, 577-583.

- Karg, H., Carter, A., Brix, M. R., & Littke, R. (2005). Late- and post-Variscan cooling and exhumation history of the northern Rhenish Massif and the southern Ruhr Basin; new constraints from fission-track analysis. Geologische Rundschau = International Journal of Earth Sciences, 94(2), 180-192.
- Keller, L. M., Fuegenschuh, B., Hess, M., Schneider, B., & Schmid, S. M. (2006). Simplon fault zone in the Western and Central Alps; mechanism of Neogene faulting and folding revisited. Geology, 34(4), 317-320.
- Keller, L. M., Hess, M., Fuegenschuh, B., & Schmid, S. M. (2005). Structural and metamorphic evolution of the Camughera-Moncucco, Antrona and Monte Rosa Units southwest of the Simplon Line, Western Alps. Eclogae Geologicae Helvetiae, 98(1), 19-49.
- Kershaw, A. P., D'Costa, D. M., Tibby, J., Wagstaff, B. E., & Heijnis, H. (2004). The last million years around Lake Keilambete, western Victoria. Proceedings of the Royal Society of Victoria, 116(1), 93-104.
- Ketcham, R. A. (2005). Forward and inverse modeling of lowtemperature thermochronometry data. Reviews in Mineralogy and Geochemistry, 58(1), 275-314.
- Ketcham, R.A. (2003). Observations on the relationship between crystallographic orientation and biasing in apatite fission-track measurements. American Mineralogist, 88, 817-829.
- Ketcham, R.A. (2005). Forward and inverse modeling of lowtemperature thermochronometry data. Reviews in Mineralogy and Geochemistry, 58, 275-314.
- Ketcham, R.A. (2005). The role of crystallographic angle in characterizing and modeling apatite fission-track length data. Radiation Measurements, 39, 595-601.
- Kissová, D., Dunkl, Daniíkl., Plaienka, D., Frisch, W. and Marschalko, R. (2005): The Pieninic exotic cordillera (Andrusov Ridge) revisited: new zircon FT ages of granitic pebbles from Cretaceous flysch conglomerates of the Pieniny Klippen Belt (Western Carpathians, Slovakia). Slovak Geol. Mag., 11, 17-28.
- Kohn B.P., O'Sullivan P.B., Gleadow A.J.W., Brown R.W., Gallagher K. and Foster D.A. (2003). Fission track thermotectonic imaging and denudation history of Tasmania: Reply to Discussion by D.E. Leaman. Australian Journal of Earth Sciences, 50: 646-650.
- Kohn, B. P., Belton, D. X., Brown, R. W., Gleadow, A. J. W., Green, P. F., & Lovering, J. F. (2003). Experimental evidence for the pressure dependence of fission track annealing in apatite; discussion [modified]. Earth and Planetary Science Letters, 215(1-2), 299-306.
- Kohn, B.P., Gleadow, A.J.W., Brown, R.W., Gallagher, K., Lorencak, M., Noble, W.P. (2005). Visualising Thermotectonic and Denudation Histories Using Apatite Fission Track Thermochronology. In: Reiners, P.W. and Ehlers, T.A. (Eds.), Low Temperature Thermochronology: Techniques, Interpretations, and Applications. Reviews in Mineralogy and Geochemistry, 58, pp. 527-565.
- Kondo, H., Tanaka, K., Mizuochi, Y., & Ninomiya, A. (2004). Long-term changes in distribution and chemistry of middle Miocene to Quaternary volcanism in the Chokai-Kurikoma area across the northeast Japan Arc. The Island Arc, 13(1), 18-46.

- Kuhlemann, J., Frisch, W. Székely, B., Dunkl, I., Daniík, M. and Krumrei, I. (2005): Würmian maximum glaciation in Corsica. Austrian Journal of Earth Sciences, 97, 68-81.
- Kuhlemann, J., Frisch, W., Dunkl, I., Kázmér, M. and Schmiedl, G. (2004): Miocene siliciclastic deposits of Naxos Island: Geodynamic and environmental implications for the evolution of the southern Aegean Sea (Greece), in Bernet, M., and Spiegel, C., eds., Detrital thermochronology - Provenance analysis, exhumation, and landscape evolution of mountain belts: Boulder, Colorado, Geological Society of America Special Paper 378, p. 5165.
- Kuhlemann, J., I. Dunkl, A. Brügel, C. Spiegel and W. Frisch (2006). From source terrains of the Eastern Alps to the Molasse Basin: Detrital record of non-steady-state exhumation. Tectonophysics 413, 301-316.
- Kumerics, C., Ring, U., Brichau, S., Glodny, J., & Monie, P. (2005). The extensional Messaria shear zone and associated brittle detachment faults, Aegean Sea, Greece. Journal of the Geological Society of London, 162(4), 701-721.
- Kunk, M. J., Wintsch, R. P., Naeser, C. W., Naeser, N. D., Southworth, C. S., Drake, A. A., Jr., et al. (2005). Contrasting tectonothermal domains and faulting in the Potomac Terrane, Virginia-Maryland; discrimination by (super 40) Ar/ (super 39) Ar and fission-track thermochronology. Geological Society of America Bulletin, 117(9-10), 1347-1366.
- Lal, D., Harris, N. B. W., Sharma, K. K., Gu, Z., Ding, L., Liu, T., et al. (2004). Erosion history of the Tibetan Plateau since the last interglacial; constraints from the first studies of cosmogenic (super 10) Be from Tibetan bedrock. Earth and Planetary Science Letters, 217(1-2), 33-42.
- Laurenzi, M. A., Bigazzi, G., Balestrieri, M. L., & Bouska, V. (2003). (super 40) Ar/ (super 39) Ar laser probe dating of the Central European tektite-producing impact event. Meteoritics & Planetary Science, 38(6), 887-893.
- Leaman, D. E., Kohn, B. P., O'Sullivan, P. B., Gleadow, A. J. W., Brown, R. W., Gallagher, K., et al. (2003). Shaping the Australian crust over the last 300 million years; insights from fission track thermotectonic imaging and denudation studies of key terranes; discussion and reply. Australian Journal of Earth Sciences, 50(4), 645-650.
- Leloup, P. H., Arnaud, N., Sobel, E. R. and Lacassin, R. (2005), Timing and mode of exhumation of the highest alpine range, the Mt Blanc massif: Tectonics, v. 24 TC4002, doi:10.1029/2004TC001676.
- Li, X., Wang, Y., Tan, K., & Peng, T. (2005). Meso-Cenozoic uplifting and exhumation on Yunkaidashan; evidence from fission track thermochronology. Chinese Science Bulletin, 50(9), 903-909.
- Lim, H. S., & Lee, Y. I. (2005). Cooling history of the Upper Cretaceous Palgongsan Granite, Gyeongsang Basin, SE Korea and its tectonic implication for uplift on the active continental margin. Tectonophysics, 403(1-4), 151-165.
- Lisker, F. (2004). The evolution of the geothermal gradient from Lambert Graben and Mahanadi Basin; a contribution to the Indo-Antarctic Rift debate. Gondwana Research, 7(2), 363-373.
- Lisker, F., & Olesch, M. (2003). Long-term landscape evolution of George V Land as indicated by fission track data. Terra Antartica, 10(3), 249-256.
- Lisker, F., Brown, R., & Fabel, D. (2003). Denudational and thermal history along a transect across the Lambert Graben,

northern Prince Charles Mountains, Antarctica, derived from apatite fission track thermochronology. Tectonics, 22, no.5, 14.

- Little, T. A., Cox, S., Vry, J. K., & Batt, G. (2005). Variations in exhumation level and uplift rate along the oblique-slip Alpine Fault, central Southern Alps, New Zealand. Geological Society of America Bulletin, 117(5-6), 707-723.
- Lo, D., R. L. Fleischer, E. A. Albert, and J.G. Arnason, J. Environ (2006). "Size and Location of Depleted Uranium Grains in Reservoir Sediments", Radioactivity.
- Locke, B. D., & Garver, J. I. (2005). Thermal evolution of the eastern Serrania del Interior foreland fold and thrust belt, northeastern Venezuela, based on apatite fission-track analyses. Special Paper - Geological Society of America, 394, 315-328.
- Lorencak, M., Kohn, B. P., Osadetz, K. G., & Gleadow, A. J. W. (2004). Combined apatite fission track and (U/Th)/He thermochronometry in a slowly cooled terrane; results from a 3440-m-deep drill hole in the southern Canadian Shield. Earth and Planetary Science Letters, 227(1-2), 87-104.
- MacPhee, D., Bowring, S. A., Reiners, P. W., & Anonymous. (2005). Combined (U-Th)/He and U-Pb thermochronometry of rift-flank exhumation in east-central Africa. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 304.
- Malusa M.G., Philippot P., Zattin M. & Martin S. (2006) Late stages of exhumation constrained by structural, fluid inclusion and fission track analyses (Sesia-Lanzo unit, Western European Alps). Earth and Planetary Science Letters, 243, 565-580.
- Malusa M.G., Zattin M., Ando S., Garzanti E. & Vezzoli G. (2005) - Detrital apatite fission-track ages from the Po sands: implications for the erosional evolution of the orogenic source areas. Geoacta, 4, 57-65.
- Malusa, M. G., Polino, R., Zattin, M., Bigazzi, G., Martin, S., & Piana, F. (2005). Miocene to present differential exhumation in the Western Alps; insights from fission track thermochronology. Tectonics, 24, no.3, 23.
- Marcolini, F., Bigazzi, G., Bonadonna, F. P., Centamore, E., Cioni, R., & Zanchetta, G. (2003). Tephrochronology and tephrostratigraphy of two Pleistocene continental fossiliferous successions from central Italy. JQS. Journal of Quaternary Science, 18(6), 545-556.
- Mazzoli S., Aldega L., Corrado S., Invernizzi C. & Zattin M. -Pliocene-Quaternary tectonic exhumation of 'Apulian' carbonate platform reservoir rocks, Monte Alpi, southern Apennines, Italy. GSA Sp. Publ., in press.
- McDowell, F. W., McIntosh, W. C., & Farley, K. A. (2005). A precise (super 40) Ar/ (super 39) Ar reference age for the Durango apatite (U-Th)/He and fission-track dating standard. Chemical Geology, 214(3-4), 249-263.
- McInnes, B. I. A., Evans, N. J., Fu, F. Q., & Garwin, S. (2005). Application of thermochronology to hydrothermal ore deposits. Reviews in Mineralogy and Geochemistry, 58(1), 467-498.
- Meibom, A., Anderson, D. L., Sleep, N. H., Frei, R., Chamberlain, C. P., Hren, M. T., et al. (2003). Are high (super 3) He/ (super 4) He ratios in oceanic basalts an indicator of deep-mantle plume components? Earth and Planetary Science Letters, 208(3-4), 197-204.

- Metzler, K. (2004). Formation of accretionary dust mantles in the solar nebula; evidence from preirradiated olivines in CM chondrites. Meteoritics & Planetary Science, 39(8), 1307-1319.
- Min, K. (2005). Low-temperature thermochronometry of meteorites. Reviews in Mineralogy and Geochemistry, 58(1), 567-588.
- Min, K., Farley, K. A., Renne, P. R., & Marti, K. (2003). Single grain (U-Th)/He ages from phosphates in Acapulco Meteorite and implications for thermal history. Earth and Planetary Science Letters, 209(3-4), 323-336.
- Min, K., Reiners, P. W., Nicolescu, S., & Greenwood, J. P. (2004). Age and temperature of shock metamorphism of Martian meteorite Los Angeles from (U-Th)/He thermochronometry. Geology, 32(8), 677-680.
- Mitchell, S. G., & Reiners, P. W. (2003). Influence of wildfires on apatite and zircon (U-Th)/He ages. Geology, 31(12), 1025-1028.
- Mochizuki, N., Tsunakawa, H., Shibuya, H., Tagami, T., Ozawa, A., Cassidy, J. and Smith, I. E. M. (2004). K-Ar ages of the Auckland geomagnetic excursions. Earth Planets Space, 56, 283-288.
- Morgan, L. A., & McIntosh, W. C. (2005). Timing and development of the Heise volcanic field, Snake River plain, Idaho, western USA. Geological Society of America Bulletin, 117(3-4), 288-306.
- Morley, C. K. (2004). Nested strike-slip duplexes, and other evidence for Late Cretaceous-Palaeogene transpressional tectonics before and during India-Eurasia collision, in Thailand, Myanmar and Malaysia. Journal of the Geological Society of London, 161(5), 799-812.
- Mortimer, N., & Cooper, A. F. (2004). U-Pb and Sm-Nd ages from the Alpine Schist, New Zealand. New Zealand Journal of Geology and Geophysics, 47(1), 21-28.
- Mortimer, N., Herzer, R. H., Walker, N. W., Calvert, A. T., Seward, D., & Chaproniere, G. C. H. (2003). Cavalli Seamount, Northland Plateau, SW Pacific Ocean; a Miocene metamorphic core complex? Journal of the Geological Society of London, 160(6), 971-983.
- Moser, F., Hann, H. P., Dunkl, I., & Frisch, W. (2005). Exhumation and relief history of the Southern Carpathians (Romania) as evaluated from apatite fission track chronology in crystalline basement and intramontane sedimentary rocks. Geologische Rundschau = International Journal of Earth Sciences, 94(2), 218-230.
- Murakami, M., & Tagami, T. (2004). Dating pseudotachylyte of the Nojima Fault using the zircon fission-track method. Geophysical Research Letters, 31, no.12, 14.
- Murakami, M., Yamada, R. and Tagami, T. (2006). Short-term annealing characteristics of spontaneous fission tracks in zircon: A qualitative description. Chem. Geol., 227, 214-222.
- Naeth, J., Di Primio, R., Horsfield, B., Schaefer, R. G., Shannon, P. M., Bailey, W. R., et al. (2005). Hydrocarbon seepage and carbonate mound formation; a basin modelling study from the Porcupine Basin (offshore Ireland). Journal of Petroleum Geology, 28(2), 147-166.
- Najman, Y., Carter, A., Oliver, G., & Garzanti, E. (2005). Provenance of Eocene foreland basin sediments, Nepal; constraints to the timing and diachroneity of early Himalayan orogenesis. Geology, 33(4), 309-312.

- Nasdala, L., Reiners, P. W., Garver, J. I., Kennedy, A. K., Stern, R. A., Balan, E., et al. (2004). Incomplete retention of radiation damage in zircon from Sri Lanka. American Mineralogist, 89(1), 219-231.
- Nicolescu, S., Reiners, P. W., & Anonymous. (2005). (U-Th)/He dating of epidote and andradite garnet. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 26.
- Nomade, S., Renne, P. R., Vogel, N., Deino, A. L., Sharp, W. D., Becker, T. A., et al. (2005). Alder Creek sanidine (ACs-2); a Quaternary (super 40) Ar/ (super 39) Ar dating standard tied to the Cobb Mountain geomagnetic event. Chemical Geology, 218(3-4), 315-338.
- Ohtani, T., Shigematsu, N., Fujimoto, K., Tomita, T., & Iwano, H. (2004). Geochronological constraint on the brittle-plastic deformation along the Hatagawa fault zone, NE Japan. Earth, Planets and Space, 56(12), 1201-1207.
- Ortega-Rivera, A. (2003). Geochronological constraints on the tectonic history of the Peninsular Ranges Batholith of Alta and Baja California; tectonic implications for western Mexico. Special Paper Geological Society of America, 374, 297-335.
- Osadetz K.G., Kohn B.P., Feinstein S. and Price R. (2004). Foreland belt thermal history using apatite fission track thermochronology: implications for Lewis Thrust and Flathead Fault in the Southern Canadian Cordilleran petroleum province. In Swennen, R., Roure, F. and Granath, J. (eds), Chapter 2 - Deformation, fluid flow and reservoir appraisal in foreland fold and thrust belts. AAPG Hedberg Series, no. 1, p. 21-48.
- Osorio, A. M., Iunes, P. J., Bigazzi, G., Hadler N, J. C., Laurenzi, M. A., Norelli, P., et al. (2003). Fission-track dating of Macusanite glasses with plateau and size correction methods. Radiation Measurements, 36(1-6), 407-412.
- Ozawa, A., Tagami, T. and Garcia, M. O. (2005). Unspiked K-Ar dating of the Honolulu rejuvenated and Ko'olau shield volcanism on O'ahu, Hawai'i. Earth Planet. Sci. Lett., 232, 1-11.
- Ozawa, A., Tagami, T. and Kamata, H. (2006). Argon isotopic composition of some Hawai'ian historical lavas. Chem. Geol., 226, 66-72.
- Ozawa, A., Tagami, T. and Sudo, M. (2004). K-Ar geochronology on the temporal change of eruptive stryle in the eastern Izu area, central Japan. The Island Arc, 13, 466-472.
- Ozawa, A., Tagami, T., Listanco, E. L., Arpa, C. B. and Sudo, M. (2004). Initiation and propagation of subduction along the Philippine Trench: evidence from temporal and spatial distribution of volcanoes. J. Asian Earth Sci., 23, 105-111.
- Parada, M. A., Feraud, G., Fuentes, F., Aguirre, L., Morata, D., & Larrondo, P. (2005). Ages and cooling history of the Early Cretaceous Caleu Pluton; testimony of a switch from a rifted to a compressional continental margin in central Chile. Journal of the Geological Society of London, 162(2), 273-287.
- Park, J., Okazaki, R., Kusakabe, M., Nagao, K., & Yoneda, S. (2003). Noble gas and oxygen isotopic compositions of Towada H6-chondrite from Aomori Prefecture, Japan. Geochemical Journal, 37(6), 639-648.
- Perelygin, V.P, I. G. Abdullaev, G. P. Knyazeva, L.I. Kravets, R. I. Petrova, R. Brandt, W. Ensinger, P. Vater, R.L. Fleischer, Yu.V. Bondar, and M. Rebetez (2003). "On Search and Identification of Tracks Due to Short-Lived SHE Nuclei in

Extraterrestrial Crystals," Radiation Measurements 36, 271-279.

- Perelygin, V.P, Yu. V. Bondar, R. Brandt, W. Ensinger, R. L. Fleischer, L. I. Kravets, M. Rebetez, R. Spohr, P. Vater, and S. G. Stetsenko (2003). "On Search and Identification of Relatively Short-Lived Super-Heavy Nuclei (Z≥110) by Fossil Track Studies of Meteoritic and Lunar Olivine Crystals," Phys. of Atomic Nuclei (MAIK Nauka/Interperiodica) 66, no. 8, 1569-1573.
- Perelygin, V.P., Yu.V. Bondar, R. Brandt, W. Ensinger, R.L. Fleischer, L.I. Kravets, M. Rebetez, R. Spohr, P. Vater, S.G. Stetsenko. (2003). "On Search and Identification of Relatively Short-Lived Super-Heavy Nuclei (Z >110) by Fossil Track Studies of Meteoritic and Lunar Olivine Crystals." Yadernaya Phyzika, 66, 1612-1616, in Russian.
- Perez de Armas, J. (2005). Tectonic and thermal history of the western Serrania del Interior foreland fold and thrust belt and Guarico Basin, north-central Venezuela; implications of new apatite fission-track analysis and seismic interpretation. Special Paper Geological Society of America, 394, 271-314.
- Pérez-Arlucea, M.; Carter.; A., Clemente, F.; González, D.; Nombela, M. (2005). Multi-approach, long and short-term denudation rate calculations for the Galician Coast (Ri de Virgo, Spain): AFTD, DEM analysis, river loads and sediment budgets. Journal of Coastal Research, v49, p9-14.
- Persano, C., Stuart, F. M., Barfod, D. N., Bishop, P., Brown, R. W., & Anonymous. (2005). Constraining denudation in Scotland by using a combination of low temperature thermochronometers. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 299.
- Persaud, M., & Pfiffner, O. A. (2004). Active deformation in the eastern Swiss Alps; post-glacial faults, seismicity and surface uplift. Tectonophysics, 385(1-4), 59-84.
- Peyaud JB, Barbarand J, Carter A, Pagel M. (2005). Mid-Cretaceous uplift and erosion on the northern margin of the Ligurian Tethys deduced from thermal history reconstruction Journal of Earth Sciences 94 (3): 462-474
- Pi, T., Sole, J., Taran, Y., Bierlein, F. P., Foster, D. A., Gray, D. R., et al. (2005). (U-Th)/He dating of flourite; application to the La Azul flourspar deposit in the Taxco mining district, Mexico. Mineralium Deposita, 39(8), 976-982.
- Pik, R., Marty, B., Carignan, J., & Lave, J. (2003). Stability of the upper Nile drainage network (Ethiopia) deduced from (U-Th)/He thermochronometry; implications for uplift and erosion of the Afar Plume dome. Earth and Planetary Science Letters, 215(1-2), 73-88.
- Platt, J. P., Argles, T. W., Carter, A., Kelley, S. P., Whitehouse, M. J., & Lonergan, L. (2003). Exhumation of the Ronda Peridotite and its crustal envelope; constraints from thermal modelling of a P-T-time array. Journal of the Geological Society of London, 160(5), 655-676.
- Platt, J. P., Kelley, S. P., Carter, A., & Orozco, M. (2005). Timing of tectonic events in the Alpujarride Complex, Betic Cordillera, southern Spain. Journal of the Geological Society of London, 162(3), 451-462.
- Platt, J.P., Whitehouse, M.J., Kelley, S.P., Carter, A., Hollick, L. (2003). Simultaneous extensional exhumation across the Alboran Basin: Implications for the causes of late orogenic extension. Geology, v 31,p251-254.
- Qiu, N., Zha, M., Wang, X., & Yang, H. (2005). Tectono-thermal evolution of the Junggar Basin, NW China; constraints from R

(sub o) and apatite fission track modelling. Petroleum Geoscience, 11(4), 361-372.

- Raab, M., Brown, R., Gallagher, K., Weber, K., and Gleadow, A. (2005), Denudational and Thermal History of the Early Cretaceous Brandberg and Okenyenya Igneous Complexes on Namibia's Atlantic Passive Margin, Tectonics, 24, TC3006, doi:10.1029/2004TC001688/
- Rahn, M. K., Brandon, M. T., Batt, G. E., & Garver, J. I. (2004). A zero-damage model for fission-track annealing in zircon. American Mineralogist, 89(4), 473-484.
- Ravenhurst, C. E., Roden-Tice, M. K., & Miller, D. S. (2003). Thermal annealing of fission tracks in fluorapatite, chlorapatite, manganoanapatite, and Durango Apatite; experimental results. Canadian Journal of Earth Sciences = Revue Canadienne des Sciences de la Terre, 40(7), 995-1007.
- Re, G. H., Jordan, T. E., & Kelley, S. (2003). Cronologia y paleogeografia del Terciario de la cuenca intermontana de Iglesia septentrional, Andes de San Juan, Argentina. Tertiary geochronology and paleogeography of the northern Iglesia intermontane basin, San Juan, Argentina. Revista de la Asociacion Geologica Argentina, 58(1), 31-48.
- Redfield, T. F., Braathen, A., Gabrielsen, R. H., Osmundsen, P. T., Torsvik, T. H., & Andriessen, P. A. M. (2005). Late Mesozoic to early Cenozoic components of vertical separation across the More-Trondelag fault complex, Norway. Tectonophysics, 395(3-4), 233-249.
- Redfield, T. F., Osmundsen, P. T., & Hendriks, B. W. H. (2005). The role of fault reactivation and growth in the uplift of western Fennoscandia. Journal of the Geological Society of London, 162(6), 1013-1030.
- Reimold, W. U., Buchanan, P. C., Ambrose, D., Koeberl, C., Franchi, I., Lalkhan, C., et al. (2004). Thuathe, a new H4/5 chondrite from Lesotho; history of the fall, petrography, and geochemistry. Meteoritics & Planetary Science, 39(8), 1321-1341.
- Reiners, P. W. (2005). Zircon (U-Th)/He thermochronometry. Reviews in Mineralogy and Geochemistry, 58(1), 151-179.
- Reiners, P. W., Campbell, I. H., Nicolescu, S., Allen, C. M., Hourigan, J. K., Garver, J. I., et al. (2005). (U-Th)/(He-Pb) double dating of detrital zircons. American Journal of Science, 305(4), 259-311.
- Reiners, P. W., Campbell, I. H., Nicolescu, S., Allen, C., Garver, J. I., & Anonymous. (2005). Continental-scale tectonics; zircon He-FT-Pb triple-dating of modern river sediment. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 297.
- Reiners, P. W., Spell, T. L., Nicolescu, S., & Zanetti, K. A. (2004). Zircon (U-Th)/He thermochronometry; He diffusion and comparisons with (super 40) Ar/ (super 39) Ar dating. Geochimica et Cosmochimica Acta, 68(8), 1857-1887.
- Ren, Z., Xiao, H., Liu, L., Zhang, S., Lei, L., Qin, Y., et al. (2005). Determination of Cenozoic uplift-cooling event in Qinshui Basin. Shiyou Yu Tianranqi Dizhi = Oil & Gas Geology, 26(1), 109-113.
- Rocchi, S., Armienti, P., & Di Vincenzo, G. (2005). No plume, no rift magmatism in the West Antarctic Rift. Special Paper Geological Society of America, 388, 435-447.
- Roden-Tice, M. K., & Tice, S. J. (2005). Regional-scale Mid-Jurassic to Late Cretaceous unroofing from the Adirondack Mountains through central New England based on apatite

fission-track and (U-Th)/He thermochronology. Journal of Geology,  $113(5),\,535{-}552.$ 

- Roy, M., Kelley, S., Pazzaglia, F. J., Cather, S., & House, M. A. (2004). Middle Tertiary buoyancy modification and its relationship to rock exhumation, cooling, and subsequent extension at the eastern margin of the Colorado Plateau. Geology, 32(10), 925-928.
- Safran, E. B., Bierman, P. R., Aalto, R., Dunne, T., Whipple, K. X., & Caffee, M. W. (2005). Erosion rates driven by channel network incision in the Bolivian Andes. Earth Surface Processes and Landforms, 30(8), 1007-1024.
- Sano, H., Sherrod, D.R. and Tagami T. (2006). K-Ar dating reveals no rejuvenated-stage in Kaho'olawe, Hawai'i. J. Volcanol. Geotherm. Res., 152, 91-96.
- Schaller, M., Hovius, N., Willett, S. D., Ivy-Ochs, S., Synal, H. A., & Chen, M. C. (2005). Fluvial bedrock incision in the active mountain belt of Taiwan from in situ-produced cosmogenic nuclides. Earth Surface Processes and Landforms, 30(8), 955-971.
- Schultz, P. H., Zarate, M., Hames, B., Koeberl, C., Bunch, T., Storzer, D., et al. (2004). The Quaternary impact record from the Pampas, Argentina. Earth and Planetary Science Letters, 219(3-4), 221-238.
- Sell, I., Poupeau, G., Gonzalez-Casado, J. M., & Lopez-Martinez, J. (2004). A fission track thermochronological study of King George and Livingston Islands, South Shetland Islands (West Antarctica). Antarctic Science, 16(2), 191-197.
- Senglaub, Y., Brix, M. R., Adriasola, A. C., & Littke, R. (2005). New information on the thermal history of the southwestern Lower Saxony Basin, northern Germany, based on fissiontrack analysis. Geologische Rundschau = International Journal of Earth Sciences, 94(5-6), 876-896.
- Sherlock, S. C., Kelley, S. P., Parnell, J., Green, P., Lee, P., Osinski, G. R., et al. (2005). Re-evaluating the age of the Haughton impact event. Meteoritics & Planetary Science, 40(12), 1777-1787.
- Sherrod, D. R., Nishimitsu, Y. and Tagami, T. (2003). New K-Ar ages and the geologic evidence against rejuvenated-stage volcanism at Haleakala, East Maui, a postshield-stage volcano of the Hawaiian island chain. Geol. Soc. Am. Bull., 115, 683-694.
- Shi XB, Qiu XL, Liu HL (2006). Thermochronological analyses on the cooling history of the Lincang granitoid bathotith, Western Yunnan Acta Petrologica Sinica 22 (2): 465-479.
- Shi XB, Qiu XL, Liu HL, (2006). Cenozoic cooling history of Lincang granitoid batholith, western Yunnan: Evidence from fission track data Chinese Journal of Geophysics-Chinese Edition 49 (1): 135-142
- Shuster, D. L., & Farley, K. A. (2004). (super 4) He/ (super 3) He thermochronometry. Earth and Planetary Science Letters, 217(1-2), 1-17.
- Shuster, D. L., Ehlers, T. A., Rusmore, M. E., & Farley, K. A. (2005). Rapid glacial erosion at 1.8 Ma revealed by (super 4) He/ (super 3) He thermochronometry. Science, 310(5754), 1668-1670.
- Shuster, D. L., Vasconcelos, P. M., Heim, J. A., & Farley, K. A. (2005). Weathering geochronology by (U-Th)/He dating of goethite. Geochimica et Cosmochimica Acta, 69(3), 659-673.

- Siddoway, C. S., Baldwin, S. L., Fitzgerald, P. G., Fanning, C. M., & Luyendyk, B. P. (2004). Ross Sea mylonites and the timing of intracontinental extension within the West Antarctic rift system. Geology, 32(1), 57-60.
- Sinclair, H. D., Gibson, M., Naylor, M., & Morris, R. G. (2005). Asymmetric growth of the Pyrenees revealed through measurement and modelling of orogenic fluxes. American Journal of Science, 305(5), 369-406.
- Singh, J., Singh, L., & Kher, S. (2003). A comparison of fission track and laser fluorometry techniques for uranium analysis in water samples. Radiation Measurements, 36(1-6), 517-519.
- Sniderman, J. M. K., O'Sullivan, P. B., Hollis, J. D., & Kershaw, A. P. (2004). Late Pliocene vegetation and climate change at Stony Creek Basin, Western Uplands of Victoria, Australia. Proceedings of the Royal Society of Victoria, 116(1), 77-92.
- Sobel, E. R. and Strecker, M. R. (2003), Uplift, exhumation, and precipitation: Tectonic and climatic control of Late Cenozoic landscape evolution in the northern Sierras Pampeanas, Argentina: Basin Research, v. 15, doi: 10.1046/j.1365-2117.2003.00214.x.
- Sobel, E. R., Oskin, M., Burbank, D., and Mikolaichuk, A. (2006), Exhumation of basement-cored uplifts: Example of the Kyrgyz Range quantified with apatite fission-track thermochronology: Tectonics, v. 25, TC2008, doi:10.1029/2005TC001809.
- Sobel, E.R., Chen, J., and Heermance, R.V., in press, Late Oligocene - Early Miocene initiation of shortening in the Southwestern Chinese Tian Shan: Implications for Neogene shortening rate variations: Earth and Planetary Science Letters.
- Spiegel, C., Kuhlemann, J., Frisch, W. (in press). Tracing sediment pathway by zircon fission track analysis: Oligocene marine connections in Central Europe. International Journal of Earth Sciences.
- Spiegel, C., Sachsenhofer, R. F., Privalov, V. A., Zhykalyak, M. V., & Panova, E. A. (2004). Thermotectonic evolution of the Ukrainian Donbas fold belt; evidence from zircon and apatite fission track data. Tectonophysics, 383(3-4), 193-215.
- Spiegel, C., Siebel, W., Kuhlemann, J., Frisch, W. (2004). Toward a comprehensive provenance analysis: a multimethod approach and its implications for the evolution of the Central Alps. In: "Detrital thermochronology Provenance Analysis, Exhumation, and Landscape Evolution in Mountain Belts, edited by Matthias Bernet and Cornelia Spiegel. GSA Special Paper 378, 37-50.
- Spikings, R. A., Winkler, W., Hughes, R. A., & Handler, R. (2005). Thermochronology of allochthonous terranes in Ecuador; unravelling the accretionary and post-accretionary history of the Northern Andes. Tectonophysics, 399(1-4), 195-220.
- Spikings, R.A., Foster, D.A. and Kohn, B.P. (2006). Low temperature (<110°C) th/ermal history of the Mt. Isa and Murphy Imliers, northeast Australia: evidence from apatite fission track thermochronology. Australian Journal of Earth Sciences, 53: 151-165.
- Spotila, J. A. (2005). Applications of low-temperature thermochronometry to quantification of recent exhumation in mountain belts. Reviews in Mineralogy and Geochemistry, 58(1), 449-466.
- Spotila, J. A., Buscher, J. T., Meigs, A. J., & Reiners, P. W. (2004). Long-term glacial erosion of active mountain belts;

example of the Chugach-St. Elias Range, Alaska. Geology, 32(6), 501-504.

- Stefani C., Zattin M. & Grandesso P. Petrographic signature of Paleogene turbiditic sedimentation in north-eastern Italy. GSA Sp. Publ., in press.
- Stephenson, J., Gallagher, K. and Holmes, C. (2006) Low temperature thermochronology and modelling strategies for multiple samples 2 : Partition modeling for 2D and 3D distributions with discontinuities, Earth Planet Sci. Letts. 241, 557-570.
- Stewart, R. J., & Brandon, M. T. (2004). Detrital-zircon fissiontrack ages for the "Hoh Formation"; implications for late Cenozoic evolution of the Cascadia subduction wedge. Geological Society of America Bulletin, 116(1-2), 60-75.
- Stockli, D. F. (2005). Application of low-temperature thermochronometry to extensional tectonic settings. Reviews in Mineralogy and Geochemistry, 58(1), 411-448.
- Stockli, D. F., & Anonymous. (2005). Application of multiple (U-Th)/He geo- and thermochronometers with closure temperatures < approximately 250 degrees C. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 27.
- Stockli, D. F., & Farley, K. A. (2004). Empirical constraints on the titanite (U-Th)/He partial retention zone from the KTB drill hole. Chemical Geology, 207(3-4), 223-236.
- Stockli, D. F., Farley, K. A., Walker, J. D., Blackburn, T. J., & Anonymous. (2005). He diffusion and (U-Th)/He thermochronometry of monazite and rutile. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 8.
- Stuart, F. M., Persano, C., Balestrieri, M. L., Bishop, P., & Anonymous. (2005). Apatite fission track and (U-Th)/He thermochronometers constraints on the development of two high elevation passive margins. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 306.
- Sun, Z., Yang, Z., Pei, J., Ge, X., Wang, X., Yang, T., et al. (2005). Magnetostratigraphy of Paleogene sediments from northern Qaidam Basin, China; implications for tectonic uplift and block rotation in northern Tibetan Plateau. Earth and Planetary Science Letters, 237(3-4), 635-646.
- Tagami, T. (2005). Zircon fission-track thermochronology and applications to fault studies. Reviews in Mineralogy and Geochemistry, 58(1), 95-122.
- Tagami, T. and Murakami, M. (2006). Probing fault zone heterogeneity on the Nojima fault: Constraints from zircon fission-track analysis of borehole samples. Tectonophysics, in press.
- Tagami, T., & Murakami, M. (2005). Zircon fission-track thermochronology of the Nojima fault zone, Japan. AAPG Memoir, 85, 269-285.
- Tagami, T., & O'Sullivan, P. B. (2005). Fundamentals of fissiontrack thermochronology. Reviews in Mineralogy and Geochemistry, 58(1), 19-47.
- Tagami, T., Nishimitsu, Y. and Sherrod, D. R. (2003). Rejuvenated-stage volcanism after 0.6-Myr quiescence at West Maui volcano, Hawaii: new evidence from K-Ar ages and chemistry of Lahaina Volcanics. J. Volcanol. Geotherm. Res., 120, 207-214.
- Tello Saenz, C. A., Hackspacher, P. C., Hadler Neto, J. C., lunes, P. J., Guedes, S., Ribeiro, L. F. B., et al. (2003). Recognition of Cretaceous, Paleocene, and Neogene

tectonic reactivation through apatite fission-track analysis in Precambrian areas of southeast Brazil: association with the opening of the South Atlantic Ocean. Journal of South American Earth Sciences, 15(7), 765-774.

- Tello, C. A., Palissari, R., Hadler, J. C., Iunes, P. J., Guedes, S., Curvo, E. A. C., et al. (2006). Annealing experiments on induced fission tracks in apatite; measurements of horizontalconfined track lengths and track densities in basal sections and randomly oriented grains. American Mineralogist, 91(2-3), 252-260.
- Thiede, R. C., Arrowsmith, J. R., Bookhagen, B., McWilliams, M. O., Sobel, E. R., & Strecker, M. R. (2005). From tectonically to erosionally controlled development of the Himalayan Orogen. Geology, 33(8), 689-692.
- Thiede, R. C., Bookhagen, B., Arrowsmith, J. R., Sobel, E. R., & Strecker, M. R. (2004). Climatic control on rapid exhumation along the Southern Himalayan Front. Earth and Planetary Science Letters, 222(3-4), 791-806.
- Thiede, R.C., Arrowsmith, J.R., Bookhagen, B., McWilliams, M., Sobel, E.R., Strecker, M.R. (2006), Dome formation and extension in the Tethyan Himalaya, Leo Pargil, Northwest India: GSA Bulletin, v. 118, no. 5/6, p. 635-650, doi: 10.1130/B25872.1.
- Thomson, S.N., Ring U. (2006). Thermochronologic evaluation of post- collision extension in the Anatolide Orogen, western Turkey. Tectonics, 25, TC3005, doi: 10.1029/2005TC001833.
- Timar-Geng, Z., Grujic, D., & Rahn, M. (2004). Deformation at the Leventina-Simano Nappe boundary, Central Alps, Switzerland. Eclogae Geologicae Helvetiae, 97(2), 265-278.
- Tricart, P. (2004). From extension to transpression during the final exhumation of the Pelvoux and Argentera Massifs, Western Alps. Eclogae Geologicae Helvetiae, 97(3), 429-439.
- Vannay, J.C., Grasemann, B., Rahn, M., Carter, A., Baudraz, V. and Cosca, M. (2004). Miocene to Holocene exhumation of metamorphic crustal wedges in the NW Himalaya: Evidence for tectonic extrusion coupled to fluvial erosion. Tectonics, v23, pp 1014-1029
- Vega Granillo, R., & Calmus, T. (2003). Mazatan metamorphic core complex (Sonora, Mexico); structures along the detachment fault and its exhumation evolution. Journal of South American Earth Sciences, 16(4), 193-204.
- Ventura, B., & Lisker, F. (2003). Long-term landscape evolution of the northeastern margin of the Bohemian Massif; apatite fission-track data from the Erzgebirge (Germany). International Journal of Earth Sciences, 92(5), 691-700.
- Vermeesch, P., Miller, D. D., Graham, S. A., de Grave, J., & McWilliams, M. O. (2006). Multimethod detrital thermochronology of the Great Valley Group near New Idria, California. Geological Society of America Bulletin, 118(1-2), 210-218.
- Vidal, O., Wendt, A. S., & Chadderton, L. T. (2003). Experimental evidence for the pressure dependence of fission track annealing in apatite; reply [modified]. Earth and Planetary Science Letters, 215(1-2), 307-316.
- Villasante-Marcos, V., Osete, M. L., Gervilla, F., & Garcia-Duenas, V. (2003). Palaeomagnetic study of the Ronda peridotites (Betic Cordillera, southern Spain). Tectonophysics, 377(1-2), 119-141.

- Wallis, S., Moriyama, Y., & Tagami, T. (2004). Exhumation rates and age of metamorphism in the Sanbagawa Belt; new constraints from zircon fission track analysis. Journal of Metamorphic Geology, 22(1), 17-24.
- Wang, E., Xu, F., Zhou, J., Wan, J., & Burchfiel, B. C. (2006). Eastward migration of the Qaidam Basin and its implications for Cenozoic evolution of the Altyn Tagh Fault and associated river systems. Geological Society of America Bulletin, 118(3-4), 349-365.
- Wang, R.-C., Xu, S.-J., Fang, Z., Shieh, Y.-N., Li, H.-M., Li, D.-M., et al. (2004). Protolith age and exhumation history of metagranites from the Dabie UHP metamorphic belt in eastcentral China; a multi-chronological study. Geochemical Journal, 38(4), 345-362.
- Weber, K. and Raab, M.J. (2003). Tectonomorphological development of Namibia. Zbl. Geol. Paläont. Teil I: 117-129.
- Weber, U. D., Kohn, B. P., Gleadow, A. J. W., & Nelson, D. R. (2005). Low temperature Phanerozoic history of the northern Yilgarn Craton, Western Australia. Tectonophysics, 400, no.1-4, 127-151.
- Weber, U.D., Hill, K.C., Brown, R.W., Gallagher, K., Kohn, B.P., Gleadow, A.J.W. and Foster, D.A. (2004). Sediment supply to the Gippsland Basin from thermal history analysis: constraints on Emperor - Golden Beach reservoir composition. APPEA Journal 2004, 44: 397-415.
- Weber, U.D., Kohn, B.P. and Gleadow, A.J.W. (2005). Muccan Batholith, Eastern Pilbara, Western Australia. In: Anand, R.R. and de Broekert, P. (Editors), Regolith Landscape Evolution Across Australia: A compilation of regolith landscape case studies with regolith landscape evolution models, CRC LEME, Perth, pp. 338-343.
- Willett, S. D., Fisher, D., Fuller, C., Yeh, E.-C., & Lu, C.-Y. (2003). Erosion rates and orogenic-wedge kinematics in Taiwan inferred from fission-track thermochronometry. Geology, 31(11), 945-948.
- Williams, G. A., Turner, J. P., & Holford, S. P. (2005). Inversion and exhumation of the St. George's Channel basin, offshore Wales, UK. Journal of the Geological Society of London, 162(1), 97-110.
- Willner, A. P., Thomson, S. N., Kroener, A., Wartho, J.-A., Wijbrans, J. R., & Herve, F. (2005). Time markers for the evolution and exhumation history of a Late Paleozoic paired metamorphic belt in north-central Chile (34 degrees -35 degrees 30'S). Journal of Petrology, 46(9), 1835-1858.
- Willner, A.P., Hervé, F., Thomson, S.N. & Massonne, H.-J. (2004). Converging PT-paths of different HP-LT metamorphic units within a Mesozoic accretionary belt (Diego de Almagro Island, Southern Chile, 51°30`S): Evidence for juxtaposition during late shortening of an active continental margin. Mineralogy and Petrology, 81, p. 43-84.
- Yamada, K., Tagami, T. and Kamata, H. (2006). K-Ar geochronology of rhyolitic rocks in the Hohi volcanic zone, central Kyushu, Japan. J. Asian Earth Sci., in press.

- Yamada, K., Tagami, T. and Shimobayashi, N. (2003). Experimental study on hydrothermal annealing of fission tracks in zircon., Chem. Geol., 201, 351-357.
- Yang, T. F., Chen, C. H., Tien, R. L., Song, S. R., & Liu, T. K. (2003). Remnant magmatic activity in the Coastal Range of east Taiwan after arc-continent collision; fission-track data and (super 3) He/ (super 4) He ratio evidence. Radiation Measurements, 36(1-6), 343-349.
- Yuan, W., Carter, A., Dong, J., Bao, Z., An, Y. Guo (2006). Mesozoic-Tertiary exhumation history of the Altai Mountains, northern Xinjiang, China: New constraints from apatite fission track data. Tectonophysics v412, p183-193.
- Zarki-Jakni, B., van der Beek, P., Poupeau, G., Sosson, M., Labrin, E., Rossi, P., et al. (2004). Cenozoic denudation of Corsica in response to Ligurian and Tyrrhenian extension; results from apatite fission track thermochronology. Tectonics, 23, no.1, 18.
- Zattin M. & Zuffa G.G. (2004) Unravelling source rocks of episutural and foredeep Late Eocene-Miocene arenites of northern Apennines and southern Alps. Bollettino Società Geologica Italiana, 123, 67-76.
- Zattin M., Cuman A., Fantoni R., Martin S., Scotti P. & Stefani C. (2006) – From Middle Jurassic heating to Neogene cooling: the thermochronological evolution of the southern Alps. Tectonophysics, 414, 191-202.
- Zattin, M. (2003). Apatite fission-track analysis and the thermochronology of sedimentary successions. Atti Ticinensi di Scienze della Terra, 9, Serie speciale, 33-42.
- Zattin, M., Cuman, A., Fantoni, R., Martin, S., Scotti, P., & Stefani, C. (2003). Thermochronological evolution of the eastern Southern Alps from vitrinite reflectance and fissiontrack data. Atti Ticinensi di Scienze della Terra, 9, Serie speciale, 92-95.
- Zattin, M., Okay, A. I., & Cavazza, W. (2005). Fission-track evidence for late Oligocene and mid-Miocene activity along the North Anatolian Fault in south-western Thrace. Terra Nova, 17(2), 95-101.
- Zattin, M., Stefani, C., & Martin, S. (2003). Detrital fission-track analysis and sedimentary petrofacies as keys of Alpine exhumation; the example of the Venetian foreland (European Southern Alps, Italy). Journal of Sedimentary Research, 73(6), 1051-1061.
- Zhu, W., Shu, L., Sun, Y., Wang, F., Zhao, Z., & Anonymous. (2005). Mesozoic-Cenozoic exhumation history of north Tianshan, northwest China; constraints from fission track analysis. Geochimica et Cosmochimica Acta, 69(10, Suppl.), 310.
- Zwingmann, H., & Mancktelow, N. S. (2004). Timing of Alpine fault gouges. Earth and Planetary Science Letters, 223(3-4), 415-425.

# International Fission-Track Directory

This revised and extended International Fission-Track Directory is by no means complete or accurate. Although a number of people do not figure in the staff, e-mail or phone directories of the institutes under which they are mentioned, this has not been interpreted as a sign of non-existence, and they have for the moment been retained with their old affiliations. It would be much appreciated if you would let the editor know if your address has changed, or if people have joined or definitively left your lab, so that the directory can be updated.

## Abbate, Ernesto

Dipartimento di Science della Terra University of Florence Via La Pira 4, 50121 Firenze, Italy Phone: +39 55 275 7527 Fax: +39 55 218 628 E-Mail: abbate@unifi.it Web: steno.geo.unifi.it

#### Aizawa, Minoru

Department of Geology Imperial College Prince Consort Road, London SW7 2BP, UK Phone: +44 171 594 6468 Fax: +44 171 594 6464 E-Mail: m.aizawa@ic.ac.uk

## Alson, Pedro

Laboratorios de Termocronología y Geomatemáticas Facultad de Ingeniería Universidad Central de Venezuela Venezuela E-Mail: palson@cantv.net

## Andriessen, Paul, A.M.

Department of Isotope Geochemistry Faculteit Aardwetenschappen Vrije Universiteit Amsterdam De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands Phone: +31 20 4447340 Fax: +31 20 6462457 E-Mail: Paul.Andriessen@falw.vu.nl Web: www.geo.vu.nl/~isotopen

#### Armstrong, Phillip, A.

Department of Geosciences California State University P.O. Box 6850, Fullerton CA 92834-6850, USA Phone: +1 714 278 3169, +1 714 278 7266 E-Mail: parmstrong@fullerton.edu Web: geology.fullerton.edu/index.htm

## Arne, Dennis

Alpine Geological & Technical Services 126A Delany Avenue, Bright Victoria 3741, Australia E-Mail: dennis.arne@bigpond.com

#### Augusto, Pedro, F. P. M.

Departamento de Raios Cósmicos e Cronologia Instituto de Física Gleb Wataghin Universidade Estadual de Campinas Caixa Postal 6165, CEP 13083-970 UNICAMP Campinas SP, Brazil Phone: +55 19 3788 5362 Fax: +55 19 3788 5512 E-Mail: pmoreira@ifi.unicamp.br Web: www.ifi.unicamp.br/~turtelli/www0.htm

## Avé Lallemant, Hans, G.

Department of Earth Science Rice University 6100 Main St, Houston TX 77005-1892, USA Phone: +1 713 348 4880 Fax: +1 713 348 5214 E-Mail: ave@rice.edu Web: terra.rice.edu

#### Balcázar Garcia, Miguel

Departamento de Estudios del Ambiente Instituto nacional de investigaciones Nucleares Apartado Postal 18-1027, México D.F. 11801, Mexico E-Mail: mbg@nuclear.inin.mx Web: www.inin.mx

#### Baldwin, Suzanne

Heroy Geology Laboratory Department of Earth Sciences Syracuse University Syracuse, New York 13244-1070, USA Phone: +1 315 443 4920 Fax: +1 315 443 2619 E-Mail: sbaldwin@syr.edu Web: su-thermochronology.syr.edu

## Balestrieri, Maria, L.

CNR-Istituto di Geoscienze e Georisorse Sezione di Firenze Via La Pira 4, 50121 Firenze, Italy Phone: +39 55 275 7494 Fax: +39 55 218 628 E-Mail: balestrieri@jgg.cnr.it, mbalestrieri@unifi.it Web: http://www.igg.cnr.it

## Barbarand, Jocelyn

UMR Orsay Terre 8616 Université Paris-Sud Bâtiment 504, 91405 Orsay cedex, France Phone: +33 1 69 15 48 68 Fax: +33 1 69 15 48 68 E-Mail: barbaran@geol.u-psud.fr

## Barbero González, Luis, C.

Dipartimento Geologia Facultad de Ciencias del Mar y Ambientales Universidad de Cadiz 1150 Puerto Real, Cadiz, Spain Phone: +34 956 470 864 Fax: +34 956 470 811 E-Mail: luis.barbero@uca.es Web: www2.uca.es/facultad/ciencias\_mar

## Batt, Geoffrey

Department of Geology Royal Holloway University of London Egham Hill, Egham Surrey TW20 OEX, UK Phone: +44 1784 443606 E-Mail: g.batt@gl.rhul.ac.uk

## Belton, Dave, X.

CSIRO Exploration and Mining MARC - School of Physics The University of Melbourne VIC 3010, Australia Phone: +61 3 8344 5081 Fax: +61 3 8344 5081 E-Mail: David.Belton@csiro.au Web: http://www.nmp.csiro.au

## Bergman, Steven, C.

Departments of Geosciences & Science & Mathematics University of Texas at Dallas Box 830688, Richardson TX 75083-0688, USA Phone: +1 972 883 4365 Fax: +1 972 883 2537 E-Mail: bergman@utdalla.edu Web: www.utdallas.edu/dept/geosciences/Bergman.html

#### Bermudez, Mauricio

Laboratorios de Termocronología y Geomatemáticas Facultad de Ingenieria Universidad Central de Venezuela Caracas, Venezuela E-Mail: mbermude@euler.ciens.ucv.ve

## Bernet, Matthias

Department of Geosciences State University of New York New Paltz, New York, 12561, USA Phone: +1 845 257 3765 E-Mail: bernetm@newpaltz.edu

#### Bigazzi, Giulio

Istituto di Geocronologia e Geochimica Isotopica Consiglio Nazionale delle Ricerche Area delle Ricerca di Pisa Via G. Moruzzi 1, 56124 Pisa, Italy Phone: +39 50 315 2283 Fax: +39 50 315 2360 E-Mail: g.bigazzi@igg.cnr.it Web: www.area.pi.cnr.it/AREAEN/iggi.html

#### Blythe, Ann, E.

Department of Earth Sciences University of Southern California Los Angeles CA 90089-0740, USA Phone: +1 213 8211094 Fax: +1 213 740 8801 E-Mail: blythe@earth.usc.edu

#### Boettcher, Stefan, S.

Department of Geological Sciences University of Texas at Austin Austin TX 78712, USA Phone: +1 512 471 8547 Fax: +1 512 471 9425 E-Mail: sboett@maestro.geo.utexas.edu Web: www.geo.utexas.edu

#### Bojar, Ana-Voica

Insitut für Geologie und Paläontologie Karl-Franzens Universität Heinrichstraße 26, A-8010 Graz, Austria Phone: +43 316 380 8727 Fax: +43 316 380 9871 E-Mail: ana-voica.bojar@kfunigraz.ac.at/ Web: www-ang.kfunigraz.ac.at/~bojar

## Bondar, Julia

State Science Center of Environmental Radiogeochemistry 34 Palladin Avenue, 03142 Kiev, Ukraine Fax: +380 44 451 02 36 E-Mail: center@radgeo.freenet.kiev.ua

#### Bondarenko, Oleg

Deputy Director - Radiation Safety Chief SSSIE Echocentre 6 Shkolnaya Str. Chornobyl, Kievskaya obl., Ukraine Phone: +380 4493 52218 Fax: +380 4493 52398 E-Mail: boa@ic-chernobyl.kiev.ua Web: http://www.ic-chernobyl.kiev.ua/ecocentr.htm

#### Boulyga, Sergei

Central Department for Analytical Chemistry Research Centre Jülich D-52425 Jülich, Germany Phone: +49 2461 613 291 Fax: +49 2461 612 560 E-Mail: s.boulyga@fz-juelich.de

## Boztug, Durmus

Jeoloji Mühendisli\_i Bölümü Mühendislik Fakültesi Cumhuriyet Üniversitesi 58140 Sivas, Turkey Phone: +90 346 219 1010 Fax: +90 346 219 1171 E-Mail: boztug@cumhuriyet.edu.tr Web: jeoloji.cumhuriyet.edu.tr

## Brandon, Mark, T.

Kline Geology Laboratory Department of Geology and Geophysics Yale University P.O. Box 208109 210 Whitney Avenue, New Haven CT 06520-8109, USA Phone: +1 203 432 3135 Fax: +1 203 432 3134 E-Mail: mark.brandon@yale.edu Web: www.geology.yale.edu/~brandon/

#### Brix, Manfred, R.

Institut für Geologie Mineralogie und Geophysik Fakultät für Geowissenschaften Ruhr-Universität Bochum Postfach 102148, D-44721 Bochum, Germany Phone: +49 2343 223 236 Fax: +49 2343 214 572 E-Mail: manfred.r.brix@ruhr-uni-bochum.de Web: www.ruhr-uni-bochum.de/hardrock/Mitarbeiter/brix.html

## Brown, Roderick, W.

Department of Geographical and Earth Sciences Gregory Building University of Glasgow Glasgow G12 8QQ, Scotland Phone: +44 141 330 5460 Fax: +44 141 330 4817 E-Mail: r.brown@ges.gla.ac.uk Web: http://www.ges.gla.ac.uk

## **Brunel, Maurice**

ISTEEM GGP laboratory CC 058 Universite Montpellier II 1 Place E. Bataillon, 34095 Montpellier Cedex, France Phone: +33 4 67 14 36 47 Fax: +33 4 67 14 42 53 E-Mail: brunel@dstu.univ-montp2.fr Web: www.dstu.univ-montp2.fr

## Burchart, Jan

Institute of Geological Sciences Polish Academy of Sciences Zwirki i Wigury 93, 02-089 Warsaw, Poland Web: www.pan.pl/english/index.html Calmus, Thierry Instituto de Geología Universidad Nacional Autonoma de México (ERNO) Apartado Postal 1039, Hermosillo Sonora 8300, Mexico Phone: +52 662 217 5019 E-Mail: tcalmus@servidor.unam.mx Web: geologia.igeolcu.unam.mx/geol.htm

## Carlson, William, D.

Department of Geological Sciences University of Texas at Austin Austin TX 78712, USA Phone: +1 512 471 4770 Fax: +1 512 471 9425 E-Mail: wcarlson@mail.utexas.edu Web: www.geo.utexas.edu/faculty/carlson.htm

## Carpéna, Joëlle

Commissariat Energie Atomique Direction Energie Nucléaire Centre d'Etudes de Cadarache DED/SEP/Bâtiment 307, 13108 Saint Paul lez Durance CDX, France Phone: +33 4 42 25 46 94 E-Mail: carpe@bahia.cad.cea.fr Web: www-cad.cea.fr/r10.htm

## Carpenter, Stephen, B.

NIST Gaithersburg, Maryland, USA

#### Carrapa, Barbara

Institut für Geowissenschaften Universität Potsdam Postfach 60 15 53, D-14415 Potsdam, Germany Phone: +49 331 977 5840 Fax: +49 331 977 5700 E-Mail: carrapa@geo.uni-potsdam.de

## Carrière, Kevin

FS Archäometrie Heidelberger Akad. d. Wissenschaften am MPI für Kernphysik. Saupfercheckweg 1, D-69117 Heidelberg, Deutchland Phone: +49 6221 516 337 Fax: +49 6221 516 633 E-Mail: carriere@mpi-hd.mpg.de

#### Carter, Andrew

London Thermochronology Research Group Research School of Earth Sciences University & Birkbeck College London London WC1E 6B, UK Phone: +44 20 7679 2418 Fax: +44 20 7813 2802 E-Mail: a.carter@ucl.ac.uk Web: www.fissiontrack.ucl.ac.uk/Personal/Andypage/AC.html

## Cawley, Steve

BP-Amoco Farburn Industrial Estate Aberdeen AB21 7PB, UK E-Mail: cawleysj@bp.com

## Cederbom, Charlotte

Chadderton, Lewis, T. Research School of Physical Sciences and Engineering Australian National University Canberra ACT, Australia Phone: +61 2 6125 4542 Fax: +61 2 6125 0043 E-Mail: Lewis.Chadderton@anu.edu.au Web: www.rsphysse.anu.edu.au/ampl/research/rad/index.html

### Clavero, Jorge

Servicio Nacional de Geologia y Mineria Av. Sta. Maria 0104, Casilla 10465 Providencia, Santiago de Chile Phone: +56 2 737 5050 Fax: +56 2 777 1906 E-Mail: JClavero@sernageomin.cl Web: www.sernageomin.cl/prototipo/index.htm

#### Cloos, Mark

Department of Geological Sciences University of Texas at Austin Austin TX 78712, USA Phone: +1 512 471 4170 Fax: +1 512 471 9425 E-Mail: cloos@mail.utexas.edu Web: www.geo.utexas.edu/faculty/cloos.htm

## Coleman, Max

Research Institute for Sedimentology The University of Reading P.O. Box 227, Whiteknights Reading RG6 6AB, UK Phone: +44 118 931 6627 Fax: +44 118 931 0279 E-Mail: m.l.coleman@reading.ac.uk Web: www.rdg.ac.uk/AcaDepts/sl/Sed/Staff/People/mlc1.html

#### Corrigan, Jeff, D.

ARCO Exploration and Production 2300 W. Plano Parkway, Plano TX 75075, USA Phone: +1 214 509 4090 E-Mail: dprjdc@arco.com

#### Coutand, Isabelle

Université des Sciences et Technologies de Lille 1 UMR-CNRS PBDS UFR des Sciences de la Terre (bât.SN5) 59655 Villeneuve d'Ascq cedex, France Phone: +33 3 20 43 41 21 Fax: +33 3 20 43 49 10 E-Mail: Isabelle.Coutand@univ-lille1.fr

### Danhara, Tohru

Kyoto Fission-Track Co. Ltd. 44-4 Minamitajiri-cho Omiya Kita-ku, Kyoto 603-8832, Japan Phone: +81 75 493 0684 Fax: +81 75 493 0741 E-Mail: kyoto-ft@mb.neweb.ne.jp

### Danišík, Martin

Institut für Geologie und Paläontologie Eberhard Karls Universität Tübingen Sigwartstrasse 10, D-72076 Tübingen, Germany Phone: +49 7071 2972494 Fax: +49 7071 5059 E-Mail: martin.danisik@uni-tuebingen.de Web: www.uni-tuebingen.de/geo/gpi

## De Corte, Frans

Institute for Nuclear Sciences University of Gent Proeftuinstraat 86, B-9000 Gent, Belgium Phone: +32 9 264 6627 Fax: +32 9 264 6699 E-Mail: Frans.Decorte@rug.ac.be Web: allserv.rug.ac.be/~jddonder/AnalChem/incmain.htm#Home

#### Deeken, Anke

Institut fuer Geologische Wissenschaften Fachrichtung Geologie Freie Universitaet Berlin Maltheser Str. 74-100, D-12249 Berlin, Germany Phone: +49 30 83870345 E-Mail: adeeken@zedat.fu-berlin.de De Grave, Johan Geologisch Instituut Universiteit Gent Krijgslaan 281 (S8), B-9000 Gent, Belgium Phone: +32 9 264 4592 Fax: +32 9 264 4597 E-Mail: Johan.Degrave@rug.ac.be Web: allserv.rug.ac.be/-fmees/mineralogy.htm

#### De Wispelaere, Antoine

Institute for Nuclear Sciences University of Gent Proeftuinstraat 86, B-9000 Gent, Belgium Phone: +32 9 264 6627 Fax: +32 9 264 6699 E-Mail: Antoine.Dewispelaere@rug.ac.be Web: allserv.rug.ac.be/~jddonder/AnalChem/incmain.htm#Home

## De Wit, M., C.J.

De Beers Consolidated Mines Ltd. P.O. Box 7383, Centurion 0046 Pretoria, South Africa

## Dickin, Alan, P.

School of Geography and Geology McMaster University Hamilton Ontario L8S 4M1, USA Phone: +1 905 525 9140 Ext. 24365 Fax: +1 905 522 3141 E-Mail: dickin@mcmaster.ca Web: www.science.mcmaster.ca/geology/dickin.html

#### Dodson, Martin, H.

Department of Earth Sciences University of Leeds Leeds LS2 9JT, UK

## Dokka, Roy, K.

Department of Geology Louisiana State University E235 Howe Russell Building LSU, Baton Rouge Louisiana 70803-4101, USA Phone: +1 225 388 3353 Fax: +1 225 388 2302 E-Mail: dokka@geol.lsu.edu Web: www.geol.lsu.edu/department/facilities/laboratories/tectonics/in

#### Dolati, Asghar

dex.html

Institute of Geology Department of Earth Sciences ETH-Zentrum Leonhardstrasse 19 /LEB, CH-8092 Zürich, Switzerland Phone: +41 44 632 8823 Fax: +41 44 632 1030 E-Mail: Asghar.dolati@erdw.ethz.ch Web: www.erdw.ethz.ch

## Donelick, Raymond, A.

Apatite to Zircon Ltd 1521 Pine Cone Road Moscow, ID 83843, USA Phone: +1 208 892 1493 Fax: +1 208 875 8881 E-Mail: donelick@apatite.com

## Donelick, Margaret, B.

Apatite to Zircon Ltd 1521 Pine Cone Road Moscow, ID 83843, USA Phone: +1 208 892 1493 Fax: +1 208 875 8881 E-Mail: donelick@apatite.com Web: www.apatite.com

### Dorighel, Olivier

Groupe de Géophysique Nucléaire Institut Dolomieu Université Joseph Fourier 15 Rue Maurice Gignoux, 38031 Grenoble Cedex, France

## Duddy, Ian, R.

Geotrack International Pty. Ltd. 37 Melville Road, West Brunswick VIC 3055, Australia Phone: +61 3 9380 1077 Fax: +61 3 9380 1477 E-Mail: mail@geotrack.com.au Web: www.geotrack.com.au

## Dumitru, Trevor, A.

Department of Geology and Environmental Sciences Stanford University Stanford CA 94305-2115, USA Phone: +1 650 725 6155 Fax: +1 650 725 6155 E-Mail: trevor@pangea.stanford.edu Web: pangea.Stanford.edu/structure/projects/fission\_track.html

## Dunai, Tibor, J.

School of GeoSciences The University of Edinburgh Geography Building, Drummond Street, Edinburgh EH8 9XP, UK Phone: +44 131 650 2546 Fax: +44 131 650 2524 E-Mail: Tibor.Dunai@ed.ac.uk

## Dunkl, István

Geoscience Center Georg-August-Universität Göttingen, Goldschmidtstrasse 3, D-37077 Göttingen, Germany E-Mail: istvan.dunkl@geo.uni-goettingen.de

## Durrani, Saeed, A.

School of Physics and Space Research University of Birmingham Birmingham B15 2TT, UK Phone: +44 21 414 4691, +44 21 414 4655 Fax: +44 21 414 4693 E-Mail: s.a.durrani@bham.ac.uk

#### Eby, Nelson, G.

Department of Environmental Earth & Atmospheric Sciences University of Massachusetts Lowell MA 01854, USA Phone: +1 978 934 3907 Fax: +1 978 934 3069 E-Mail: Nelson\_Eby@uml.edu

#### Edwards, Michael

Institut für Geologie University of Vienna Althanstrasse 14, Vienna A-1090, Austria Phone: +49 177 783 2827 E-Mail: medwards@gmx.net

#### Ege, Harald

Institut für Geologische Wissenschaften Freie Universität Berlin Malteserstrasse 74-100, D-12249 Berlin, Germany E-Mail: ege@zedat.fu-berlin.de

## Ehlers, Todd

Department of Geological Sciences University of Michigan 2534 C.C. Little Building 425 East University, Ann Arbor Michigan 48109-1063, USA Phone: +1 734 763 5112 (temporary) Fax: +1 734 763 4690 E-Mail: tehlers@umich.edu

## Enkelmann, Eva

Earth and Environmental Sciences Department Lehigh University 31 Williams Dr., Bethlehem PA 18015, USA Phone: +1 610 758 3688 Fax: +1 610 758 3677 E-Mail: eva.enkelmann@lehigh.edu

## Evarts, Russell, C.

U.S. Geological Survey MS-999 345 Middlefield Road, Menlo Park CA 94025, USA Phone: +1 650 329 4929 Fax: +1 650 329 4936 E-Mail: revarts@usgs.gov Web: www.usgs.gov

## Fabel, Derek

Department of Geographical and Earth Sciences University of Glasgow University Avenue, Glasgow G12 8QQ, UK Phone: +44 141 330 2262 Fax: +44 141 330 4894 E-Mail: dfabel@ges.gla.ac.uk Web: http://web.ges.gla.ac.uk/~dfabel

## Farley, Kenneth, A.

Division of Geological and Planetary Sciences California Institute of Technology Caltech MS 170-25, Pasadena CA 91125, USA Phone: +1 626 395 6005 Fax: +1 626 568 0935 E-Mail: farley@gps.caltech.edu Web: www.gps.caltech.edu/~farley/farley.htm

## Fayon, Annia, K.

Department of Geology and Geophysics University of Minnesota-Twin Cities 108 Pillsbury Hall 310 Pillsbury Drive SE, Minneapolis MN 55455, USA Phone: +1 612 626 9805 Fax: +1 612 625 3819 E-Mail: annia@geolab.geo.umn.edu Web: www.geo.umn.edu/orgs/whitney/DLW\_postdoc.html

### Feinstein, Shimon

Department of Geological and Environmental Sciences Ben Gurion University of the Negev P.O. Box 653, Beer Sheva 84105, Israel Phone: +972 8 647 2622 Fax: +972 8 647 2997 E-Mail: shimon@bgumail.bgu.ac.il Web: www.bgu.ac.il/geol/index.html

## Filip, Jiri

Institute of Geology Academy of Sciences of the Czech Republic Rozvojova 135, Prague 6 CZ 165 02, Czech Republic Phone: +420 2 3308 7248 Fax: +420 2 3308 7111 E-Mail: filip@gli.cas.cz Web: www.gli.cas.cz

### Fisher, David, E.

Department of Geological Sciences University of Miami Miami FL 33124-0401, USA Phone: +1 305 284 3254 Fax: +1 305 284 4258 Web: www.as.miami.edu/geology Fitzgerald, Paul, G. Heroy Geology Laboratory Department of Earth Sciences Syracuse University Syracuse New York 13244-1070, USA Phone: +1 315 443 2871 Fax: +1 315 443 2619 E-Mail: pgfitzge@syr.edu Web: su-thermochronology.syr.edu

## Fleischer, Robert, L.

Research Professor of Geology Olin 312 Union College Schenectady, NY 12308, USA Phone: +1 518 388-6770 Fax: +1 518 388 6417 E-Mail: fleischr@union.edu

## Fletcher, John, M.

Departamento de Geología Centro de Investigación Científica y de Educación Superior de Ensenada P.O. Box 430222, Ensenada B.C., México Phone: +1 174 50 50 26031 Fax: +1 174 50 50 26021 E-Mail: jfletche@cicese.mx Web: geologia.cicese.mx/jfletche/jfletche.htm

### Foland, Sara, S.

Amoco Production Company P.O. Box 800, Denver CO 80201, USA E-Mail: sfoland@geosociety.org

## Foster, David, A.

Department of Geology University of Florida P.O. Box 112120, Gainsville FL 32611-2120, USA Phone: +1 352 392 2231 E-Mail: dafoster@ufl.edu Web: www.clas.ufl.edu/users/dfoster

## Franks, Daniel

University of Queensland St Lucia Brisbane QLD 4072, Australia E-Mail: danielfranks@hotmail.com

### Frei, Stefan

School of Earth Sciences University of Melbourne Victoria 3010, Australia Phone: +61 3 8344 4996 Fax: +61 3 8344 7761 E-Mail: s.frei@pgrad.unimelb.edu.au

### Fügenschuh, Bernhard

Geologisch Paläontologisches Institut Universität Basel Bernoullistrasse 32, Basel BS CH-4056, Switzerland Phone: +41 61 267 86 10 Fax: +41 61 267 36 13 E-Mail: bernhard.fuegenschuh@unibas.ch Web: www.unibas.ch/earth/tecto/people/fuegenschuh.html

## Galbraith, Rex, F.

Department of Statistical Science University College London Gower Street, London WC1E 6BT, UK Phone: +44 20 7679 1859 Fax: +44 20 7383 4703 E-Mail: rex@stats.ucl.ac.uk Web: www.ucl.ac.uk/geolsci/research/fissiontrack/Personal/Rexpage /Rex.html

## Gallagher, Kerry

Department of Earth Sciences and Engineering Imperial College of Science Technology and Medicine Exhibition Road, London SW7 2BT, UK Phone: +44 20 759 46424 Fax: +44 20 759 46444 E-Mail: kerry@ic.ac.uk, k.l.gallagher@ic.ac.uk Web: www.huxley.ic.ac.uk/research/Comp&Geophys/Geophys/maint ext.php3

#### Garver, John, I.

Department of Geology Union College Schenectady, NY 12308-2311, USA Phone: +1 518 388 6517 Fax: +1 518 388 6789 E-Mail: garverj@union.edu Web: zircon.union.edu/FT/FThome.html

#### George, Pete

Department of Geological Sciences University of Texas at Austin Austin TX 78712, USA Web: www.geo.utexas.edu

## Giegengack, Robert

Department of Earth and Environmental Science University of Pennsylvania Philadelphia PA 19104-6316, USA Phone: +1 215 898 5191 Fax: +1 215 573 9145 E-Mail: gieg@sas.upenn.edu Web: www.sas.upenn.edu/earth/giegenga.html

## Giles, Melvin, R.

EPT-HM Volmerlaan 8, Rijswijk, The Netherlands E-Mail: gilesm@siep.shell.com

### Glasmacher, Ulrich, A.

Geologisch-Palaeontologisches Institut Ruprecht-Karls Universität Heidelberg Im Neuenheimer Feld 234, D-69120 Heidelberg, Germany Phone: +49 6221 544866 Fax: +49 6221 545503 E-Mail: ulrich.a.glasmacher@urz.uni-heidelberg.de

## Gleadow, Andrew, J. W.

School of Earth Sciences University of Melbourne Parkville Victoria 3010, Australia Phone: +61 3 8344 9866 Fax: +61 3 8344 7761 E-Mail: gleadow@unimelb.edu.au Web: web.earthsci.unimelb.edu.au/Gleadow/Gleadow.html

## Glotzbach, Christoph

Institut für Geowissenschaften Eberhard Karls Universität Tübingen Sigwartstrasse 10, D-72076 Tübingen, Germany Phone: +49 7071 2975240 Fax: +49 7071 5059 E-Mail: christoph.glotzbach@uni-tuebingen.de

## Green, Paul, F.

Geotrack International Pty. Ltd. 37 Melville Road, West Brunswick VIC 3055, Australia Phone: +61 3 9380 1077 Fax: +61 3 9380 1477 E-Mail: mail@geotrack.com.au Web: www.geotrack.com.au Grimmer, Jens, C. Institut für Geologie Technische Universität Bergakademie Freiberg Bernhard-von-Cottastrasse 2, D-09596 Freiberg, Germany Phone: +49 3731 394599 Fax: +49 3731 393597 E-Mail: jens.grimmer@geo.tu-freiberg.de Web: www.geo.tu-freiberg.de/tektono

#### Grivet, Manuel

Laboratoire de Microanalyses Nucléaires U.F.R. des Sciences et des Techniques Université de Franche-Comté 16 route de Gray, F-25030 Besancon Cedex, France Phone: +33 3 81 66 65 01 Fax: +33 3 81 66 65 22 E-Mail: manuel.grivet@univ-fcomte.fr Web: www.univ-fcomte.fr

## Guedes de Oliveira, Sandro

Departamento de Raios Cósmicos e Cronologia Instituto de Fisica Gleb Wataghin Universidade Estadual de Campinas Caixa Postal 6165, CEP 13083-970 UNICAMP Campinas SP, Brazil Phone: +55 19 788 5522 Fax: +55 19 788 5512 E-Mail: guedes@ifi.unicamp.br Web: www.ifi.unicamp.br/~turtelli/www0.htm

## Guglielmetti, Alessandra

Instituto di Fisica Generale Applicata Università degli Studi di Milano Via Celoria 16, 20133 Milano, Italy Phone: +39 2 266 6259 Fax: +39 2 266 5717 E-Mail: guglielmetti@mi.infn.it Web: albinoni.brera.unimi.it/istituto/istituto.htm

## Gunnell, Yanni

Laboratoire de géographie physique Universite Denis Diderot Paris VII 1 Place Aristide Briand, 92195 Meudon Cedex, France Phone: +33 1 4507 5552 E-Mail: gunnell@paris7.jussieu.fr Web: www.sigu7.jussieu.fr

### Hadler Neto, Julio, C.

Departamento de Raios Cósmicos e Cronologia Instituto de Fisica Gleb Wataghin Universidade Estadual de Campinas Caixa Postal 6165, CEP 13083-970 UNICAMP Campinas SP, Brazil Phone: +55 19 788 5522 Fax: +55 19 788 5512 E-Mail: hadler@ifi.unicamp.br Web: www.ifi.unicamp.br/~turtelli/www0.htm

#### Hansen, Kirsten

#### Harrison, Mark, T.

Department of Earth and Space Sciences University of California Los Angeles 595 Charles Young Drive East, Los Angeles CA 90095-1567, USA Phone: +1 310 825 7970 Fax: +1 310 825 2779 E-Mail: tmh@argon.ess.ucla.edu Web: www.ess.ucla.edu/faculty/harrison

#### Hasebe, Noriko

Institute of Nature and Environmental Technology Kanazawa University Kanazawa 920-1192, Japan Phone: +81 76 264 5727 Fax: +81 76 264 5746 E-Mail: hasebe@kenroku.kanazawa-u.ac.jp

### Hashemi-Nezhad, Reza, S.

Department of High Energy Physics School of Physics University of Sydney A28 Sydney NSW 2006, Australia Phone: +61 2 9351 5964 Fax: +61 2 9351 7727 E-Mail: reza@physics.usyd.edu.au Web: www.physics.usyd.edu.au/hienergy/index.html

#### Hejl, Ewald, R.

Institut für Geologie und Paläontologie Universitat Salzburg Hellbrunnerstraße 34/III, A-5020 Salzburg, Austria Phone: +43 662 8044 5437 Fax: +43 662 8044 621 E-Mail: ewald.hejl@sbg.ac.at Web: www.sbg.ac.at/gew

## Hendriks, Bart, W.H.

Norges Geologiske Undersøkelse Leiv Eirikssons vei 39, 7491 Trondheim, Norway Phone: +47 73904440, Mob.:+47 45261691 Fax: +47 73921620 E-Mail: bart.hendriks@ngu.no

## Hill, Kevin, C.

School of Earth Sciences University of Melbourne Parkville Victoria 3010, Australia Phone: +61 3 8344 9708, +61 3 8344 7675 Fax: +61 3 8344 7761 E-Mail: kevin.hill@unimelb.edu.au Web: www.earthsci.unimelb.edu.au

#### Himeno, Osamu

Department of Mining Engineering Kyushu Sangyo University Hakozaki Fukuoka 812-81, Japan Phone: +81 92 642 3635 E-Mail: himeno@mine.kyushu-u.ac.jp

#### Hobro, James, W. D.

Bullard Laboratories Department of Earth Sciences University of Cambridge Madingley Road, Cambridge CB3 0EZ, UK Phone: +44 1223 337 191 Fax: +44 1223 360 779 E-Mail: hobro@esc.cam.ac.uk Web: bullard.esc.cam.ac.uk/~hobro

## Honda, Teruyyki

Atomic Energy Research Laboratory Musashi Institute of Technology Kawasaki 215, Japan Web: atomsun2.atom.musashi-tech.ac.jp

## Hoshino, Hidehiro

Kanazawa University Kakuma town Kanazawa City Ishikawa Pref., Japan E-Mail: hide43@nihonkai.kanazawa-u.ac.jp

#### House, Martha

Division of Geological and Planetary Sciences California Institute of Technology Caltech MS 100-23, Pasadena CA 91125, USA Phone: +1 626 395 6023 Fax: +1 626 568 0935 E-Mail: mhouse@gps.caltech.edu Web: www.gps.caltech.edu/%7Emhouse

## Huigen, Yvette

Department of Isotope Geochemistry Faculty of Earth sciences Vrije Universiteit Amsterdam De Boelelaan 1085, Amsterdam 1081 HV, The Netherlands Phone: +31 20 4447397 Fax: +31 20 6462457 E-Mail: huiy@geo.vu.nl Web: www.geo.vu.nl

#### Hungerbühler, Dominik

Department Erdwissenschaften ETH-Zentrum CH-8092 Zurich, Switzerland Web: www.erdw.ethz.ch

### Hurford, Anthony, J.

London Thermochronology Research Group Department of Earth Sciences University College London London WC1E 6B, UK Phone: +44 20 7679 7704 Fax: +44 20 7813 2802 E-Mail: t.hurford@ucl.ac.uk Web: www.ucl.ac.uk/geolsci/research/fissiontrack/Indexpage/index.html

## Issler, Dale, R.

Natural Resources Canada Geological Survey of Canada 3303-33rd St., NW Calgary Alberta T2L 2A7, Canada Phone: +1 403 292 7172 Fax: +1 403 292 5377 E-Mail: dissler@nrcan.gc.ca Web: www.nrcan.gc.ca/gsc/calgary/index\_e.html

#### Ito, Hisatoshi

Central Research Institute of Electric Power Industry 1646 Abiko, Abiko city Chiba 270-1194, Japan Phone: +81 70 6568 9621 Fax: +81 471 83 3182 E-Mail: ito\_hisa@criepi.denken.or.jp Web: http://criepi.denken.or.jp/en/civil/

## lunes, P., J.

Departamento de Raios Cósmicos e Cronologia Instituto de Fisica Gleb Wataghin Universidade Estadual de Campinas Caixa Postal 6165, CEP 13083-970 UNICAMP Campinas SP, Brazil Phone: +55 19 788 5522 Fax: +55 19 788 5512 E-Mail: iunes@fif.unicamp.br Web: www.ifi.unicamp.br/~turtelli/www0.htm

## Iwano, Hidekilwano, Hideki

Kyoto Fission-Track Co. Ltd. 44-4 Minamitajiri-cho Omiya Kita-ku, Kyoto 603-8832, Japan Phone: +81 75 493 0684 Fax: +81 75 493 0741 E-Mail: kyoto-ft@mb.neweb.ne.jp Iwano, Hidekic Kyoto Fission Track Co. Ltd. 44--4 Minamitajiri-cho Omiya Kita-ku, Kyoto 603-8832, Japan Phone: +81 75 493 0684 Fax: +81 75 493 0741 E-Mail: kyoto-ft@mb.neweb.ne.jp

#### Jacobs, Joachim

Department of Earth Science University of Bergen Allegaten 41, N-5007 Bergen, Norway Phone: +47 55 58 34 32 Fax: +47 55 58 36 60 E-Mail: joachim.jacobs@geo.uib.no

## Jaramillo, Jose, M.

National University Bogota, Colombia E-Mail: jose-maria.jaramillo@tepmanet.com

#### Jolivet, Marc

Laboratoire Dynamique de la Lithosphère UMR 5573 CNRS UMII Cc060 Université Montpellier II 5 Pl. E. Bataillon 34095 Montpellier cédex 05, France Phone: +33 4 67 14 37 28 Fax: +33 4 67 52 39 08 E-Mail: jolivet@dstu.univ-montp2.fr

## Jonckheere, Raymond

Geologisches Institut Technische Universität - Bergakademie Freiberg Bernhard-von-Cotta-Straße 2, D-09599 Freiberg (Sachsen), Deutschland Phone: +49 3731 393160 Fax: +49 3731 393597 E-Mail: raymond.jonckheere@geo.tu-freiberg.de

## Juez-Larre, Joaquim

Department of Isotope Geochemistry Faculty of Earth sciences Vrije Universiteit Amsterdam De Boelelaan 1085, Amsterdam 1081 HV, The Netherlands Phone: +31 20 444 7395 Fax: +31 20 646 2457 E-Mail: juej@geo.vu.nl Web: www.geo.vu.nl/~isotopen

## Kamp, Peter, J.J.

Department of Earth Sciences University of Waikato Private Bag 3105, Hamilton 2001, New Zealand Phone: +64 7 838 4876 Fax: +64 7 856 0115 E-Mail: p.kamp@waikato.ac.nz Web: sci.waikato.ac.nz/cgi-bin/dynapage.pl?pagename=erth

## Ketcham, Richard, A.

Department of Geological Sciences University of Texas at Austin Austin TX 78712, USA Phone: +1 512 471 6942 Fax: +1 512 471 9425 E-Mail: ketcham@mail.utexas.edu Web: www.geo.utexas.edu/scientist/ketcham.htm

## Kelley, Shari, A.

Department of Earth and Environmental Sciences New Mexico Institute of Mining and Technology 801 Leroy Place, Socorro NM 8780, USA Phone: +1 505 661 6171 E-Mail: sakelley@ix.netcom.com Web: www.ees.nmt.edu/faculty/kelley.html

#### Khattak, N., U.

Radiation Physics Division PINSTECH P.O. Nilore, Islamabad, Pakistan Phone: +92 051 2207725 E-Mail: niamat@pinstech.org.pk

#### Kohlmann, Fabian

School of Earth Sciences University of Melbourne Victoria 3010, Australia Phone: +61 3 8344 4996 Fax: +61 3 8344 7761 E-Mail: f.kohlmann@pgrad.unimelb.edu.au

## Kohn, Barry, P.

School of Earth Sciences University of Melbourne Parkville Victoria 3010, Australia Phone: +61 3 8344 7217, +61 3 8344 7675 Fax: +61 3 8344 7761 E-Mail: b.kohn@unimelb.edu.au

### Kowallis, Bart, J.

Brigham Young University Provo, UT 84602, USA Phone: +1 801 422 2467 E-Mail: bart\_kowallis@byu.edu

## Krochmal, Michael

Autoscan Systems Pty. Ltd. P.O. Box 112, Ormond VIC 3204, Australia Phone: +61 39 596 8065, +61 39 596 8092 Fax: +61 39 596 8369 E-Mail: autoscan@autoscan.com.au Web: members.ozemail.com.au/~autoscan/index.html

#### Krugh, Chris

Department of Earth Sciences Institute of Geology ETH-Zentrum Sonneggstrasse 5, 8092 Zürich, Switzerland E-Mail: krugh@erdw.ethz.ch

#### Kumar, Devender

Scientist Letter Box no. 11 National Geophysical Research Institute (NGRI) Uppal Road, Hyderabad - 500 007, India Phone: +91 40 27170141/27174433 extn: 2346/2347 Fax: +91 40 27171564/27170491 E-Mail: devender@ngri.res.in, devngri@rediffmail.com

## Laken, Garey

Autoscan Systems Pty. Ltd. P.O. Box 112, Ormond VIC 3204, Australia Phone: +61 39 596 8065, +61 39 596 8092 Fax: +61 39 596 8369 E-Mail: autoscan@autoscan.com.au Web: members.ozemail.com.au/~autoscan/index.html

#### Lal, Nand

Department of Earth Sciences Kurukshetra University Kurukshetra 136119, India Phone: +91 1 744 20410 Fax: +91 1 744 20277

#### Lange, Jan-Michael

Museum für Mineralogie und Geologie Staatliche Naturhistorische Sammlungen Dresden Königsbrücker Landstraße 159, D-01109 Dresden, Germany Phone: +49 351 892 6403 Fax: +49 351 892 6404 E-Mail: lange@snsd.de Web: www.snsd.de

#### Larsen, lan

Autoscan Systems Pty. Ltd. P.O. Box 112, Ormond VIC 3204, Australia Phone: +61 39 596 8065, +61 39 596 8092 Fax: +61 39 596 8369 E-Mail: autoscan@autoscan.com.au Web: members.ozemail.com.au/~autoscan/index.html

#### Laslett, Geoff

CSIRO Mathematical and Information Sciences Private Bag 10, Clayton South MDC Vic 3169, Australia Phone: +61 3 9545 8018 Fax: +61 3 9545 8080 E-Mail: geoff.laslett@cmis.csiro.au Web: www.cmis.csiro.au/Geoff.Laslett/index.htm

## Lee, Yong, I.

School of Earth and Environmental Sciences Seoul National University Korea E-Mail: lee2602@plaza.snu.ac.kr Web: sees.snu.ac.kr/bken/index.html

## Lewis, Cherry, L.E.

History of Geology Group The Geological Society of London Wells Cottage 21 Fowler Street, Macclesfield Cheshire SK10 2AN, UK Phone: +44 1625 260049 Fax: +44 1625 523690 E-Mail: clelewis@aol.com Web: www.geolsoc.org.uk

### Lim, Hyoun, S.

Lab. of Sedimentary System School of Earth and Environmental Sciences Seoul National University NS80, Seoul 151-747, Korea Phone: +82 2 878 0252 E-Mail: tracker@freechal.com, tracker@nate.com

#### Link, Katharina

Institut für Mineralogie Petrologie und Geochemie Albert-Ludwigs-Universität Albertstrasse 23b, 79104 Freiburg, Germany Phone: +49 761 203 6425 Fax: +49 761 203 6396 E-Mail: kalink@uni-freiburg.de Web: www.uni-freiburg.de/minpet/welcome.html

#### Linn, Jon

Technical Applications and Geographic Information Systems Unit West Virginia Department of Environmental Protection 10 McJunkin Rd, Nitro WV 25143, USA Phone: +1 304 759 0519 Fax: +1 304 759 0565 E-Mail: jlinn@crimson.osmre.gov Web: www.dep.state.wv.us/ito/tagis.html

## Lisker, Frank

Fachbereich 5 Geowissenschaften Universität Bremen Postfach 330440, D-28334 Bremen, Germany Phone: +49 421 218 3994 Fax: +49 421 218 3993 E-Mail: flisker@uni-bremen.de Web: www.palmod.unibremen.de/FB5/agpolar/englisch/index1.htm Liu, Shunsheng Changsha Institute of Geotectonics Academia Sinica Changsha 410013, Hunan Province, PR China Phone: +86 731 885 9165 Fax: +86 731 885 9137 E-Mail: liuss@ms.csig.ac.cn Web: www.csig.ac.cn

#### Lopez, Arturo

Departamento de Estudios del Ambiente Instituto nacional de investigaciones Nucleares Apartado Postal 18-1027, México D.F. 11801, Mexico E-Mail: arturo.lopez@mpi-hd.mpg.de Web: www.inin.mx

#### Lovering, John

School of Earth Sciences University of Melbourne 66A Molesworth Street, Kew Victoria 3101, Australia Phone: +61 3 9853 2868 Fax: +61 3 9853 3640 E-Mail: fitzlove@alphalink.com.au

#### Luft, Fernanda, F.

School of Earth Sciences University of Melbourne Victoria 3010, Australia Phone: +61 3 8344 9669 Fax: +61 3 8344 7761 E-Mail: ffe@pgrad.unimelb.edu.au

# Lutz, Timothy, M.

Department of Geology and Astronomy West Chester University West Chester PA 19383, USA Phone: +1 610 436 3498 E-Mail: tlutz@wcupa.edu Web: geology.wcupa.edu

#### Maksaev, Victor

Department of Geology University of Chile Casilla 13518, Correo 21, Santiago de Chile Phone: +56 2 678 4112 Fax: +56 2 696 3050 E-Mail: vmaksaev@tamarugo.cec.uchile.cl Web: www.cec.uchile.cl/~geologia

#### McInnes, Brent

Thermochronology Laboratory CSIRO Exploration & Mining 26 Dick Perry Avenue, Kensington WA Australia 6151, Australia Phone: +61 8 6436 8694 Fax: +61 8 6436 8586 E-Mail: brent.mcinnes@csiro.au Web: www.em.csiro.au

## Miller, Elizabeth, L.

Department of Geology and Environmental Sciences Stanford University Stanford, CA 94305-2115, USA Phone: +1 650 723 1149 E-Mail: miller@pangea.stanford.edu Web: pangea.Stanford.EDU/structure/projects/fission\_track.html

### Miller, Jack, A.

Department of Earth Sciences University of Cambridge Downing Street, Cambridge CB2 3EQ, UK Phone: +44 1223 337 184 Fax: +44 1223 333 450 E-Mail: jam2@esc.cam.ac.uk Web: www.esc.cam.ac.uk Mora, Andrès Institut für Geowissenschaften Universität Potsdam Postfach 60 15 53, D-14415 Potsdam, Germany Phone: +49 331 977 5880 Fax: +49 331 977 55700 E-Mail: mora@geo.uni-potsdam.de

#### Mora, Jorge

Laboratorios de Termocronología y Geomatemáticas Facultad de Ingeniería Universidad Central de Venezuela Caracas, Venezuela E-Mail: jImoram@lycos.com

## Murphy, John, M.

3901 Goodell Lane Ft. Collins, CO 80526-2924, USA Phone: +1 970 282 9570 E-Mail: zutfripple@cs.com

## Murtazaev, Khatam

General Physics Sub-Faculty Khujand State University 735700 Khujand, Tajikistan Phone: +992 3422 53212 23775 E-Mail: murtaza@hotam.khj.td.silk.org, murtaza@khj.tajik.net

## Naeser, Chuck

U.S. Geological Survey 926a National Center Reston, VA 20192, USA Phone: +1 703 648 6964 Fax: +1 703 648 6953 E-Mail: cnaeser@usgs.gov Web: greenwood.cr.usgs.gov/capabilities/gronemtrac/geochron/fissio n/contacts.html

#### Naeser, Nancy

U.S. Geological Survey 926a National Center Reston, VA 20192, USA Phone: +1 703 648 5328 Fax: +1 703 648 6953 E-Mail: nnaeser@usgs.gov Web: greenwood.cr.usgs.gov/capabilities/gronemtrac/geochron/fissio

## Neumann, Reinhard

n/contacts.html

Materials Research Gesellschaft für Schwerionenforschung Planckstrasse 1, D-64291 Darmstadt, Germany Phone: +49 6159 71 2172 Fax: +49 6159 71 3266 E-Mail: r.neumann@gsi.de Web: www-wnt.gsi.de/mr

## Nishimura, Susumu

Department of Geology and Mineralogy Division of Earth and Planetary Sciences Graduate School of Science Kyoto University Sakyo-ku, Kyoto 606-8502, Japan Phone: +81 75 753 4150 Fax: +81 75 753 4189 Web: terra.kueps.kyoto-u.ac.jp/index\_eg.html

#### Oddone, Massimo

Dipartimento di Chimica Generale Università degli Studi di Pavia Viale Taramelli 12, 27100 Pavia, Italy Phone: +39 382 507 334 Fax: +39 382 528 544 E-Mail: oddone@ipv36.unipv.it Web: chifis1.unipv.it/genchem.htm

#### Olesch, Martin

Fachbereich 5 Geowissenschaften Universität Bremen Postfach 330440, D-28334 Bremen, Germany Phone: +49 421 218 3940 Fax: +49 421 218 3993 E-Mail: olesch@geopol.uni-bremen.de Web:

www.palmod.unibremen.de/FB5/agpolar/englisch/index1.htm

## Omar, Gomaa, I.

Department of Earth and Environmental Science University of Pennsylvania 240 South 33rd Street Philadelphia, PA 19104-6316, USA Phone: +1 215 898 6908 Fax: +1 215 898 0964 E-Mail: gomar@sas.upenn.edu Web: www.sas.upenn.edu/earth/omar.html

## Ono, Masako

Department of Earth and Planetary Materials Science Faculty of Science Hokkaido University Kyoyo-chigaku N17-W8 Kita-ku, Sapporo 060, Japan Phone: +81 11 716 2111 ext. 5309 Fax: +81 11 736 3290 E-Mail: mo@epms.hokudai.ac.jp Web: www.ep.sci.hokudai.ac.jp/epms/index\_e.htm

## Osadetz, Kirk

Energy and Environment Geological Survey of Canada 3303-33rd St. NW Calgary, Alberta T2L 2A7, Canada Phone: +1 403 292 7022 Fax: +1 403 292 7159 E-Mail: kosadetz@nrcan.gc.ca Web: www.nrcan.gc.ca/gsc/calgary/index\_e.html

## O'Sullivan, Paul

Apatite to Zircon Inc. 1521 Pine Cone Road Moscow, ID 83843, USA Phone: +1 208 892 1493 Fax: +1 208 875 8881 E-Mail: OSullivan@apatite.com Web: http://www.apatite.com

### Pagel, Maurice

Laboratoire de Géochimie Faculté des sciences d'Orsay Université Paris Sud XI Bâtiment 504, F-91405 Orsay-Cedex, France Phone: +33 1 6915 6755 Fax: +33 1 6815 4882 E-Mail: pagel@geol.u-psud.fr Web: geosciences.geol.u-psud.fr/recherch/weblab/pagel.html

#### Pan, Yun

Department of Geological Sciences State University of New York at Albany DEAS - ES351, Albany NY 12222, USA E-Mail: ypan@spsu.edu;ynatl@bellsouth.net

## Parand, Simon

Geological Survey of Iran Meraj Blvd. Azadi Sq., P.O. Box 13185-1494 Tehran, Iran Fax: +98 21 600 9338 Web: www.gsi-iran.org Parra, Mauricio Institut für Geowissenschaften Universität Potsdam Postfach 60 15 53, D-14415 Potsdam, Germany Phone: +49 331 977 5880 Fax: +49 331 977 5700 E-Mail: mauricio@geo.uni-potsdam.de

#### Patel, Ramesh, C.

Department of Geophysics Kurukshetra University Kurukshetra 138 119, India Phone: +91 1744 238056 Fax: +91 1744 238277/238035 E-Mail: patelramesh\_chandra@rediffmail.com

#### Perelygin, Vladimir, P.

Flerov Laboratory of Nuclear Reactions Joint Institute for Nuclear Research Post Office Box 79, 101 000 Moscow, Russia Phone: +7 96 216 2102 Fax: +7 96 216 5955 E-Mail: pergam@cv.jinr.ru Web: sungraph.jinr.dubna.su/cap/index.html

## Petford, Nick

Centre for Earth and Environmental Science Research School of Earth Sciences and Geography Kingston University Penrhyn Road Kingston on Thames, Surrey KT1 2EE, UK Phone: +44 208 547 7518 Fax: +44 208 547 7497 E-Mail: n.pet@kingston.ac.uk Web: www.king.ac.uk/geog

### Poupeau, Gerard, R.

## Price, Buford, P.

Department of Physics University of California Berkeley, CA 94720, USA Phone: +1 510 642 4982 Fax: +1 510 643 8497 E-Mail: bprice@uclink4.berkeley.edu, price@lbl.qov Web: physics.berkeley.edu/research/price

#### Qvale, Henning

Institute for Energy Technology P.O. Box 40, N 2007 Kjeller, Norway Phone: +47 6380 6122 Fax: +47 6381 5553 E-Mail: hq@varney.ife.no Web: www.ife.no/english

## Raab, Matthias

School of Earth Sciences University of Melbourne Parkville, Victoria 3010, Australia Phone: +61 3 8344 9868, +61 3 8344 7675 Fax: +61 3 8344 7761 E-Mail: mraab@unimelb.edu.au

## Rahn, Meinert

HSK 5232 Villigen-HSK, Switzerland Phone: +41 56 310 39 03 Fax: +41 56 310 49 03 E-Mail: rahn@hsk.psi.ch

#### Ratschbacher, Lothar

Tektonophysik - Institut für Geologie Technische Universität Bergakademie Freiberg Bernhard-von-Cottastrasse 2, D-09596 Freiberg, Germany Phone: +49 3731 393 758 Fax: +49 3731 393 295 E-Mail: lothar@geo.tu-freiberg.de Web: www.geo.tu-freiberg.de/tektono

#### Raza, Asaf

School of Earth Sciences University of Melbourne Parkville, Victoria 3010, Australia Phone: +61 3 8344 9977, +61 3 8344 7675 Fax: +61 3 8344 7761 E-Mail: asaf@unimelb.edu.au Web: www.earthsci.unimelb.edu.au

#### Redfield, Thomas, F.

Technical Exploration Services Norges Geologiske Undersøkelse Leiv Eirikssons vei 39, 7491 Trondheim, Norway Fax: +47 73921620 E-Mail: tim.redfield@ngu.no

## Reinecker, John

Institut für Geologie und Paläontologie Eberhard Karls Universität Tübingen Sigwartstrasse 10, D-72076 Tübingen, Germany Fax: +49 7071 5059 E-Mail: john.reinecker@gpi.uni-karlsruhe.de Web: www.uni-tuebingen.de/geo/gpi

#### **Reiners**, Peter

Department of Geology and Geophysics Yale University P.O. Box 208109 New Haven, CT 06520-8109, USA Phone: +1 203 432 3761 E-Mail: peter.reiners@yale.edu Web: www.geology.yale.edu/~reiners/

#### Riley, Brook, C.D.

Department of Geological Sciences Cal Poly University Pomona 3801 West Temple Avenue Pomona, California 91768, USA Phone: +1 909 869 3459 E-Mail: bcdr7@yahoo.com, bcriley@csupomona.edu

## Ring, Uwe

Department of Geological Sciences University of Canterbury Private Bag 4800, Christchurch 8004, New Zealand Phone: +64 3 364-2987 ext. 7723 Fax: +64 3 364-2769 E-Mail: uwe.ring@canterbury.ac.nz Web: www.geol.canterbury.ac.nz

#### Roden-Tice, Mary, K.

Center for Earth and Environmental Science State University of New York Plattsburgh 101 Broad Street Plattsburgh, NY 12901, USA Phone: +1 518 564 2019 Fax: +1 518 564 3152 E-Mail: mary.rodentice@plattsburgh.edu Web: faculty.plattsburgh.edu/mary.rodentice

#### Ruiz, Geoffrey

Isotope Geochemistry Faculty of Earth and Life Sciences Vrije University Amsterdam De Boelelaan 1085, 1081 HV AMSTERDAM, The Netherlands Phone: +31 20 44 47 326, Mob:+31 6 11 02 81 54 Fax: +31 20 64 62 457 , ruig@geo.vu.nl

#### Sahu, Himansu

School of Earth Sciences University of Melbourne Victoria 3010, Australia Phone: +61 3 8344 7673 Fax: +61 3 8344 7761 E-Mail: hsahu@pgrad.unimelb.edu.au

#### Sandhu, Amanjit

Environmental Ściences Physical Sciences Division University of Toronto Scarborough Campus 1265 Military Trail, Scarborough ON M1C 1A4, Canada Fax: +1 416 287 7279 E-Mail: amanjitsandhu@usa.net Web: glacier.scar.utoronto.ca

## Schäfer, Thorsten

Fachbereich 5 Geowissenschaften Universität Bremen Postfach 330440, D-28334 Bremen, Germany Phone: +49 421 218 3994 Fax: +49 421 218 3993 E-Mail: postgrad@geopol.unibremen.de Web: www.palmod.unibremen.de/FB5/agpolar/englisch/index1.htm

#### Schuller, Volker

Institut für Geologie und Paläontologie Eberhard Karls Universität Tübingen Sigwartstrasse 10, D-72076 Tübingen, Germany Phone: +49 7071 2975 240 Fax: +49 7071 5059 E-Mail: volker.schuller@uni-tuebingen.de Web: www.uni-tuebingen.de/geo/gpi/agfrisch/mitarbeiter/schuller

#### Schwab, Martina

Institut für Geologie und Paläontologie Eberhard Karls Universität Tübingen Sigwartstrasse 10, D-72076 Tübingen, Germany Phone: +49 7071 2974 703 Fax: +49 7071 5059 E-Mail: martina.schwab@uni-tuebingen.de Web: www.uni-tuebingen.de/geo/gpi/index.html

#### Schwabe, Erika

Heroy Geology Laboratory Department of Earth Sciences Syracuse University Syracuse, New York 13244-1070, USA Phone: +1 315 443 3828 Fax: +1 315 443 3363 E-Mail: eschwabe@syr.edu Web: su-thermochronology.syr.edu

#### Sekyra, Gerd

Institut für Geologie und Paläontologie Universitat Salzburg Hellbrunnerstraße 34/III, A-5020 Salzburg, Austria Phone: +43 662 8044 5437 Fax: +43 662 8044 621 Web: www.sbg.ac.at/gew

#### Seiler, Christian

School of Earth Sciences University of Melbourne Parkville, Victoria 3010, Australia Phone: +61 3 8344 4996 Fax: +61 3 8344 7761 E-Mail: c.seiler@pgrad.unimelb.edu.au

#### Sélo, Madeleine

Laboratoire de Minéralogie Museum d'Histoire Naturelle 61 Rue Buffon, 75231 Paris Cedex 05, France Fax: +33 1 40 79 35 24 E-Mail: selo@mnhn.fr Web: www.mnhn.fr/mnhn/mineralogie

#### Seward, Diane

Institute of Geology, Department of Earth Sciences ETH-Zentrum Sonneggstrasse 5, 8092 Zürich, Switzerland Phone: +41 1 632 3675 Fax: +41 1 632 1080 E-Mail: diane.seward@erdw.ethz.ch, diane@erdw.ethz.ch Web: www.erdw.ethz.ch

#### Shi, Xiaobin

Key Laboratory of Marginal Sea Geology South China Sea Institute of Oceanology Chinese Academy of Sciences Guangzhou 510301, China Phone: +86 20 89023153 E-Mail: xbshi@scsio.ac.cn Web: http://www.scsio.ac.cn

#### Siddall, Ruth

Fission Track Research Group, Department of Earth Sciences University College London London WC1E-6B, UK Phone: +44 20 7679 2758 Fax: +44 20 7813 2802 E-Mail: r.siddall@ucl.ac.uk Web: www.ucl.ac.uk/geolsci/research/fissiontrack/Personal/Ruthpag e/RS.html

#### Sobel, Edward

Institut für Geowissenschaften Universität Potsdam Postfach 60 15 53, D-14415 Potsdam, Germany Phone: +49 331 977 5585 Fax: +49 331 977 5700 E-Mail: sobel@rz.uni-potsdam.de Web: www.geo.uni-potsdam.de/Personal/personal.html

#### Sohrabi, Mehdi

Radiation Protection Department Atomic Energy Organization of Iran P.O. Box 14155-4494, Tehran, Iran Fax: +98 21 690 6095 E-Mail: aeoicom@dci.iran.com

## Spencer, Steven

School of Earth Sciences University of Melbourne Victoria 3010, Australia Phone: +61 3 8344 9669 Fax: +61 3 8344 7761 E-Mail: s.spencer2@pgrad.unimelb.edu.au

#### Spiegel, Cornelia

Institut für Geowissenschaften Eberhard Karls Universität Tübingen Sigwartstrasse 10, D-72076 Tübingen, Germany Phone: +49 7071 2972494 Fax: +49 7071 5059 E-Mail: cornelia.spiegel@uni-tuebingen.de Spikings, Richard Section des Sciences de la Terre University of Geneva 13 Rue des Maraichers, 1211 Geneva 4, Switzerland Phone: +41 22 379 6826, Lab: +41 22 379 6639 Fax: +41 22 379 3210 E-Mail: Richard.Spikings@terre.unige.ch

#### Stapel, Gerco

Department of Isotope Geochemistry Faculty of Earth sciences Vrije Universiteit Amsterdam De Boelelaan 1085, Amsterdam 1081 HV, The Netherlands Fax: +31 20 646 2457 E-Mail: stag@geo.vu.nl Web: www.geo.vu.nl/~isotopen

## Steckler, Michael, S.

Lamont-Doherty Earth Observatory Columbia University Palisades, NY 10964, USA Phone: +1 914 365 8479 Fax: +1 914 365 8179 E-Mail: steckler@ldeo.columbia.edu Web: www.ldeo.columbia.edu

## Stiberg, Jan-Petter

Fission and Vision AS Ammerudgrenda 271, 0960 Oslo, Norway E-Mail: fission@online.no

## Stockli, Daniel

Department of Geology University of Kansas 1475 Jayhawk Boulevard 120 Lindley Hall Lawrence, KS 66045-7613, USA Phone: Lindley office:+1 785 864 4995, IGL office:+1 785 864 7714, Lab:+1 785 864 7709 Fax: +1 785 864 5276 E-Mail: stockli@ku.edu Web: http://www.people.ku.edu/~stockli/

## Stockmal, Glen, S.

Institute of Sedimentography and Petroleum Geology Geological Survey of Canada 3303-33rd Street, N.W. Calgary Alberta T2L 2A7, Canada Phone: +1 403 292 7173 Fax: +1 403 292 5377 E-Mail: gstockma@nrcan.gc.ca Web: www.nrcan.gc.ca/gsc/calgary/staff/stockmal e.html

#### Stone, John

Department of Geological Sciences University of Washington P.O. Box 351310, Seattle Washington 98195-1310, USA Phone: +1 206 685 9514 Fax: +1 206 543 3836 E-Mail: stone@geology.washington.edu Web: www.geology.washington.edu/faculty/stone-bio.html

#### Stump, Edmund

Department of Geological Sciences Arizona State University Box 871404 Tempe, AZ 85287-1404, USA Phone: +1 480 965 3971 Fax: +1 480 965 8102 E-Mail: ed.stump@asu.edu Web: geology.asu.edu/people/faculty/stump/index.html

#### Stüwe, Kurt

Institut für Geologie und Paläontologie Karl-Franzens Universität Heinrichstraße 26, A-8010 Graz, Austria Phone: +43 316 380 5682 Fax: +43 316 380 9870 E-Mail: kurt.stuewe@kfunigraz.ac.at Web: wegener.kfunigraz.ac.at

## Summerfield, Michael, A.

Department of Geography School of Earth Sciences University of Edinburgh Edinburgh EH8 9XP, UK Phone: +44 131 650 2519 Fax: +44 131 650 2524 E-Mail: mas@geo.ed.ac.uk Web: www.geo.ed.ac.uk/home/geoghome.html

#### Suzuki, Tatsuo

Department of Geology Faculty of Education Kagoshima University Korimoto 1-20-6, Kagoshima 890-0065, Japan Phone: +81 99 285 7807 Fax: +81 99 285 7807 E-Mail: t\_suzuki@edu.kagoshima-u.ac.jp Web: www.edu.kagoshima-u.ac.jp/english.html

#### Tagami, Takahiro

Department of Geology and Mineralogy Kyoto University Kitashirakawa Oiwakecho Sakyo-ku, Kyoto 606-8502, Japan Phone: +81 75 753 4153 Fax: +81 75 753 4189 E-Mail: tagami@terra.kueps.kyoto-u.ac.jp Web: www.kueps.kyoto-u.ac.jp/~tagami/

#### Talbot, James

K/T GeoServices Inc. P.O. Box 126 Argyle, TX 76226-0126, USA Phone: +1 214 403 6342 Fax: +1 940 387 9980 E-Mail: james@ktgeo.com Web: www.ktgeo.com/tCl.html

#### Tamanyu, Shiro

Geothermal Resources Research Group Institute for Geo-resources and Environment National Institute of Advanced Industrial Science and Technology Central 7 1-1-1 Higashi, Tsukuba 305-8567, Japan Phone: +81 298 61 3941, +81 298 61 3737 Fax: +81 298 61 3717 E-Mail: s.tamanyu@aist.go.jp Web: unit.aist.go.jp/georesenv/English/index.html

#### Tello Sáenz, Carlos, A.

Departamento de Raios Cósmicos e Cronologia Instituto de Fisica Gleb Wataghin Universidade Estadual de Campinas Caixa Postal 6165 CEP 13083-970 UNICAMP, Campinas SP, Brazil Phone: +55 19 788 5522 Fax: +55 19 788 5512 E-Mail: saenz@ifi.unicamp.br Web: www.ifi.unicamp.br/~turtelli/www0.htm

## Thiede, Rasmus

Institut für Geowissenschaften Universität Potsdam Postfach 60 15 53, D-14415 Potsdam, Germany Phone: +49 331 977 5784 Fax: +49 331 977 5700 E-Mail: thiede@geo.uni-potsdam.de Thomson, Stuart Department of Geology & Geophysics Yale University P.O. Box 208109 New Haven, CT 06520-8109, USA Phone: +1 203 432 1269 E-Mail: stuart.thomson@yale.edu Web: http://earth.geology.yale.edu/~snt6/

#### **Tingate**, Peter

National Centre For Petroleum Geology and Geophysics Thebarton Campus University of Adelaide Adelaide SA 5005, Australia Phone: +61 8 8303 4296 Fax: +61 8 8303 4345 E-Mail: peter.tingate@adelaide.edu.au Web: www.ncpgg.adelaide.edu.au/staff/peter tingate.htm

#### Trautmann, Christina

Materials Research Gesellschaft für Schwerionenforschung Planckstrasse 1, D-64291 Darmstadt, Germany Phone: +49 6159 71 2716 Fax: +49 6159 71 3266 E-Mail: c.trautmann@gsi.de Web: www-wnt.gsi.de/mr

## Van den haute, Peter

Geologisch Instituut Universiteit Gent Krijgslaan 281 (S8), B-9000 Gent, Belgium Phone: +32 9 264 4592 Fax: +32 9 264 4594 E-Mail: peter.vandenhaute@rug.ac.be Web: allserv.rug.ac.be/-fmees/mineralogy.htm

## Van der Beek, Peter

Labo de Géodynamique des Chaînes Alpines Observatoire des Sciences de l'Univers Grenoble Université Joseph Fourier Maison des Géosciences BP 53, 38041 Grenoble, France Phone: +33 476 514 062 Fax: +33 476 514 058 E-Mail: pvdbeek@ujf-grenoble.fr Web: www.univsavoie.fr/labos/lgca/perso/cherc/pvand/pvand.htm

Van der Wateren, F., M.

Department of Isotope Geochemistry Faculteit Aardwetenschappen Vrije Universiteit Amsterdam De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands Fax: +31 20 646 2457 Web: www.geo.vu.nl/~isotopen

## Vasconcelos, Paulo

Department of Earth Sciences University of Queensland St Lucia, Brisbane QLD 4072, Australia Phone: +61 7 3365 2297 Fax: +61 7 3365 1277 E-Mail: p.vasconcelos@earth.uq.edu.au;, paulo@bgc.org Web: www.earth.uq.edu.au

## Ventura, Barbara

Fachbereich 5 Geowissenschaften Universität Bremen Postfach 330440, D-28334 Bremen, Germany Phone: +49 421 218 3994 Fax: +49 421 218 3993 E-Mail: bventura@geopol.uni-bremen.de Web:

www.palmod.unibremen.de/FB5/agpolar/englisch/index1.htm

#### Vidal, Olivier

Laboratoire de Géodynamique des Chaînes Alpines Maison des Géosciences Université Joseph Fourier 1381 Rue de la Piscine, 38400 St-Martin d'Hères, France Phone: +33 4 76 53 59 24 Fax: +33 4 76 51 40 58 E-Mail: Olivier.Vidal@ujf-grenoble.fr Web: www.univsavoie.fr/labos/lgca/perso/cherc/ovid/ovid.htm

#### Viola, Giulio

Geological Survey of Norway Trondheim, 7491, Norway E-Mail: gulio.viola@ngu.no

## Wadatsumi, Kiyoshi

Department of Geosciences Osaka City University 3-3-138 Sugimoto Sumiyoshi-ku, Osaka 558, Japan Phone: +81 6 605 2593 Fax: +81 6 605 3071 E-Mail: h1602@ocugw.cc.osaka-cu.ac Web: www.sci.osaka-cu.ac.jp/geos/english/geoeng.html

#### Wagner, Günther, A.

Max-Planck-Institut für Kernphysik Postfach 103980, D-69029 Heidelberg, Germany Phone: +49 6221 516 289 Fax: +49 6221 516 633 E-Mail: g.wagner@mpi-hd.mpg.de Web: www.mpi-hd.mpg.de/english/index.html

## Walgenwitz, Frederic

CSTJF Elf Aquitaine Production Avenue Larribau, 64018 Pau Cedex, France E-Mail: frederic.walgenwitz@elf-p.fr

## Wallace, Wesley, K.

Department of Geology and Geophysics University of Alaska Fairbanks, AK 99775-5780, USA Phone: +1 907 474 5386 Fax: +1 907 474 5163 E-Mail: wes.wallace@gi.alaska.edu Web: www.gi.alaska.edu/tsrg/people/wallace

#### Wang, Shicheng

Institute of High Energy Physics Chinese Academy of Sciences P.O. Box 2732, Beijing 100080, P.R. China E-Mail: wangshch@ihepa.ac.cn Web: www.ihep.ac.cn/english/index.htm

#### Watanabe, Hiroaki

Department of Earth Sciences Kanazawa University Kanazawa 920-1192, Japan Fax: +81 76 264 5746 E-Mail: hiro741@kenroku.kanazawa-u.ac.jp Web: earth.s.kanazawau.ac.jp/chronology/EnglishPage/indexE.html

#### Watanabe, Koichiro

Department of Mining Faculty of Engineering Kyushu University 36 Hakozaki Higashi-ku, Fukuoka 812-8581, Japan Phone: +81 92 642 3634 E-Mail: wat@mine.kyushu-u.ac.jp Web: www.eng.kyushu-u.ac.jp/index-e.html

## Westgate, John, A.

Environmental Sciences Physical Sciences Division University of Toronto Scarborough Campus 1265 Military Trail, Scarborough ON M1C 1A4, Canada Phone: +1 416 287 7234 Fax: +1 416 287 7279 E-Mail: westgate@scar.utoronto.ca Web: glacier.scar.utoronto.ca

## Wiesinger, Maria

Institut für Geologie und Paläontologie Universität Salzburg Hellbrunner Straße 34, A-5020 Salzburg, Austria E-Mail: maria1.wiesinger@sbg.ac.at

### Willet, Sean, D.

Department of Geological Sciences University of Washington P.O. Box 351310, Seattle Washington 98195-1310, USA Phone: +1 206 543 8653 Fax: +1 206 543 3836 E-Mail: swillett@u.washington.edu Web: www.geology.washington.edu/~willett

#### Wölfler, Andreas

Institut für Geowissenschaften Eberhard Karls Universität Tübingen Sigwartstrasse 10, D-72076 Tübingen, Germany Phone: +49 7071 2974702 Fax: +49 7071 5059 E-Mail: andreas.woelfler@uni-tuebingen.de

## Wu, Qianhong

Changsha Institute of Geotectonics Academia Sinica Changsha 410013, Hunan Province, P.R. China Phone: +86 731 885 9165 Fax: +86 731 885 9137 E-Mail: wqh@ms.csig.ac.cn Web: www.csig.ac.cn

### Xu, Ganqing

Department of Earth Sciences University of Waikato Private Bag 3105, Hamilton 2001, New Zealand Phone: +64 7 838 4024 Fax: +64 7 856 0115 E-Mail: xu1@waikato.ac.nz Web: sci.waikato.ac.nz/cgi-bin/dynapage.pl?pagename=erth

#### Yamada, Ryuji

Nagaoka Institute of Snow and Ice Studies National Research Institute for Earth Science and Disaster Prevention 9400082 Suyoshi Nagaoka Niigata, Japan Phone: +81 258 35 7522 Fax: +81 258 35 0020 E-Mail: ryamada@nagaoka.bosai.go.jp Web: www.bosai.go.jp/ad/Eng/division.htm

## Yamashita, Tohru

Kyoto Fission-Track Co. Ltd. 44-4 Minamitajiri-cho Omiya Kita-ku, Kyoto 603-8832, Japan Phone: +81 75 493 0684 Fax: +81 75 493 0741 E-Mail: kyoto-ft@mb.neweb.ne.jp

## Yang, Tsanyao, F.

Department of Geology National Taiwan University 245 Chou-shan Road, Taipei 106-17, Taiwan Phone: +886 2 2369 5568 Fax: +886 2 2363 6095 E-Mail: tyyang@ccms.ntu.edu.tw Web: www.gl.ntu.edu.tw/english\_mode

#### Yegingil, Zehra

Physics Department Cukurova University P.O. Box 171, 01330 Adana, Turkey E-Mail: zehra@cu.edu.tr Web: www.cu.edu.tr/fakulteler/fef

#### Yoshioka, Tetsu

Kyoto Fission-Track Co. Ltd. 44-4 Minamitajiri-cho Omiya Kita-ku, Kyoto 603-8832, Japan Phone: +81 75 493 0684 Fax: +81 75 493 0741 E-Mail: kyoto-ft@mb.neweb.ne.jp

## Yuan, Wanming

Institute of High Energy Physics Chinese Academy of Sciences P.O. Box 2732, Beijing 100080, P.R. China E-Mail: yuanwm@ihepa.ac.cn Web: www.ihep.ac.cn/english/index.htm

### Zattin, Massimiliano

Dipartimento di Scienze Geologiche Universita di Bologna Via Zamboni 67, 40127 Bologna, Italy Phone: +39 51209 4579 Fax: +39 51209 4522 E-Mail: zattin@geomin.unibo.it Web: www.geomin.unibo.it

#### Zeitler, Peter, K.

Department of Earth and Environmental Sciences Lehigh University 31 Williams Drive Bethlehem, PA 18015, USA Phone: +1 215 758 3671 Fax: +1 215 758 3677 E-Mail: peter.zeitler@lehigh.edu Web: www.ees.lehigh.edu

## Zentilli, Marcos

Department of Earth Sciences Dalhousie University Halifax Nova Scotia, B3H 3J5, Canada Phone: +1 902 494 3873 Fax: +1 902 494 6889 E-Mail: marcos.zentilli@dal.ca Web: meguma.earthsciences.dal.ca/staff/zentilli/zentilli.htm

## Zhou, Zuyi

Department of Marine Geology and Geophysics Tongji University 1239 Siping Road, 200092 Shanghai, China Phone: +86 21 6598 2358 Fax: +86 21 6502 6278 E-Mail: zzydmg@tju.ihep.ac.cn Web: www.tongji.edu.cn/lib\_english/enindex.htm

#### Zuffa, Gian, G.

Dipartimento di Scienze Geologiche Universita di Bologna Via Zamboni 67, 40127 Bologna, Italy Phone: +39 51209 4540 Fax: +39 51209 4522 E-Mail: zuffa@geomin.unibo.it Web: www.geomin.unibo.it

#### Zwingmann, Horst

CSIRO Petroleum Curtin University PO Box 1130, Bentleigh, WA 6102, Australia Phone: +61 8 9266 4681 E-Mail: horst.zwingman@csiro.au

# Call for Contributions

The next issue of OnTrack is scheduled for early in 2007 and we are (always) looking for contributions. OnTrack welcomes contributions of virtually any kind, including:

- scientific articles,
- present PhD studies,
- recent PhD theses abstracts,
- job openings,
- news,
- descriptions of new lab techniques,
- descriptions of what you are doing in your research,
- reviews of useful products,
- ravings about what your lab is doing perfectly,
- meeting announcements,
- cartoons and gossip,
- ...

OnTrack includes a list of recent and forthcoming fission-track papers. If you know of a paper that was published recently or is in press and should appear in the list, please let the editor know so that it can be added to the list. Also, if you happen to change location due to a change in jobs or finishing off the thesis and graduating, please inform the editor. Thesis abstracts are particularly encouraged! OnTrack is also happy to print advertisements. Please contact the editor for advertising rates.

## Send contributions to:

Matthias Raab School of Earth Sciences University of Melbourne Parkville, Victoria 3010, Australia Phone: +61 3 8344 9868 /7675 Fax: +61 3 8344 7761 E-mail: mraab@unimelb.edu.au Web: www.earthsci.unimelb.edu.au


**Rex F. Galbraith**, *University College*, *London*, *UK* A volume in the **Interdisciplinary Statistics** series Series editors: Niels Keiding; Byron J.T. Morgan, and Peter Van der Heijden

## THE TOOLS FOR EXPLORING GEOLOGICAL HISTORY

Statistical analyses of the numbers, lengths, and orientations of fission tracks etched in minerals yield dating and thermal history information valuable in geological and geoscience applications, particularly in oil exploration. Fission tracks can be represented mathematically by a stochastic process of randomly oriented line segments in three dimensions, and this "line segment" model can describe and explain the essential statistical features of the data, providing a rigorous foundation for quantitative modelling and simulation studies.

**Statistics for Fission Track Analysis** explores the line segment model and its consequences for the analysis and interpretation of data. The author derives the equations for fission track data and the theoretical probability distributions for the number, orientation, and length measurements of the tracks. He sets out the theory of fission track dating and through numerical examples, presents methods for analyzing and interpreting fission track counts. Later chapters address statistical models for situations in which samples contain mixtures of fission track ages. These methods, along with observation features of the various measurements, are illustrated by real examples. Finally, the author brings together the theoretical and observation aspects to formulate a joint likelihood function of counts, lengths, and angles as a basis for parametric thermal history modelling.

Designed for broad accessibility, this is the first book to fully cover the statistical foundations of fission track analysis. Whether you work in a fission track lab, in archaeological, geological, or geochronological research, or in geological applications of statistics, you will find the background material and practical tools you need to optimize the use of fission track analysis in your work and to make further advances in the field.

## **FEATURES**

- Establishes the line segment model for fission track analysis from physical considerations and verifies empirically key aspects of the theory
- Includes detailed worked examples of data analysis obtained by both the "external detector" and "population" experimental methods
- Briefly discusses the statistical aspects of fission track annealing studies and the calculation of palaeotemperatures
- Contains clear explanations of the mathematical and statistical concepts employed

## CONTENTS

### INTRODUCTION

What are Fission Tracks? How are they Observed? Why are they Useful? Applications of Fission Track analysis Mathematical Representation of Fission Tracks Fission Track Dating and Provenance Studies Thermal Histories and Track Length Distributions Sampling by Plane Section Initial Formation of Tracks Shortening of Tracks by Heat Properties of Apatite **THE POISSON LINE SEGMENT MODEL** Joint Distribution of Length and Orientation

The Number of Tracks with a Given Attribute

The Expected Number of Tracks Intersecting a Plane

Track Density and Equivalent Isotropic Length Tracks Intersecting a Prismatic Face Effect of Non-Prismatic Face on Track Density Track Counts from a Dosimeter Glass Spatial and Temporal Variation Remarks TRACK COUNTS AND DENSITIES: FISSION

#### TRACK COUNTS AND DENSITIES: FISSION TRACK DATING

The Mathematical Basis of Fission Track Dating The External Detector Method Observed and Theoretical Track Densities A Short Digression Estimates of Fission Track Age Inspection of Single Grain Data Radial Plot of Single Grain Ages Chi-Square Age Homogeneity Test A Measure of Age Dispersion A Protocol for Data Analysis Dealing with Small Counts Practical Considerations Remarks Historical Note **THE POPULATION METHOD** Experimental Method and Data

Experimental Method and Data Theoretical and Observed Track Densities An Estimate of the Uranium Dispersion A Uranium Homogeneity Test Estimates of Fission Track Age Summarising and Inspecting the Data Age Homogeneity Test *contents continued on reverse* 

Taylor & Francis Group

Chapman & Hall/CRC

# **Contents continued STATISTICS FOR FISSION TRACK ANALYSIS**

A Measure of Dispersion of True Fission Track Ages Counts over Unequal Areas A Protocol for Data Analysis Discussion The Population-Subtraction Method Remarks **DISCRETE MIXTURES OF AGES** Maximum Likelihood Estimation of a Common Age **Discrete Mixture Models** Example: A Synthetic Mixture of Two Ages Example: Apatite Data from the Bengal Fan Maximum Likelihood Estimation Formulae How Many Ages to Fit? Example: Zircon Ages from Mount Tom Data from More than One Irradiation **CONTINUOUS MIXTURES OF AGES** Example: Otway Data from Victoria, Australia **General Approach** A Random Effects Model with Binomial Errors Maximum Likelihood Estimation Formulae A Random Effects Model with Normal Errors Examples Finite Mixtures of Random Effects Models A Minimum Age Model

Data and Statistical Model with Binomial Errors Maximum Likelihood Estimation Formulae A Minimum Age Model with Normal Errors Example: Apatite Data from China A Synthetic Mixture Revisited Grain Age Distributions Remarks **PROBABILITY DISTRIBUTIONS OF LENGTHS AND ANGLES** All Tracks Having the Same Length Each Track Having One of Two Lengths Several Different Lengths A General Isotropic Length Distribution A General Anisotropic Length Distribution Distributions on a Prismatic Face Horizontal Confined Track Lengths Some Explicit Formulae A Two-Component Mixture of Anisotropic Lengths

Quantitative Effects of Anisotropy Parametric Models for Length Against Angle OBSERVATIONAL FEATURES OF

## TRACK MEASUREMENTS

Horizontal Confined Tracks Length Bias The Loaded Dog Experiments Empirical Verification of Length Bias Fracture-Thickness Bias Orientation Bias Surface Proximity Bias Estimate of  $\mu$  from Horizontal Confined Tracks Projected Semi-Track Lengths and Angles Semi-Track Lengths and Angles

## FURTHER DEVELOPMENTS

Thermal History Parameters Combined Likelihood for Track Measurements Annealing Experiments Annealing Data Annealing Models Fitting Annealing Models Calculating the Length Distribution from a thermal history Inferring Times and Temperatures from Lengths Multi-Compositional Annealing Models APPENDIX Poisson Processes in One, Two, and Three Dimensions Notes on the Poisson Distribution Relation Between Binomial and Poisson Distribution Standard Errors and Confidence Intervals Components of Error Statistical Significance Tests and *p*-Values **Radial Plots** Histograms and "Probability Density" Plots Parametric Models and Likelihood Inference

Each chapter also contains Bibliographic Notes

## Please use this ORDER FORM, CALL or ORDER ONLINE at WWW.CRCPRESS.COM

STATISTICS FOR FISSION TRACK ANALYSIS ......Catalog no. C5335, ISBN: 1-58488-533-5 at \$79.95 / £44.99 each.

### Other titles of interest:

Country

HANDBOOK OF GEOPHYSICAL EXPLORATION AT SEA ..........Catalog no. 4252, ISBN: 0-8493-4252-X at \$319.95 / £183.00 each.

STANDARD METHODS OF GEOPHYSICAL FORMATION EVALUATION ......Catalog no. L1261, ISBN: 1-5667-0261-5 at \$99.95 / £57.99 each.

INTRODUCTION TO GEOPHYSICAL FORMATION EVALUATION ........Catalog no. L1262, ISBN: 1-5667-0262-3 at \$84.95 / £48.99 each.

Ordering Information: Orders must be prepaid or accompanied by a purchase order. Checks should be made payable to CRC Press. Please add the appropriate shipping and handling charge for each book ordered. All prices are subject to change without notice. <u>U.S./Canada</u>: All orders must be paid in U.S. dollars. TAX: As required by law, please add applicable state and local taxes on all merchandise delivered to CA, CT, FL, KY, MO, NY, and PA. For Canadian orders, please add GST. We will add tax on all credit card orders. **European Orders:** All orders must be paid in U.K. £. VAT will be added at the rate applicable. <u>Textbooks:</u> Special prices for course adopted textbooks may be available for certain titles. To review a book for class adoption, contact our Academic Sales Department or submit your textbook evaluation request online at www.crcpress.com/eval.htm <u>Satisfaction Guaranteed</u>: If the book supplied does not meet your expectations, it may be returned to us in a saleable condition within 30 days of receipt for a full refund.

CHIDDING AND HANDLING

SHIFFING AND HANDLING				
Region	Delivery Time	First Title	Additional Title	For priority
USA/Canada	3-5 Days	\$5.99	\$1.99	mail services,
South America	7-14 Days	\$9.99	\$3.99	please contact
Europe	3-5 Days	£2.99	£0.99	CRC PRESS
Rest of World	7-21 Days	£4.99	£2.99	office.
Signature and Telephone Num	ber required on all	orders	Exp. Date	Aonth Year
Signature			PO#	
Telephone				
If you would like to receive information from us by e-mail, please provide your e-mail address below.				

E-Mail Address .

