A service of the U.S. National Library of Medicine ? and the National In itutes of Health [Sign In] [Register] All Databases PubMed Nucleotide Protein Genome Structure **OMIMO PMC** Journals Books Search PubMed for Go Clear Preview/Index Limits History Clipboard **Details** Display **AbstractPlus** 20 Sort By Send to Show Review: 0 Links **1:** <u>J Endocrinol Invest.</u> 2003 Jul;26(7):679-82. Coexistence of a parathyroid adenoma and parathyroid cyst Related Links causing primary hyperparathyroidism. A parathyroid cyst with adenoma on thallium-201/technetium-99m Ardito G, Fadda G, Danese D, Modugno P, Giordano A, subtraction imagif@n Nucl Med. 1987] Revelli L, Ardito F, Pontecorvi A. [A parathyroid cyst and an Institute of Surgical Semeiothic, Catholic University of Sacred Heart, adenoma associated with primary Rome, Italy. hyperparathyroidism: report of a [Hinyokika Kiyo. 1990] The association of a functional parathyroid cyst with a parathyroid adenoma is an uncommon finding. In this report Parathyroid cyst and hyperparathyroidisminiz Megklarges1 we describe the clinical history of a 60-yr-old man, presenting with the following findings: hypercalcemia (18.9 [Giant intrathyroidal parathyroid mg/dl), elevated serum parathormone levels (1320 pg/dl), cyst with hyperparathyroidism: a hypercalciuria (228 mg/dl), and hyperphosphaturia (155 case report] [Ann Ital Chir. 2005] mg/dl). Neck ultrasound, magnetic resonance imaging (MRI) [Cystic parathyroid adenoma. A and 99Tc Sestamibi scintigraphy led to the identification of a rare cause of primary left parathyroid adenoma, located at the lower pole of the hyperparathyroidisgeskr Laeger. 2002] left thyroid gland lobe, associated with a parathyroid cyst, located at the upper extremity of the same thyroid lobe. See all Related Articles... Parathyroidectomy was performed and the histological examination confirmed the diagnosis of a parathyroid adenoma with aspects of cystic degeneration and an upper parathyroid cyst. Analysis of the crystal clear intracystic fluid showed elevated parathyroid hormone (PTH) levels (137.000 pg/ml). The patient is normocalcemic at 2 yr after surgery without signs of recurrent parathyroid enlargements. Aetiology, diagnosis and management of parathyroid cyst will be discussed. PMID: 14594122 [PubMed - indexed for MEDLINE] Display AbstractPlus Show 20 Sort By Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer